

# SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی




عضویت در خبرنامه



فیلم های آموزشی

## کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



مباحث پیشرفته یادگیری عمیق؛  
شبکه های توجه گرافی  
(Graph Attention Networks)



کارگاه آنلاین آموزش استفاده از  
وب آو ساینس



کارگاه آنلاین مقاله روزمره انگلیسی

## Original Article

## Suicidal mothers

Salvatore Gentile <sup>a,\*</sup><sup>a</sup> Department of Mental Health, Piazza Galdi, Italy.

## KEY WORDS

Depression  
Pregnancy  
Suicide

**Abstract:**

**Background:** Epidemiological research has demonstrated that suicidal ideation is a relatively frequent complication of pregnancy in both developed and developing countries. Hence, the aims of this study are: to assess whether or not pregnancy may be considered a period highly susceptible to suicidal acts; to recognize potential contributing factors to suicidal behaviors; to describe the repercussions of suicide attempts on maternal, fetal, and neonatal outcome; to identify a typical profile of women at high risk of suicide during pregnancy.

**Methods:** Medical literature information published in any language since 1950 was identified using MEDLINE/PubMed, Scopus, and Google Scholar databases. Search terms were: "pregnancy", (antenatal) "depression", "suicide". Searches were last updated on 28 September 2010. Forty-six articles assessing the suicidal risk during pregnancy and obstetrical outcome of pregnancies complicated by suicide attempts were analyzed, without methodological limitations.

**Results:** Worldwide, frequency of suicidal attempts and the rate of death by suicidal acts are low. Although this clinical event is rare, the consequences of a suicidal attempt are medically and psychologically devastating for the mother-infant pair. We also found that common behaviors exist in women at high risk for suicide during pregnancy. Review data indeed suggest that a characteristic profile can prenatally identify those at highest risk for gestational suicide attempts.

**Conclusions:** Social and health organizations should make all possible efforts to identify women at high suicidal risk, in order to establish specific programs to prevent this tragic event. The available data informs health policy makers with a typical profile to screen women at high risk of suicide during pregnancy. Those women who have a current or past history of psychiatric disorders, are young, unmarried, unemployed, have incurred an unplanned pregnancy (eventually terminated with an induced abortion), are addicted to illicit drugs and/or alcohol, lack effective psychosocial support, have suffered from episodes of sexual or physical violence are particularly vulnerable.

Accepted 2010-

© 2011 KUMS, All right reserved

**\*Corresponding Author at:**

Salvatore Gentile, MD, ASL Salerno (Italy) - Department of Mental Health, Mental Health Center n. 63 Cava de' Tirreni, Operative Unit n. 63 Cava de' Tirreni - Amalfi Coast, Piazza Galdi, 84013 Cava de' Tirreni (Salerno), tel. +39 089 4455439, fax.+39 089 4455440, e-mail: [mailto:salvatore\\_gentile@alice.it](mailto:mailto:salvatore_gentile@alice.it) (Salvatore Gentile).

© 2011 KUMS, All right reserved

**Introduction**

A World Health Organization report has named depression as the greatest disease burden for women worldwide. <sup>1</sup> In particular, childbearing years are a time of in-

creased vulnerability to the onset or recurrence of major depressive disorder, thus placing young women at risk of suffering from severe affective impairment during pregnancy. <sup>2,3,4</sup> Indeed, antenatal depression is a significant public health problem. Information drawn from scientific literature under-

lines that, in developed countries, between 3% and 16% of pregnant women fulfill the diagnostic criteria for unipolar major depression.<sup>5,6</sup> In specific populations, such as marginalized minority groups and unmarried teenagers, the rate of clinically relevant mood symptoms in pregnancy may be as high as 51%.<sup>7</sup> In developing countries, the prevalence of antenatal depression is estimated at 33%.<sup>8</sup>

It is well-known that antenatal depression adversely impacts on several obstetrical, fetal, and infantile outcomes. In fact, untreated maternal mood symptoms have been associated with an increase in the rate of pregnancy complications (such as pre-eclampsia, premature delivery, impaired fetoplacental function, and low fetal growth) and perinatal problems.<sup>9,10,11,12,13,14</sup>

Maternal depression also induces significant effects on neonatal physiology. Elevated cortisol and norepinephrine levels, lower dopamine levels, and greater relative right frontal EEG asymmetry have all been described in newborns of mothers suffering from depression during pregnancy.<sup>15,16</sup> Findings from a preliminary, recent study<sup>17</sup> have also suggested that there is an elevated risk of lower birth weight among infants born to adolescent mothers who report depressive symptoms complicated by self-harm ideation and behavior compared to those born to adolescents who report depressive features without suicidal symptoms.

The detrimental impact of untreated maternal depression on the infant neurocognitive and psychological development is also well-known. Many untreated women continue to be depressed in the postnatal period and are at risk of impaired interaction with their infant at a time when the child is in a particularly sensitive developmental period<sup>18,19,20,21,22</sup>, and this may lead to poorer cognitive functioning and compromised social adaptation during childhood, adolescence, and young adulthood.<sup>23,24,25</sup>

Prevalently, however, depression is characterized by a dramatic increase in the risk of suicide or serious suicide attempts, since more than 90% of suicide victims have a diagnosable mental illness and approximately 60% of all suicides occur in persons with mood disorders.<sup>26</sup>

Suicide during pregnancy is included among the indirect causes of *maternal death*. The World Health Organization defines *maternal deaths* as all deaths occurring any time during pregnancy and up to 42 days after parturition.<sup>27</sup> Indirect causes of maternal deaths include all those conditions not directly related to obstetric cause but worsened by the pregnant status, such as complications due to cardiac disease and mental disorders. When all maternal deaths within one year after delivery are considered, suicide is one of the four leading causes of maternal deaths overall (together with throm-

boembolisms, obesity, and cardiac events).<sup>28,29</sup> For this reason, maternal mortality associated with psychiatric illness should remain a focus of high clinical interest in the developed countries. In these countries, in fact, the rate of maternal deaths due to other conditions, such as poor antenatal care and malnutrition, have progressively decreased during the last century.<sup>30</sup> Moreover, the epidemiological studies have been mainly focused on the increased risk of suicidal attempts during the puerperal period, since suicides account for up to 20% of postpartum deaths.<sup>31</sup>

Hence, the primary aim of this study is to assess whether or not pregnancy may be considered a period highly susceptible to suicidal events. Other scopes of this review are: to recognize potential contributing factors to suicidal behaviors; to describe the repercussions of suicide attempts on maternal, fetal, and neonatal outcomes; to identify a typical profile of women at high risk of suicide during pregnancy, in order to establish effective strategies of prevention.

## Methods

Medical literature information published in any language since 1950 was identified using MEDLINE/PubMed, Scopus, and Google Scholar databases. Additional references were identified from the reference lists of published articles. Search terms were: “pregnancy”, (antenatal) “depression”, “suicide”. Searches were last updated on 28 September 2010. Selected on the basis of their abstract or the full-text article when the abstract was unavailable, all articles assessing the suicidal risk during pregnancy and obstetrical outcome of pregnancies complicated by suicide attempts were acquired and analyzed, without methodological limitations. Forty-six articles met the wide inclusion criteria. The Author was the only reviewer who performed both data selection and extraction.

## Results

### Prevalence of suicidal ideation in pregnancy

In the Western world, recent epidemiological researches have demonstrated that suicidal ideation may be detectable in a percentage ranging from 13.1% to 33.0% of pregnant women.<sup>32,33</sup> Symptoms frequently associated with suicidal ideation include depressed mood, lack of concentration, anxiety, preoccupations, obsessive compulsive symptoms, tiredness, worry concerning bodily functions, and compulsion.<sup>34</sup>

In developing countries, too, nearly 14% of pregnant women admit to thoughts of self-harm.<sup>8</sup> In these regions of the world, the occurrence of suicidal ideation during the gestational period is a relevant concern especially among adoles-

cents, since it has been reported in up to 17% of young mothers.<sup>35</sup>

#### A. How many mothers die by suicide?

##### Europe

A Danish register/case audit study based on data from the registers of the Danish Medical Health Board, death certificates, and hospital records covering the period 2002-2006 identified 26 cases of maternal death, leading to a maternal mortality ratio of 8.0/100,000 live births.<sup>36</sup> No cases of death due to suicide were identified. Similar results derived from data extrapolated from three national Finnish health registers<sup>37</sup> and two Swedish national registers.<sup>38</sup>

Only five cases of suicidal deaths during pregnancy have been described in the United Kingdom report *Why mothers die 2000-2002*.<sup>39</sup> In three of these cases, the women had been diagnosed with severe form of psychiatric disorders, such as major depression (two cases) or schizophrenia (one case).

##### North America

An epidemiological U.S. study performed on a large State of California database showed that the cumulative incidence of suicide attempts during pregnancy was four per 10,000 pregnancies.<sup>40</sup> The contribution of several injuries to maternal mortality was assessed through a review of New York City medical examiner records of 2331 women 15-44 years of age who died of injuries during 1987-91.<sup>41</sup> The largest proportion of injury-related deaths were homicides (63%), whereas suicide accounted for only 11 cases. To further determine the prevalence of suicide during pregnancy, the autopsy reports of all female residents of the same city, 10-44 years old, who committed suicide from 1990 to 1993 were assessed for pregnancy.<sup>42</sup> The standardized mortality ratio for suicide during pregnancy was 0.33; that is, the number of suicides of pregnant women was only one-third of that expected.

A retrospective study of abortion and maternal deaths in Minnesota during the period 1950-1965 including data on more than 1,400,000 women reported only four cases of suicides that occurred during the gestational period<sup>43</sup>, whereas the Maternity Mortality Collaborative (a U.S. program focused to investigate incidence, risk factors, and causes of maternal deaths in 19 regional areas) identified 13 cases of suicide among 712 deaths of women during pregnancy that occurred between 1980 and 1985.<sup>44</sup>

In the year 2000, a collaborative effort involving World Health Organization (WHO), UNICEF, and UNFPA estimated 660 maternal deaths in the United States. This av-

erages out at 11 maternal deaths per 100,000 live births reported.<sup>45</sup> The most common causes of maternal death varied somewhat from region to region. Several medical and obstetric complications, but not suicidal attempts, were identified as significant contributors to maternal mortality in these populations. Moreover, only five cases of maternal suicidal deaths were identified through a follow-up investigation of 41 injury-related maternal deaths identified from 1992 to 1994 in North Carolina.<sup>46</sup> Canadian data also excluded a significant risk of suicidal death during pregnancy.<sup>47</sup>

##### Asia

An epidemiological study<sup>48</sup> looked at maternal deaths in Singapore from 2000 to 2004, by linking coronial cases of female suicide in the reproductive age group 15-45 years, with the birth registration database, to identify both early and late maternal deaths. Among the 589 cases of identified maternal death due to suicide, none of these occurred during pregnancy.

##### Africa

Similar results emerged after analyzing data from African countries. An estimation of maternal mortality rates for 1982-83 in Zambia led to identify just one case of suicide (induced by an illegitimate pregnancy).<sup>49</sup> Of 175 maternal deaths reported in two district of Zimbabwe over two years (1989-1990), four were listed as suicide owing to unwanted pregnancy.<sup>50</sup>

B. Factors contributing to increase the suicidal risk in pregnancy are summarized in [Table 1](#).

**Table 1. Contributing factors to suicidal behaviors in pregnancy\***

Developing countries	Developed countries
Intimate partner violence, both physical and sexual	Intimate partner violence, both physical and sexual
Unmarried status	Unmarried status
Teen age #	Teen age #
Pregnancy outside marriage €	Unplanned pregnancy
Lack of effective familial and/or social government institution supports	Lack of effective familial and/or social government institution supports
Conception of a female fetus, especially in those traditional societies where a strong familial preference exists towards the birth of male infants	Previous or present history of any psychiatric disorder
Poor mother-in-law support	Street drug/alcohol abuse
	Unemployment
	Induced abortion

\* These findings are not in a ranking numeric order

# Age ranges are not available

### Developed countries

The rate of suicidal ideation in pregnancy is significantly associated with psychiatric disorder and, especially, with current major depressive episode and comorbid anxiety disorder and/or substance abuse.<sup>8,32</sup> Psychosocial factors which may also contribute to increase the rate of maternal suicide attempts during pregnancy are teen age, unplanned pregnancy, unmarried status or recent divorce, unemployment, and difficult access to safe abortion service.<sup>52,53</sup> Moreover, hospital-based, cohort studies, and literature reviews have suggested that intimate partner violence, previous experience of sexual assaults, interpersonal conflicts also represent specific factors associated with an increased suicidal risk.<sup>54,55,56</sup> Experiencing intimate partner violence may increase the risk of suicidal behaviors by enhancing the risk of impulsive responses and facilitating the onset of posttraumatic stress disorder, since both situations are intrinsically associated with an increased suicidal risk.<sup>57,58</sup> In addition, both severe emotional distress<sup>59</sup> and overall suicidal risk is increased after voluntary pregnancy termination. Indeed, the risk of suicidal ideation and attempt is at least doubled in women who experience abortion<sup>60,61</sup>; though other studies<sup>62</sup> report that the suicide rate among women who have abortions may be as much as six times higher than that of women who have given birth in the prior year, and double that of women who experience spontaneous abortion. Of note, however, the strongest evidence-based confirmation regarding the relationship between voluntary pregnancy termination and an increase in the suicidal risk derives from a review available from a journal issued by a catholic association.<sup>63</sup> A possible explanation of the increased risk of suicidality after induced abortion is that this decision, for many women, rather than being a relief and/or an answer to their problems, may be additional proof of their worthlessness and may contribute to suicidality<sup>64</sup>; indeed, it has been reported that a post-abortion suicide attempt may be an impulsive act of despair.<sup>65</sup>

C. Most frequently used methods for attempting suicide in pregnancy.

Almost invariably, pregnant women choose poison ingestion or drug overdose for their suicide attempts.<sup>66</sup> Indeed, a Thailand study demonstrated that poisoning is the commonest method of suicide, organophosphate pesticides such as parathion being the poison of choice.<sup>67,68</sup>

In contrast, in Western countries several licit drugs, such as benzodiazepines, are the most frequently used drugs for attempting suicide.<sup>69</sup> However, different medications, including analgesics (most notably acetaminophen), vitamins or iron,

antibiotics, antihistamines or decongestants may be used in order to attempt suicide.<sup>70</sup> Moreover, despite the use of barbiturates being progressively decreased in both neurological and psychiatric practice, such medications are still used by suicidal pregnant women.<sup>71</sup> The period of pregnancy at highest risk of suicide attempts seems to be the second trimester.<sup>72</sup>

D. Maternal and neonatal outcome after attempted suicide

Interesting information comes from studies focused on investigating whether or not suicide attempts by drug overdose may impair obstetric, fetal, and neonatal outcomes. In answer to this question, women diagnosed with drug intoxication during pregnancy were identified in the Regional Hospital Discharge Registry of North Jutland from 1977 to 1999 by linkage of diagnoses for abortion and delivery with diagnoses for intoxication. Hospital medical records were reviewed to obtain data on drug use, dosage, and pregnancy outcome<sup>73</sup>: of 122 women studied, 17 experienced spontaneous abortion. Hence, in general, a drug overdose shortly before or during pregnancy might have been associated with a doubled risk of spontaneous abortion whereas, compared with the background population, there was no increased risk of congenital abnormalities or prematurity. No increase in the structural teratogenic risk was demonstrated in pregnant women who attempted suicide by specifically ingesting very large doses of different barbiturates.<sup>72</sup> Cognitive status and performances at behavioral scales of children exposed to barbiturates throughout placenta also did not indicate any developmental impairment.<sup>72</sup>

Recently, however, very large dose of specific benzodiazepines (such as nitrazepam) used for suicide attempts during pregnancy was associated with increased rates of congenital birth defects (which may be connected with the disruption of protein metabolism in fetal mesenchyma).<sup>74</sup> Moreover, a linked Vital Statistics-Patient Discharge State of California database confirmed that suicide attempts during pregnancy might be associated with significantly higher rates of maternal and perinatal morbidity, and in some cases, perinatal mortality.<sup>38</sup> In fact, such women showed an increase in the risk of premature labor, caesarean delivery, and need for blood transfusion. Analysis of neonatal outcomes also revealed increases in respiratory distress syndrome and low birth weight.<sup>38</sup> In addition, maternal death due to suicide may seriously compromise the future development of orphans. In fact, children who lose their own mother because of suicidal acts show a greater risk of hospitalization for all types of psychiatric disorders and suicide attempts than offspring of parents who have died of other causes.<sup>75</sup>

## Discussion

Review data seem to suggest that prevalence of unipolar depression and suicidal ideation in pregnancy is relatively high<sup>5-8,32,33,35</sup>. However, frequency of suicidal attempts and, especially, rate of death by suicidal acts are low in both developed and developing countries.<sup>36-50</sup> Although rare, consequences of a suicidal attempt are likely to be devastating for the mother-infant pair.<sup>40,73-75</sup>

Moreover, local social and health organizations may consider such women unable to carry out sound maternal functions<sup>76</sup>; hence, infants who survive maternal suicide attempts during pregnancy could be at increased risk of institutionalization. For these reasons, social and health organizations should make all possible efforts to identify women at high suicidal risk, in order to actuate specific programs focused on preventing tragic outcomes. Suicidal literature allows to identify the typical profile of women at high risk of suicide during pregnancy. Hence, the first step towards preventing suicidal events in pregnancy is to identify pregnant women who are at elevated risk factors.<sup>8</sup> In developed countries, those women who have a current or past history of psychiatric disorder<sup>51</sup>, are young, unmarried, have incurred an unplanned pregnancy which has terminated in abortion<sup>52,53</sup>, are addicted to illicit drugs and/or alcohol, lack effective psychosocial support, have suffered from episodes of sexual or physical violence are particularly vulnerable.<sup>8,54-56</sup>

The second step of an effective prevention-plan - since between 4% and 8% of all women experience intimate partner violence during pregnancy<sup>77</sup> - should consist of organizing specific social task-forces to implement and publicize (in both developing and developed countries and either in disadvantaged or privileged social context) specific programs which may facilitate women experiencing abusive relationships to withdraw from such perverse attachments.<sup>8</sup> Such programs should also involve those women who have temporarily chosen to continue the affective bonding of a desire to ensure a "normal" family to their children. Obviously, these programs must include both practical support (e.g., safe accommodation, decent work for unemployed women, and facilitated access to gynecologic care and legal assistance) as well as psychological support, finalized to facilitate the choice of continuing gestation.<sup>78</sup> Alternatively, in the case of an unwanted conception, easy access to public abortion services should be available to those women who choose to interrupt pregnancy.<sup>8,53</sup>

The third step is to promote close cooperation between community psychiatric services, family physicians, hospital gynecological units, and local social services in order to provide the best available care for all women diagnosed with psy-

chiatric disorders who have become mothers. Regrettably, however, up to 70% of patients with mental problems withdraw psychotropic medication during pregnancy and refuse psychiatric counseling<sup>79</sup>; of note, care does not resume during the postpartum period.<sup>80</sup> Thus, such women should be informed, at least during early pregnancy, that the risk of untreated mental disorder outweighs the risk for the fetus of exposure to most psychotropic medications.<sup>81</sup> In fact, several classes of psychotropic seem devoid of significant structural teratogenic risks. Among antipsychotics, chlorpromazine is the drug which shows the highest number of studies suggesting its reproductive safety<sup>82</sup> whereas, among antidepressants, fluoxetine and sertraline should be preferred, since both (anecdotally) have been associated with slight increase in fetal malformations.<sup>81,83</sup> It is also possible to safely start or continue some mood stabilizing agents, since the slight increase in the risk of orofacial cleft suggested for lamotrigine<sup>2</sup> has not been confirmed in recent studies.<sup>84</sup> However, nearly all these medications (with the exception of lamotrigine) have been associated with an increased risk of perinatal complications if used in late pregnancy.<sup>85</sup> Therefore, if clinicians deem indispensable the continuance of psychotropic medications during the last stage of the gestational period, they must firmly suggest that delivery happens in hospitals equipped with specialized neonatal intensive nursery units.<sup>86</sup>

However, the vast majority of maternal deaths due to suicide occur after parturition.<sup>31</sup> Thus, social, psychological, and clinical support provided to such mothers should be continued for at least the first six months of puerperium. The prolonged support may reduce the risk of other tragic events, such as neonaticide (child homicide on the first 24 hours of life) and infanticide (child homicide within the first year of life), which could be brought by mothers who develop antenatal depression and continue to be depressed after parturition.<sup>87</sup> The support resources should gradually be discontinued only when the personal and clinical situation of the mothers appears to be satisfactorily and definitively stabilized.

## Conclusions

Although reviewed studies do not allow us to differentiate between factors associated with suicidal ideation and factors which may contribute to the occurrence of violent and definitive suicidal acts, available data clearly indicate a noteworthy discrepancy between the prevalence of suicidal ideation during pregnancy (relatively high) and the rate of death due to intentional self-harm (extremely low). This discrepancy seems to suggest that maternal death due to suicide may occur merely in the presence of several, concomitant triggers

which involve different areas of the complex maternal universe such as compromised mental stability, weakness of the desire to experience maternity, unhealthy life-style, poor affective liaisons and family support, disadvantaged economic conditions, difficulty in adherence to rigid cultural and religious rules (especially in contexts which stigmatize any female non-orthodox behaviors), and a previous history of personal

traumatic experiences (including intimate partner violence and induced abortion).

**Funding:** none

**Competing interest:** The author declares that he has no competing interests

**Ethical approval:** none

## References

1. Murray CJL, Lopez AD, eds. The Global Burden of Disease: A comprehensive assessment of mortality and disability from disease, injuries and risk factors in 1990 and projected to 2020. Cambridge, MA: Harvard University Press, 1996.
2. Gentile S. Prophylactic treatment of bipolar disorder in pregnancy and breastfeeding: focus on emerging mood stabilizers. *Bipol Disord*. 2006 Jun;8(3):207-20.
3. Burt VK, Rasgon N. Special considerations in treating bipolar disorder in women. *Bipol Disord*. 2004 Feb;6(1):2-13.
4. Yonkers KA, Wisner KL, Stowe Z, Leibenluft E, Cohen L, Miller L, et al. Management of bipolar disorder during pregnancy and postpartum period. *Am J Psychiatry*. 2004 Apr;161(4):608-20.
5. Kumar R, Robson KM. A prospective study of emotional disorders in childbearing women. *Br J Psychiatry*. 1984 Jan;144:35-47.
6. Sundstrom IM, Bixio M, Bjorn I, et al. Prevalence of psychiatric disorders in gynaecologic outpatients. *Am J Obstet Gynecol*. 2001;184(2):8-13.
7. Hallbreich U. Prevalence of mood symptoms and depression during pregnancy: implications for clinical practice and research. *CNS Spectr*. 2004 Mar;9(3):177-84.
8. Gausia K, Fisher C, Ali M, Oosthuizen J. Antenatal depression and suicidal ideation among rural Bangladeshi women: a community-based study. *Arch Womens Ment Health*. 2009 Oct;12(5):351-8.
9. Orr ST, Miller CA. Maternal depressive symptoms and the risk of poor pregnancy outcome. Review of the literature and preliminary findings. *Epidemiol Rev*. 1995;17(1):165-71.
10. Stott DH. Follow-up study from birth of the effects of prenatal stresses. *Dev Med Child Neurol*. 1973 Dec ;15(6):770-87.
11. Kurki T, Hiilesmaa V, Raitasalo R, Mattila H, Ylikorkala O. Depression and anxiety in early pregnancy and risk of preeclampsia. *Obstet Gynecol*. 2000 Apr;95(4):487-90.
12. Freeman MP. Perinatal Psychiatry: risk factors, treatment data, and specific challenges for clinical researchers. *J Clin Psychiatry*. 2008 Apr;69(4):633-4.
13. Steer RA, Scholl TO, Hediger ML, Fischer RL. Self-reported depression and negative pregnancy outcomes. *J Clin Epidemiol*. 1992 Oct;45(10):1093-9.
14. Hollins K. Consequences of antenatal mental health problems for child health and development. *Curr Opin Obstet Gynecol*. 2007 Dec;19(6):568-72.
15. Diego MA, Field T, Hernandez-Reif M, Cullen C, Schanberg S, Kuhn C. Prepartum, postpartum, and chronic depression effects on newborns. *Psychiatry*. 2004 Spring;67(1):63-80.
16. Surkarn PJ, Kawachi I, Ryan LM, Berkman LF, Carvalho Vieira LM, Peterson KE. Maternal depressive symptoms, parenting self-efficacy, and child growth. *Am J Public Health*. 2008 Jan;98(1):125-32.
17. Hodgkinson SC, Colantuoni E, Roberts D, Berg-Cross L, Belcher HM. Depressive symptoms and birth outcomes among pregnant teenagers. *J Pediatr Adolesc Gynecol*. 2010 Feb; 23(1):16-22.
18. Heron J, O'Connor TG, Evans J, Golding J, Glover V, ALSPAC Study Team. The course of anxiety and depression through pregnancy and the postpartum in a community sample. *J Affect Disord*. 2004 May;80(1):65-73.
19. Murray L, Fiori-Cowley A, Hooper R, Cooper P. The impact of postnatal depression and associated adversity on early mother-infant interactions and later infant outcome. *Child Dev*. 1996 Oct;67(5):2512-26.
20. Figueiredo B, Costa R, Pacheco A, Pais A. Mother-to infant emotional involvement at birth. *Maternal Child Health J*. 2009 Jul;13(4):539-49.
21. Cohn JF, Tronick E. Specificity of infants' response to mothers' affective behavior. *J Am Acad Child Adolesc Psychiatry*. 1989 Mar;28(2):242-8.
22. Hart S, Field T, del Valle C, Pelaez-Nogueras M. Depressed mothers' interactions with their one-year-old infants. *Infant Behav Dev*. 1998;21:519-25.
23. Dunham P, Dunham F, Hurshman A, Alexander T. Social contingency effects on subsequent perceptual-cognitive tasks in young infants. *Child Dev*. 1989 Dec;60(6):1486-96.

24. Murray L, Hipwell A, Hooper R, Stein A, Cooper P. The cognitive development of 5-year-old children of postnatally depressed mothers. *J Child Psychol Psychiatry*. 1996 Nov;37(8):927-35.
25. Hay DF, Pawlby S, Waters CS, Perra O, Sharp D. Mothers' antenatal depression and their children's antisocial outcomes. *Child Dev*. 2010 Jan-Feb;81(1):149-65.
26. Beautrais AL, Joyce PR, Mulder RT, Fergusson DM, Deavoll BJ, Nightingale SK. Prevalence and comorbidity of mental disorders in persons making serious suicide attempts: a case-control study. *Am J Psychiatry*. 1996 Aug;153(8):1009-14.
27. World Health Organization. *International statistical classification of disease and related health problems, 10th revision (ICD-10)*. Geneva: WHO, 1993.
28. Lewis G, Drife J. *Why mothers die 1997-1999: the fifth report of the confidential enquiry into maternal deaths in the United Kingdom*. London: RCOG Press, 2001.
29. White C. Depression and obesity are major causes of maternal deaths in Britain. *BMJ*. 2004; 329:1205.
30. Högberg U, Innala E, Sandström A. Maternal mortality in Sweden, 1980-1988. *Obstet Gynecol*. 1994 Aug;84(2):240-4.
31. Lindahl V, Pearson JL, Colpe L. Prevalence of suicidality during pregnancy and the postpartum. *Arch Womens Ment Health*. 2005 Jun;8(2):77-87.
32. Frautschi S, Cerulli A, Maine D. Suicide during pregnancy and its neglects as a component of maternal mortality. *Int J Gynaecol Obstet*. 1994;47(3):275-84.
33. Newport DJ, Levey LC, Pennell PB, Ragan K, Stowe ZN. Suicidal ideation in pregnancy: assessment and clinical implications. *Arch Womens Ment Health*. 2007;10(5):181-7.
34. Lester D. The timing of attempted suicide during pregnancy. *Acta Paediatr Hung*. 1987;28(3-4):259-60.
35. Freitas GV, Botega NJ. Prevalence of depression, anxiety and suicide ideation in pregnant adolescents. *Rev Assoc Med Bras*. 2002 Jul;48(3):245-9.
36. Bødker B, Hvidman L, Weber T, Møller M, Aarre A, Nielsen KM, et al. Maternal deaths in Denmark 2002-2006. *Acta Obstet Gynecol Scand*. 2009;88(5):556-62.
37. Gissler M, Berg C, Bouvier-Colle MH, Buekens P. Injury deaths, suicides and homicides associated with pregnancy, Finland 1987-2000. *Eur J Public Health*. 2005 Oct;15(5):459-63.
38. Högberg U, Innala E, Sandström A. Maternal mortality in Sweden, 1980-1988. *Obstet Gynecol*. 1994 Aug;84(2):240-4.
39. Department of health. *Why Mothers Die*. Stationery office, <http://www.archive.official-documents.co.uk/document/doh/wmd/wmd-12.htm>, Accessed 30 March 2010.
40. Gandhi SG, Gilbert WM, McElvy SS, El Kady D, Danielson B, Xing G, et al. Maternal and neonatal outcome after attempted suicide. *Obstet Gynecol*. 2006 May;107(5):984-90.
41. Dannenber AL, Carter DM, Lawson HW, Ashton DM, Dorfman SF, Graham EH. Homicide and other injuries as causes of maternal death in New York City, 1987 through 1991. *Am J Obstet Gynecol*. 1995 May;172(5):1557-64.
42. Marzuk PM, Tardiff K, Leon AC, Hirsch CS, Portera L, Hartwell N, et al. Lower risk of suicide during pregnancy. *Am J Psychiatry*. 1997 Jan;154(1):122-3.
43. Barno A. Criminal abortion deaths, illegitimate pregnancy deaths, and suicides in pregnancy. Minnesota, 1950-1965. *Am J Obstet Gynecol*. 1967 Jun 1;98(3):356-67.
44. Rochat RW, Koonin LM, Atrash HK, Jewett JF. Maternal mortality in the United States: report from the Maternal Mortality Collaborative. *Obstet Gynecol*. 1988 Jul;72(1):91-7.
45. Christiansen LR, Collins KA. Pregnancy-associated deaths: a 15-year retrospective study and overall review of maternal pathophysiology. *Am J Forensic Med Pathol*. 2006 Mar;27(1):11-9.
46. Parson LH, Harper MA. Violent maternal deaths in North Carolina. *Obstet Gynecol*. 1999 Dec;94(6):990-3.
47. Turner LA, Kramer MS, Liu S; Maternal mortality and morbidity Study Group of the Canadian Perinatal Surveillance System. Cause-specific mortality during and after pregnancy and the definition of maternal death. *Chronic Dis Can*. 2002;23(1):31-6.
48. Chen YH, Lau G. Maternal deaths from suicide in Singapore. *Singapore Med J*. 2008 Sep;49(9):694-7.
49. Mhango C, Rochat R, Arktu A. Reproductive mortality in Lusaka, Zambia, 1982-1983. *Stud Fam Plann*. 1986 Sep;17(5):243-51.
50. Mbizvo MT, Fawcus S, Lindmark G, Nyström L. Maternal mortality in rural and urban Zimbabwe: social and reproductive factors in an incident case-referent study. *Soc Sci Med*. 1993 May;36(9):1197-205.
51. Frautschi S, Cerulli A, Maine D. Suicide during pregnancy and its neglect as a component of maternal mortality. *Int J Gynecol Obstet*. 1994 Dec;47(3):275-84.
52. Appleby L. Suicide during pregnancy and in the first postnatal year. *BMJ*. 1991 Jan 19;302(6769):137-40.
53. Gentile S. Schizoaffective disorder in women with childbearing potential: focus on treatment with newer and emerging mood stabilizers. In: Murray WH ed: *Schizoaffective Disorder: New research*. New York: Nova Publishers, 2006: 187-220.



54. Martin SL, Macy RJ, Sullivan K, Magee ML. Pregnancy-associated violent deaths: the role of intimate partner violence. *Trauma Violence Abuse*. 2007 Apr;8(2):135-48.
55. Rayburn W, Aronow R, DeLancey B, Hogan MJ. Drug overdose during pregnancy: an overview from a metropolitan poison control center. *Obstet Gynecol*. 1984 Nov;64(5):611-4.
56. Rodriguez MA, Bauer HM, McLoughlin E, Grumbach K. Screening and intervention for intimate partner abuse: practices and attitudes of primary care physicians. *JAMA*. 1999 Aug 4;282(5):468-74.
57. Whitlock FA, Edwards JE. Pregnancy and attempted suicide. *Compr Psychiatry*. 1968 Jan;9(1):1-12.
58. Gradus JL, Qin P, Lincoln AK, Miller M, Lawler E, Sørensen HT, et al. Acute stress reaction and completed suicide. *Int J Epidemiol*. 2010 Jul 12.
59. Söderberg H, Janzon L, Sjöberg NO. Emotional distress following induced abortion: a study of its incidence and determinants among abortees in Malmö, Sweden. *Eur J Obstet Gynecol Reprod Biol*. 1998 Aug;79(2):173-8.
60. Mota NP, Burnett M, Sareen J. Associations between abortion, mental disorders, and suicidal behaviour in a nationally representative sample. *Can J Psychiatry*. 2010 Apr;55(4):239-47.
61. Reardon DC, Ney PG, Scheuren F, Cogle J, Coleman PK, Strahan TW. Deaths associated with pregnancy outcome: a record linkage study of low income women. *South Med J*. 2002 Aug;95(8):834-41.
62. Gissler M, Berg C, Bouvier-Colle MH, Buekens P. Injury deaths, suicides and homicides associated with pregnancy, Finland 1987-2000. *Eur J Public Health*. 2005 Oct;15(5):459-63.
63. Thorp JM, Hartmann KE, Shadigan E. Long-term physical and psychological health consequences of induced abortion: a review of evidence. *Linacre Q*. 2005 Feb;72(1):44-69.
64. Gissler M, Hemminki E, Lönnquist J. Suicides after pregnancy. Author's reply. *BMJ*. 1997;314:902-3.
65. Reardon DC. The abortion/suicide connection. Available at: [http://www.abortionfacts.com/reardon/abortion\\_and\\_suicide.asp](http://www.abortionfacts.com/reardon/abortion_and_suicide.asp), Accessed 9 Sept 2010.
66. Rebar RW. Maternal and perinatal morbidity associated with suicide attempts in pregnant women. *J Watch Gen Med* 2006, available at: <http://general-medicine.jwatch.org/cgi/content/full/2006/613/4>, Accessed 9 July 2007.
67. Eungprabhanth V. Suicide in Thailand. *Forensic Sci*. 1975 Feb;5(1):43-51.
68. Jajoo M, Saxena S, Pandey M. Transplacentally acquired organophosphorus poisoning in a newborn: case report. *Ann Trop Paediatr*. 2010;30(2):137-9.
69. Sein Anand J, Chodorowski Z, Ciechanowicz R, Klimaszuk D, Lukasik-Glebocka M. Acute suicidal self-poisonings during pregnancy. *Przegl Lek*. 2005;62(6):434-5.
70. Rayburn W, Aronow R, DeLancey B, Hogan MJ. Drug overdose during pregnancy: an overview from a metropolitan poison control center. *Obstet Gynecol*. 1984 Nov;64(5):611-4.
71. Timmermann G, Czeizel AE, Bánhidly F, Acs N. A study of the teratogenic and fetotoxic effects of large doses of barbital, hexobarbital and butobarbital used for suicide attempts by pregnant women. *Toxicol Ind Health*. 2008 Feb;24(1-2):109-19.
72. Lester D. The timing of attempted suicide during pregnancy. *Acta Paediatr Hung*. 1987;28(3-4):259-60.
73. Flint C, Larsen H, Nielsen GL, Olsen J, Sørensen HT. Pregnancy outcome after suicide attempt by drug use: a Danish population-based study. *Acta Obstet Gynecol Scand*. 2002 Jun;81(6):516-22.
74. Gidai J, Acs N, Bánhidly F, Czeizel AE. Congenital abnormalities in children of 43 pregnant women who attempted suicide with large doses of nitrazepam. *Pharmacoepidemiol Drug Saf*. 2010 Feb;19(2):175-82.
75. Wilcox HC, Kuramoto SJ, Lichtenstein P, Långström N, Brent DA, Runeson B. Psychiatric morbidity, violent crime, and suicide among children and adolescents exposed to parental death. *J Am Acad Child Adolesc Psychiatry*. 2010 May;49(5):514-23.
76. Coverdale JH, Aruffo JA. Family planning needs of female chronic psychiatric outpatients. *Am J Psychiatry*. 1989 Nov;146(11):1489-91.
77. New York City Department of Health and Mental Hygiene. Intimate partner violence. <http://www.nyc.gov/html/doh/html/epi/domviol.shtml>, Accessed 9 April 2010.
78. Gentile S. Special care-needs for schizoaffective women in childbearing age [Abstract]. *Turk J Psychiatry*. 2006;2( suppl 1):271.
79. Goodman JH. Women's attitudes, preferences, and perceived barriers to treatment for perinatal depression. *Birth*. 2009 Mar;36(1):60-9.
80. Bennett IM, Marcus SC, Palmer SC, Coyne JC. Pregnancy-related discontinuation of antidepressants and depression care visits among Medicaid recipients. *Psychiatr Serv*. 2010 Apr;61(4):386-91.
81. Gentile S. The safety of newer antidepressants in pregnancy and breastfeeding. *Drug Saf*. 2005;28(2):137-52.
82. Gentile S. Antipsychotic therapy during early and late pregnancy. A systematic review. *Schizophr Bull*. 2010 May;36(3):518-44.
83. Gentile S, Bellantuono C. Selective serotonin reuptake inhibitor exposure during early pregnancy and the risk of fetal major malformations: focus on paroxetine. *J Clin Psychiatry*. 2009 Mar;70(3):414-22.

*Archive of SID*

84. Gentile S. The safety of mood stabilizing medication in pregnancy and lactation [Abstract]. *World J Biol Psychiatry*. 2009;10(suppl1): 87.
85. Gentile S. On categorizing gestational, birth, and neonatal complications following late pregnancy exposure to antidepressants: the prenatal antidepressant exposure syndrome. *CNS Spectr*. 2010 Mar;15(3):167-85.
86. Gentile S. Prolonged exposure to SRIs during pregnancy increases risks of adverse neonatal outcomes. *Evid Based Ment Health*. 2009 Feb;12(1):27.
87. Spinelli MG. Maternal infanticide associated with mental illness: prevention and the promise of saved lives. *Am J Psychiatry*. 2004 Sep;161(9):1548–57.

# SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



عضویت در خبرنامه



فیلم های آموزشی

## کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



مباحث پیشرفته یادگیری عمیق؛  
شبکه های توجه گرافی  
(Graph Attention Networks)



کارگاه آنلاین آموزش استفاده از  
وب آوساینس



کارگاه آنلاین مقاله روزمره انگلیسی