

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



عضویت در خبرنامه




فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



کارگاه آنلاین آشنایی با پایگاه های اطلاعات علمی بین المللی و ترند های جستجو



مباحث پیشرفته یادگیری عمیق؛ شبکه های توجه گرافی (Graph Attention Networks)



کارگاه آنلاین مقاله نویسی IEEE و ISI ویژه فنی و مهندسی

Original Article

Comparison of interdental papilla around single implants in the anterior maxilla between two implant systems: A cohort study

Masoumeh Khoshhal¹ • Fariborz Vafaei² • Mahsa Najafi³ • Masoumeh Nikkhah^{4*}

¹Dental Implants Research Center, Hamadan University of Medical Sciences, Hamadan, Iran

²Assistant Professor, Department of Prosthodontics, Faculty of Dentistry, Hamadan University of Medical Sciences, Hamadan, Iran

³Dentist, Hamadan, Iran

⁴Assistant Professor, Department of Periodontics, Faculty of Dentistry, Guilan University of Medical Sciences, Rasht, Iran

*Corresponding Author; E-mail: nikkhah_m65@yahoo.com

Received: 26 February 2017; Accepted: 15 January 2018

J Dent Res Dent Clin Dent Prospect 2018; 12(1):38-44 | doi: 10.15171/joddd.2018.006

This article is available from: <http://joddd.tbzmed.ac.ir>

© 2018 Khoshhal et al. This is an Open Access article published and distributed by Tabriz University of Medical Sciences under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background. In successful replacement of a tooth with a dental implant, soft tissue esthetic is as important as stability and function of the implant. Quality and quantity of the peri-implant mucosa can influence esthetic outcomes. This study assessed implant esthetic success of two different implant systems. In this regard the interdental papilla was evaluated and the relationship between implant type and crestal bone loss adjacent to implant was assessed.

Methods. Eighteen patients (11 males, 7 females) with a total of 18 implants participated in this historical cohort study. Patients were divided into two groups based on the type of implants: Implantium group and SPI group; 36 interproximal papillae were evaluated photographically, using Jemt's papillary presence index (PPI). Radiographic analysis was carried out to find out the relation between bone loss and type of implant. Analysis of data was performed with SPSS 18, using Fisher's exact test, independent t-test, Spearman's correlation coefficient and ANOVA.

Results. Comparison of photographs did not show a statistically significant difference in PPI between the two groups (P=0.94). Radiographic evaluation of crestal bone loss adjacent to implant shoulder did not reveal significant differences between the two groups (P=0.30).

Conclusion. Implant therapy in the anterior maxilla, using Implantium or SPI system, did not result in significant differences in esthetics. In this study, there was an inverse relationship between the distance of contact point to bone crest and papilla index (P=0.002 in the SPI group) (P=0.02 in the Implantium group).

Key words: Interdental papilla, maxilla, single-tooth implants.

Introduction

A maxillary anterior missing tooth is usually replaced for functional and aesthetic reasons.¹

Implants are a quick way to replace teeth and achieve the desired results. Therefore, reconstructing the function and esthetic appearance by using a single-tooth implant compared to tooth-supported restorations has become a common treatment.

Although dental implants have long-term success in the reconstruction of edentulous patients, achieving desired beauty of soft tissue around anterior single implants and continued successful results is an equally important and concerned issue.

Gingival facial appearance and interproximal papilla are the most important factors that make a beautiful soft tissue.²

The interproximal space depends on several factors that can compromise the interdental papilla. Improper contours of restorations or prosthetic crowns, abnormal tooth shape, traumatic flossing and interproximal hygiene procedures and, importantly, periodontal diseases can cause recession in the interdental papilla.³

It is important to note that the marginal bone level can be maintained in the face of functional loading by a properly designed implant.⁴

The marginal bone level of the peri-implant and soft tissue are strongly related to each other, which determine the aesthetic outcome.¹

Studies have shown that the height of papilla depends on the distance from the contact point to the crest of the alveolar bone of the adjacent teeth. On the other hand, anatomy of the adjacent teeth and the distance between the implants and natural teeth can affect the papilla formation.^{5,6}

Although a number of studies have suggested that peri-implant bone level is a decisive factor for the presence of papilla between the tooth and implant, some studies have found that connective tissue attachment and clinical attachment level are also important.^{7,8}

During a 6-month period, after implant surgical procedure, the level of soft tissue was enhanced considerably at tooth-facing sites (a mean of 1.1 mm) at proximal space, but no significant increase was found at implant-facing sites.⁹ Studies have shown differences between the level of the papilla in the mesial and distal aspects around single implant-supported restorations. The distal papilla has a lower score in Jemt classification compared to the mesial papilla adjacent to tooth surfaces.^{5,10} In 1997 Jemt¹⁰ reported that during a 1–3-year period after single-implant restorations, the adjacent papilla regenerates partially without any clinical intervention. The reason for this spontaneous recovery of the papillae is unclear, but it might be suggested that the inflammation caused by accumulation of plaque in the proximal space leads to swelling of the soft tissue.

Won Lee et al¹¹ showed that the height of the interproximal papilla between the single implant and the adjacent tooth is approximately equal to the interproximal papilla on the contra-lateral side. The alveolar

bone loss on both sides of a single implant site would be similar.

A distance of 3 mm between two implants has been recommended to ensure the presence of papilla. On the other hand, some studies have shown that the interproximal papilla is not influenced by the distance between the implant and tooth.^{12,13}

Several studies have been performed on the effect of the distance from the contact point of the implant restoration and the adjacent tooth to the crestal bone and the effect of implant design and diameter, implant position and soft tissue biotype on papilla around the implant.¹⁴⁻¹⁷

Many researchers have assessed the effect of different variables on peri-implant bone but have not considered its effect on soft tissues. However, evaluation of peri-implant tissues with different implant designs is mostly limited to assessing peri-implant bone level.^{18,19} The aim of this study was to compare and assess soft tissue around two commonly used implant systems (SPI and Implantium) with different designs with or without microthreads.

Methods

In the present cohort study, 18 patients were selected among patients who had received single implants in the anterior area of the maxilla 1.5 years previously in a private clinic in Hamadan. The subjects were divided into two groups (n=9). The first group was treated by the Implantium system (Dentium, Korea) and the latter was treated by the SPI implant system (Thommen, Switzerland).

Features of these two implant systems were as follows: Body shape of these two types of implants was cylindrical. The abutment connection of these implants was internal and abutment connection type was internal hex. The implant body surface of the SPI system was acid etched, sandblasted and conditioned by hydroxylation. The implant body surface of Implantium was SLA (Sandblast Large grit Acid etch). The implant body thread of SPI was V-shaped and that of Implantium was buttress. The basic difference between these two implants systems was the presence of microthread in the neck of Implantium.

The inclusion criteria of the patients consisted of no systemic disease, no pregnancy, no smoking and drinking, no periodontal disease, good oral hygiene, no bleeding on probing (BOP) (BI=0), no pocket depths >3 mm, no bone loss (the distance between CEJ of the adjacent tooth and crestal bone was not >2 mm), implant in a safe distance from the adjacent tooth (at least 1.5 mm), thick gingival biotype, no history of radiotherapy and chemotherapy, and no use of

immunosuppressive drugs. Exclusion criteria included uncontrolled active systemic disease, the need for bone grafts (GBR), thin gingival biotype, and prolonged steroid therapy.

In all the patients all the surgical procedures were performed by one periodontist and implant crowns were prepared by one prosthodontist. All the surgical and prosthetic procedures were the same. All PFM restorations were prepared by one technician and laboratory. All the implant restorations were cemented. All the patients had primary standard parallel periapical radiography. At the time of study, 1.5 years after implant insertion, standard parallel periapical radiography was obtained. To assess the presence of interproximal papilla, standard photographs were taken using a SLR camera (Canon 550D) with a 100-mm macro lens and ring flash.

In the initial radiograph, the upper edge of the implant shoulder and in the radiograph 1.5 years after the surgical procedure the distance between the implant and abutment was a considered reference line.

In the initial radiograph the distance from the contact point of the implant and bone to the reference line and also the distance from the CEJ of the adjacent tooth to the contact point of the crestal bone and tooth were measured. In the radiograph 1.5 years after implant insertion, the distance from the contact point of the implant restoration and the adjacent tooth to the crestal bone was measured. In addition, the distance from the contact point of the crestal bone and implant to the reference line and also the distance from the CEJ of the adjacent tooth to the contact point of the crestal bone and tooth were measured.

Presence of papilla between the implant and the adjacent tooth was measured and recorded based on Jemt index 1.5 years after implant insertion surgery. The distance from the height of the crestal bone and the implant shoulder was measured with a digital Vernier on initial the radiograph and the radiograph 1.5 years after surgery and recorded in mm. The distance between the height of the contact point of implant restoration and the adjacent tooth to crestal bone was measured using a digital Vernier on the radiograph 1.5 years after implant insertion surgery and recorded in mm. The distance between the CEJ of the adjacent tooth of implant to crestal bone was measured using a digital Vernier on the initial radiograph and the radiograph 1.5 years after surgery and recorded in mm.

All the measurements were performed by one person twice within two weeks and an average of the two measurements was used as the final measure.

The presence of papilla around implants was evaluated by photography and Jemt index.

Papilla index (PI) grading was as follows (Figure 1):

- Score 0: no papilla in the interproximal space
- Score 1: presence of less than 50% of the papilla height
 - Score 2: presence of at least 50% of the papilla height but not all the interproximal space
 - Score 3: the papilla completely fills the interproximal space and is coordinated by the adjacent papillae, with a favorable gingival contour
 - Score 4: the hyperplastic papillae that covers too much of the single implant restoration and/or the adjacent tooth, with unfavorable gingival contour²⁰

For this purpose, the line that connects the most apical point of the implant restoration to the most apical point of the crown of the adjacent tooth (zenith) was considered as a reference. The vertical distance from the contact point to this line was divided into two parts. Then the position of the papilla tip was evaluated by Jemt index.

SPSS 18 was used for data analysis. To compare the presence of the interdental papilla in the two groups, Fisher's exact test was used. Independent t-test was used to compare the distance between the crestal bone and the CEJ of the adjacent teeth, implant shoulder, the contact point of the implant restoration and the adjacent tooth on the initial and secondary radiographs. Spearman's correlation coefficient and ANOVA were used to examine the relationship between different variables and the presence of papilla. Level of significance was set at $P < 0.05$.

Results

The mean age of the subjects was 40 years old, ranging from 26 to 55. Seven females and 11 males took part in the study. Totally 36 papillae and 18 interdental areas and papillae adjacent to implants were assessed in each group. A total of 12, 14 and 10 papillae were evaluated around canines, lateral incisors and central incisors, respectively.

The results in the two groups were matched in terms of the length and diameter of the implants.

Kolmogorov-Smirnov test showed that the distribution of all the data in Table 1 was normal. The mean changes in the distance between the crestal bone and implant shoulder in the mesial and distal aspects in the Implantium group were 0.68 mm and 1.31 mm, respectively with 1.09 and 0.7 mm, respectively, in the SPI group. The distance from the implant shoulder to the crestal bone increased significantly, only in the distal aspect of Implantium ($P=0.008$) (Table 1). The mean bone loss adjacent to the implant shoulder in the Implantium group was 1.32 mm, with 0.7 mm in the

Table 1. Mean distances from the implant shoulder to the crestal bone in mm in the two groups

Implant type	Zone	Initial radiography	After 18 months	The mean changes	Paired samples t-test
Implantium	Mesial	1.21	1.9	0.68	P= 0.09
	Distal	0.79	2.11	1.31	P= 0.008
SPI	Mesial	1.95	3.05	1.09	P= 0.09
	Distal	1.76	2.47	0.7	P= 0.37

Significant level $P > 0.05$

SPI group. Independent t-test did not reveal significant differences between the two groups (Table 2). Based on ANOVA there was no significant correlation between bone loss and papilla index in the two groups (Table 2). Kolmogorov-Smirnov test showed normal distribution of all the data in Table 3. As shown in Table 3 in the interval between implant insertion surgery and 18 months later, changes in bone level of the teeth adjacent to implant was not significant (Table 3). As shown in Table 4, there was no significant correlation between bone loss around the tooth adjacent to the implant in the mesial and distal aspects and papilla index in the two groups (Table 4). Based on Table 5, in both groups there was significant correlation between the distance from the contact point of the implant restoration and the adjacent tooth to crestal bone and papilla index. Correlation coefficients in the Implantium and SPI groups were -0.28 was -0.15, respectively, indicating an inverse relationship between the two variables. This means that by increasing the distance between the contact point and crest, the papilla index decreased (Table 5).

Discussion

Replacing lost teeth with implants is an ideal and successful treatment option. However, currently the main concern is not osseointegration of dental implants. Successful implant treatment means achieving the

best esthetic results, in addition to stability and function of the implant.²¹

From an esthetic point of view presence and maintenance of interdental papilla is one of the main factors. In fact, esthetic results of implant treatment are not exclusively associated with the form of the crown, but they are mainly influenced by the topography of the surrounding soft tissue.

The design of this study was based on the recommendations of a study of Sorni-Broker et al, who suggested further studies to examine the effect of the micro- and macro-structure of the implant on the position of soft tissues, especially papilla.²²

In this study, we compared interdental papillae in two different commonly used implant systems (Implantium and SPI). Implantium system has microthreads in the coronal area of the fixture and the SPI system does not have it.

The strength of this study in comparison with other studies was attention to interdental papilla as a key factor in the esthetic outcomes and success of implants.

It should be noted that most of the studies did not consider our subject and a small number of studies, such as studies by Jemt et al³¹ (2004) and Pier (2011), in which interdental papillae around different designs of implants were compared, had designs and variables different from those of the present study. Therefore, those studies cannot be directly compared with our study.

Table 2. Bone loss adjacent to the implant shoulder and the relationship between bone loss adjacent to the implant shoulder and papilla index in the Implantium and SPI groups

	Implantium	SPI	Level of significance
Mean \pm SD	1.32 \pm 0.97	0.70 \pm 1.41	P = 0.30
F index in ANOVA	4	3.2	P (Implantium) = 0.06 P (SPI) = 0.08

Significant level $P > 0.05$

Table 3. The mean distance between the CEJ of the adjacent tooth and the crestal bone in mm in the two groups

Implant type	Zone	Initial radiography	Secondary radiography	The mean changes	Level of significance
Implantium	Mesial	2.22	2.56	0.33	P= 0.31
	Distal	2.21	2.31	0.09	P= 0.06
SPI	Mesial	2.21	2.67	0.46	P= 0.31
	Distal	2.67	2.80	0.12	P=0.68

Significant level $P > 0.05$

Table 4. The relationship between crestal bone loss of the adjacent teeth and papilla index in the mesial and distal aspects in the two groups

Implant type	Zone	Level of significance based on variance analysis
Implantium	Mesial	P = 0.41
	Distal	P = 0.77
SPI	Mesial	P = 0.35
	Distal	P = 0.29

Significant level $P > 0.05$

Table 5. The relationship between the distance from the contact point of the implant restoration and the adjacent tooth to crestal bone and papilla index

Implant type	Spearman's correlation coefficient	Level of significance based on variance analysis
Implantium	-0.28	P = 0.02
SPI	-0.15	P = 0.002

Significant level $P > 0.05$

The results of this study did not reveal any significant differences in the amount of crestal bone loss around implants in the two groups with and without microthreads.

This is different from the results of studies by Lee et al,²³ Nickening et al²⁴ and Bratu et al.²⁵ These studies suggested that microthreads in the coronal area of the fixture were effective in maintaining bone, and the amount of bone loss around these implants was significantly lower.

The differences between the results of this study and the study by Lee et al might be attributed to the study periods. Lee et al,²³ during the 3-year study period, observed a significant difference between the groups with and without microthreads. However, in our study we evaluated the effect of different implants on papilla after 1.5 years of implant insertion surgery.

Nickening et al²⁴ and Bratu et al²⁵ used implants that were different from the types used in this study. They compared implants with and without microthreads (machined-neck implants). However, in the present study, implants with rough neck surface and with microthreads in one group, and implants with polished neck surface were used in the other group.

A study by Amid et al²⁶ suggested that the implant neck with a microthread design relieved stresses in the bone crest. However, Schrottenboer et al²⁷ suggested that microthreads increased crestal stress upon loading. These differences are explained by different implant systems and research methods of these studies.

Hansson²⁸ suggested that, in the implant neck with a retentive feature, marginal bone loss will be prevented, decreasing peak interfacial shear stress. This study compared two different types of titanium implants by finite element analysis. Lee et al²⁹ also observed that the implant with a rough surface and microthreads at the neck might have a chance of maintaining the level of the marginal bone. Differences in the results of the above studies and the present study

might be explained by different types and surfaces of implants and different study periods.

The results of the present study are consistent with those of studies by Hartog et al¹ and Shin et al.¹⁹ Hartog et al¹ showed that after 18 months of follow-up, no considerable difference was found in the loss of marginal bone on radiographs between implants with smooth and rough necks. Shin et al¹⁹ reported no significant difference between rough-surfaced implants with and without microthreads in the amount of bone loss.

In this study there was no significant correlation between crestal bone loss adjacent to the implant shoulder and also between interproximal bone loss of the tooth adjacent to the implant and papilla index.

Studies by Nisapakultron et al,³⁰ Hartog et al¹ and Kan et al¹⁶ showed that the level of the interproximal bone crest of the adjacent tooth had a significant effect on the papilla level around the maxillary anterior single-tooth implants and was not very much related to bone level around the implant shoulder. This difference might be explained by the different designs of the studies.

Studies by Jemt et al¹⁰ and Henriksson et al³¹ showed that the distance of the contact point of the implant restoration and the adjacent tooth to the crestal bone had no significant effect on papilla presence. However, Choquet et al¹⁴ showed that when the distance between the contact point to the crest was ≤ 5 mm, the papilla was present almost in all the cases. When the distance was ≥ 6 mm, the papilla was present in half of the cases or less.

The results of this study were different from the observations reported by Jemt et al¹⁰ and Henriksson et al³¹ and the presence of a significant inverse relationship between the papilla index and the distance of the contact point of the implant restoration and the adjacent tooth.

This discrepancy might be attributed to different

variables evaluated in studies by Jemt et al and Henriksson et al. They assessed the effect of different implant systems with different abutments on soft tissues.³¹

Overall, in the present study papilla formation in all the groups was acceptable, and none of them showed score 0. Frequencies of scores 2 and 3 in the two groups were the same. There was no significant difference in the amount of bone loss between the two groups. Therefore, both implant types exhibited almost good and comparable esthetic results in the anterior area of the maxilla.

Conclusion

1. Presence of papilla and also the amount of bone loss adjacent to two implant systems did not exhibit significant differences.
2. There was a significant relationship between papilla presence and the distance of the contact point of the implant restoration and the adjacent tooth to the crestal bone.
3. There was no significant relationship between papilla presence and peri-implant bone loss and also bone loss around the adjacent tooth.

For future studies, it is suggested that a greater number of implants be included in each group for comparisons and also the effect of other types of implants on the interdental papilla be evaluated.

Authors' contributions

MKH presented the concept of the study. MKH and FV designed the study. MKH and FV were responsible for the intellectual content of the study. MN and MKH carried out the literature review. The clinical study was conducted by MKH and FV. MN was responsible for data acquisition and data analysis. The manuscript was prepared and edited by MN. The manuscript was reviewed by MKH and MN.

Acknowledgments

This study was part of a thesis by Mahsa Najafi for a DDS degree. The thesis supervisor was Dr. Masoumeh Khoshhah and the counseling professor was Dr. Fariborz Vafaei.

Funding

Hamadan University of Medical Sciences financially supported this study.

Competing interests

The authors declare that they have no significant competing financial, professional or personal interests that might have influenced the performance or presentation of the work described in this manuscript.

Ethics approval

This study was approved by the Medical Ethics and Research Committee of Hamadan University of Medical Sciences.

References

1. den Hartog L, Meijer HJ, Stegenga B, Tymstra N, Vissink A, Raghoobar GM. Single implants with different neck designs in the aesthetic zone: a randomized clinical trial. *Clin Oral Implants Res.* 2011;22(11):1289-97. doi: [10.1111/j.1600-0501.2010.02109.x](https://doi.org/10.1111/j.1600-0501.2010.02109.x)
2. Salama H, Salama M, Garber D, Adar P. Developing optimal peri-implant papillae within the esthetic zone: guided soft tissue augmentation. *J Esthet Dent.* 1995;7(3):125-9. doi: [10.1111/j.1708-8240.1995.tb00729.x](https://doi.org/10.1111/j.1708-8240.1995.tb00729.x)
3. Prato GP, Rotundo R, Cortellini P, Tinti C, Azzi R. Interdental papilla management: a review and classification of the therapeutic approaches. *Int J Periodontics Restorative Dent.* 2004;24(3):246-55. doi: [10.1016/j.prosdent.2004.07.002](https://doi.org/10.1016/j.prosdent.2004.07.002)
4. Misch CE, Qu Z, Bidez MW. Mechanical properties of trabecular bone in the human mandible: implications for dental implant treatment planning and surgical placement. *J Oral Maxillofac Surg.* 1999;57(6):700-6; discussion 6-8. doi: [10.1016/S0278-2391\(99\)90437-8](https://doi.org/10.1016/S0278-2391(99)90437-8)
5. Chang M, Wennstrom JL, Odman P, Andersson B. Implant supported single-tooth replacements compared to contralateral natural teeth. Crown and soft tissue dimensions. *Clin Oral Implants Res.* 1999;10(3):185-94. doi: [10.1034/j.1600-0501.1999.100301.x](https://doi.org/10.1034/j.1600-0501.1999.100301.x)
6. Kourkouta S, Dedi KD, Paquette DW, Mol A. Interproximal tissue dimensions in relation to adjacent implants in the anterior maxilla: clinical observations and patient aesthetic evaluation. *Clin Oral Implants Res.* 2009;20(12):1375-85. doi: [10.1111/j.1600-0501.2009.01761.x](https://doi.org/10.1111/j.1600-0501.2009.01761.x)
7. Grunder U. Stability of the mucosal topography around single-tooth implants and adjacent teeth: 1-year results. *Int J Periodontics Restorative Dent.* 2000;20(1):11-7. doi: [10.11607/prd.00.0354](https://doi.org/10.11607/prd.00.0354)
8. Palmer RM, Farkondeh N, Palmer PJ, Wilson RF. Astra Tech single-tooth implants: an audit of patient satisfaction and soft tissue form. *J Clin Periodontol.* 2007;34(7):633-8. doi: [10.1111/j.1600-051X.2007.01101.x](https://doi.org/10.1111/j.1600-051X.2007.01101.x)
9. Chang M, Wennstrom JL. Peri-implant soft tissue and bone crest alterations at fixed dental prostheses: a 3-year prospective study. *Clin Oral Implants Res.* 2010;21(5):527-34. doi: [10.1111/j.1600-0501.2009.01874.x](https://doi.org/10.1111/j.1600-0501.2009.01874.x)
10. Jemt T. Regeneration of gingival papillae after single-implant treatment. *Int J Periodontics Restorative Dent.* 1997;17(4):326-33. doi: [10.11607/prd.00.0224](https://doi.org/10.11607/prd.00.0224)
11. Lee DW, Huh JK, Park KH, Chai JK, Kim CK, Moon IS. Comparison of interproximal soft tissue height for single implants and contra-lateral natural teeth. *Clin Oral Implants Res.* 2009;20(12):1320-5. doi: [10.1111/j.1600-0501.2009.01737.x](https://doi.org/10.1111/j.1600-0501.2009.01737.x)
12. Gastaldo JF, Cury PR, Sendyk WR. Effect of the vertical and horizontal distances between adjacent implants and between a tooth and an implant on the incidence of interproximal papilla. *J Periodontol.* 2004;75(9):1242-6. doi: [10.1902/jop.2004.75.9.1242](https://doi.org/10.1902/jop.2004.75.9.1242)
13. Zetu L, Wang HL. Management of inter-dental/inter-implant papilla. *J Clin Periodontol.* 2005;32(7):831-9. doi: [10.1111/j.1600-051X.2005.00748.x](https://doi.org/10.1111/j.1600-051X.2005.00748.x)
14. Choquet V, Hermans M, Adriaenssens P, Daelemans P, Tarnow DP, Malevez C. Clinical and radiographic evaluation of the papilla level adjacent to single-tooth dental implants. A

- retrospective study in the maxillary anterior region. *J Periodontol.* 2001;72(10):1364-71. doi:10.1902/jop.2001.72.10.1364
15. Lops D, Chiapasco M, Rossi A, Bressan E, Romeo E. Incidence of inter-proximal papilla between a tooth and an adjacent immediate implant placed into a fresh extraction socket: 1-year prospective study. *Clin Oral Implants Res.* 2008;19(11):1135-40. doi:10.1111/j.1600-0501.2008.01580.x
 16. Kan JY, Rungcharassaeng K. Interimplant papilla preservation in the esthetic zone: a report of six consecutive cases. *Int J Periodontics Restorative Dent.* 2003;23(3):249-59. doi:10.11607/prd.1346
 17. Pradeep AR, Karthikeyan BV. Peri-implant papilla reconstruction: realities and limitations. *J Periodontol.* 2006;77(3):534-44. doi:10.1902/jop.2006.050068
 18. Aloy-Prosper A, Maestre-Ferrin L, Penarrocha-Oltra D, Penarrocha-Diago M. Marginal bone loss in relation to the implant neck surface: an update. *Med Oral Patol Oral Cir Bucal.* 2011;16(3):e365-8. doi:10.4317/medoral.16.e365
 19. Shin YK, Han CH, Heo SJ, Kim S, Chun HJ. Radiographic evaluation of marginal bone level around implants with different neck designs after 1 year. *Int J Oral Maxillofac Implants.* 2006;21(5):789-94.
 20. Jemt T. Regeneration of gingival papillae after single-implant treatment. *Int J Periodontics Restorative Dent.* 1997;17(4):326-33. doi:10.1016/j.prosdent.2005.02.009
 21. Somanathan RV, Simunek A, Bukac J, Brazda T, Kopecka D. Soft tissue esthetics in implant dentistry. *Acta medica (Hradec Kralove).* 2007;50(3):183-6.
 22. Sorni-Broker M, Penarrocha-Diago M, Penarrocha-Diago M. Factors that influence the position of the peri-implant soft tissues: a review. *Med Oral Patol Oral Cir Bucal.* 2009;14(9):e475-9.
 23. Lee DW, Choi YS, Park KH, Kim CS, Moon IS. Effect of microthread on the maintenance of marginal bone level: a 3-year prospective study. *Clin Oral Implants Res.* 2007;18(4):465-70. doi:10.1111/j.1600-0501.2007.01302.x
 24. Nickenig HJ, Wichmann M, Schlegel KA, Nkenke E, Eitner S. Radiographic evaluation of marginal bone levels adjacent to parallel-screw cylinder microthreaded implants and rough-surfaced microthreaded implants using digitized panoramic radiographs. *Clin Oral Implants Res.* 2009;20(6):550-4. doi:10.1111/j.1600-0501.2008.01684.x
 25. Bratu EA, Tandlich M, Shapira L. A rough surface implant neck with microthreads reduces the amount of marginal bone loss: a prospective clinical study. *Clin Oral Implants Res.* 2009;20(8):827-32. doi:10.1111/j.1600-0501.2009.01730.x
 26. Amid R, Raoofi S, Kakhodazadeh M, Movahhedi MR, Khademi M. Effect of microthread design of dental implants on stress and strain patterns: a three-dimensional finite element analysis. *Biomed Tech (Berl).* 2013;58(5):457-67. doi:10.1515/bmt-2012-0108
 27. Schrottenboer J, Tsao YP, Kinariwala V, Wang HL. Effect of microthreads and platform switching on crestal bone stress levels: a finite element analysis. *J Periodontol.* 2008;79(11):2166-72. doi:10.1902/jop.2008.080178
 28. Hansson S. The implant neck: smooth or provided with retention elements. A biomechanical approach. *Clin Oral Implants Res.* 1999;10(5):394-405. doi:10.1034/j.1600-0501.1999.100506.x
 29. Lee SY, Piao CM, Koak JY, Kim SK, Kim YS, Ku Y, et al. A 3-year prospective radiographic evaluation of marginal bone level around different implant systems. *J Oral Rehabil.* 2010;37(7):538-44. doi:10.1111/j.1365-2842.2010.02083.x
 30. Nisapakultorn K, Suphanantachai S, Silkosessak O, Rattana-mongkolgul S. Factors affecting soft tissue level around anterior maxillary single-tooth implants. *Clin Oral Implants Res.* 2010;21(6):662-70. doi:10.1111/j.1600-0501.2009.01887.x
 31. Henriksson K, Jemt T. Measurements of soft tissue volume in association with single-implant restorations: a 1-year comparative study after abutment connection surgery. *Clin Implant Dent Relat Res.* 2004;6(4):181-9. doi:10.1111/j.1708-8208.2004.tb00034.x

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



عضویت در خبرنامه




فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



کارگاه آنلاین آشنایی با پایگاه های اطلاعات علمی بین المللی و ترند های جستجو



مباحث پیشرفته یادگیری عمیق؛ شبکه های توجه گرافی (Graph Attention Networks)



کارگاه آنلاین مقاله نویسی IEEE و ISI ویژه فنی و مهندسی