

SID



سرویس های
ویژه



سرویس ترجمه
تخصصی



کارگاه های
آموزشی



بلاگ
مرکز اطلاعات علمی



سامانه ویراستاری
STES



فیلم های
آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی

کارگاه آنلاین
بررسی مقابله ای متون (مقدماتی)

کارگاه آنلاین
پروپوزال نویسی و پایان نامه نویسی

کارگاه آنلاین آشنایی با پایگاه های اطلاعات علمی
بین المللی و
ترند های جستجو

Effect of Extremely Low Frequency Electromagnetic Field and GABA_A Receptors on Serum Testosterone Level of Male Rats

Mahnaz Taherianfard ^{1,*}, Aminolah Bahaddini ², Sara Keshtkar ³, Mehdi Fazeli ⁴, Tahora Shomali ⁴

¹Department of Physiology, School of Veterinary Medicine, Shiraz University, Shiraz, IR Iran

²Department of Biology, Faculty of Basic Science, Shiraz University, Shiraz, IR Iran

³Department of Biology, University of Scientific-Practical, Shiraz University, Shiraz, IR Iran

⁴Department of Pharmacology, School of Veterinary Medicine, Shiraz University, Shiraz, IR Iran

*Corresponding author: Mahnaz Taherianfard, Department of Physiology, School of Veterinary Medicine, Shiraz University, Shiraz, IR Iran. P.O.Box: 1731, Postcode: 71345. Tel: +98-7112286950, Fax: +98-7112286940, E-mail: taherian@shirazu.ac.ir.

Received: March 03, 2013; Revised: July 09, 2013; Accepted: July 27, 2013

Background: GABA can influence the steroidogenesis in peripheral and central nervous systems.

Objectives: The present study investigates the interactive effect of GABA_A receptors and extremely low frequency electromagnetic field on serum testosterone level of male rats.

Patients and Methods: Fifty adult male rats were randomly assigned into 10 groups. Groups 2, 4, 6, 8, and 10 were exposed to ELF-EMF for 30 days 8hrs per day; while, the remaining groups (1, 3, 5, 7, and 9) were sham exposed animals. At the end of the experiment, animals in groups 1 and 2 received normal saline; while, animals in groups 3 and 4 were treated with 1 mg/kg of bicuculline methiodide, and for animals of groups 5 and 6, 3 mg/kg of bicuculline was injected. Animals of groups 7 and 8 were treated with 0.5 mg/kg of muscimol hydrobromide and rats in groups 9 and 10 received 2 mg/kg muscimol hydrobromide. About forty minutes after the injection, blood samples were collected and serum testosterone level was assayed using RIA.

Results: Administration of muscimol hydrobromide at both doses to sham exposed rats significantly decreased serum testosterone level as compared to sham exposed animals which received saline. Administration of bicuculline methiodide without exposure to ELF-EMF, had no significant effect on testosterone level as compared to group 1. Serum testosterone levels of rats in different groups, exposed to ELF-EMF were statistically the same. Moreover, serum testosterone of exposed and sham exposed rats in each treatment showed no significant difference.

Conclusions: No interactivity is present in modulatory effects of GABA_A receptors and ELF-EMFs on serum testosterone of male rats.

Keywords: Extremely Low Frequency Electromagnetic Fields; Muscimol Hydrobromide; bicuculline; Testosterone; Rat

1. Background

Testosterone is the major male sex hormone which is primarily synthesized by Leydig cells of the testes and plays a critical role in the function and development of male reproductive system. Gamma-amino butyric acid (GABA) is well known as an important inhibitory neurotransmitter in the vertebrate central nervous system; however, it has been demonstrated that some endocrine organs such as somatotrophs of the anterior pituitary (1-3) and pancreatic islet cells (4-6) can also locally synthesize GABA and GABA receptors are present in these areas. Ritta et al. evaluated the effect of GABA on "in vitro" androgen production by rat testes and observed that the highest concentration of GABA was able to modify the basal and hCG-stimulated androgen production from adult and pubertal testes (7). A local GABAergic system has been recognized in adult Leydig cells in rodent and

human testes (8). Naumenko et al. demonstrated that excitation of GABA_A receptors leads to the suppression, while excitation of the GABA_B receptors leads to the intensification, of the compensatory rise in the peripheral blood level of testosterone, after its decrease induced by hemi castration (9).

Extremely low frequency electromagnetic fields (ELF-EMFs) are present wherever electric power is used. They are emitted by power lines, televisions, hair driers, cellular phones, etc (10). The possible health effects of ELF-EMFs on reproduction have been extensively studied; however, the results are often inconsistent and contradictory (11-14). These controversies are present when the testosterone levels are assayed in animals exposed to ELF-EMFs; while some authors observed no significant change in testosterone levels in rodents (14-16); others have reported an appreciable decrease of this hormone

Implication for health policy/practice/research/medical education:

All the procedures involving animals were reviewed and approved by the Institutional Research Ethics Committee of the School of Veterinary Medicine of Shiraz University.

Copyright © 2013, Research Institute For Endocrine Sciences and Iran Endocrine Society; Published by Kowsar. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

level when rats (17, 18) or guinea pigs (19) were exposed to these magnetic fields. Interactive effects of ELF-EMFs and GABA_A receptors for modulating the testosterone producing function of testicular Leydig cells have not been clarified yet.

2. Objectives

Since the peripheral GABAergic system can modulate steroidogenesis in testis, and EMF exposure can modulate testosterone synthesis in testis; so the aim of the present investigation was to study the interactive effect of GABA_A receptors and ELF-EMF (50 Hz, 0.5 mT) on serum testosterone level of male rats.

3. Patients and Methods

3.1. Animals and Experimental Design

Fifty adult male Sprague-Dawley rats with a mean body weight of 200g were randomly assigned into 10 experimental groups (n=5 each). Groups 2, 4, 6, 8, and 10 were exposed to 50 Hz, 0.5 mT ELF-EMF for 30 days 8 hrs per day while the remaining groups (1,3,5,7,9) were sham exposed animals. At the end of this period, animals in groups 1 and 2 received normal saline; while, animals in groups 3 and 4 were treated with 1 mg/kg (low dose) of bicuculline methiodide (Sigma), and for animals of groups 5 and 6 3 mg/kg (high dose) of bicuculline methiodide was injected. Animals of groups 7 and 8 were treated with 0.5 mg/kg (low dose) of muscimol hydrobromide (sigma) and rats in groups 9 and 10 received 2 mg/kg (high dose) muscimol hydrobromide (20, 21). All these treatments were performed by IP injections and the volume of injection was kept equal for rats with same weights.

During the experiment animals had free access to commercial pellets and tap water. The environmental conditions included 12 hrs light/12hrs dark cycles and temperature of about 22°C. All procedures performed in this study were in accordance with the institutional guidelines of School of Veterinary Medicine, Shiraz University for using laboratory animals in scientific experiments.

3.2. The Exposure System

The magnetic field chamber used in the present study consisted of a 70×120 cm wooden cage with 30 cm height. Three coils of electrically insulated 1mm copper wire with 200 turns each were wound around the outer surface at equal distance. The coils were connected in parallel and sealed with adhesive bandage. The electrical

source was an autotransformer with the input of 50 Hz and 220 V (22). The magnetic field inside the chamber was measured at different locations using a hand held Gauss/Tesla Meter. The field was homogenous in a zone with 21 cm distance from the transverse borders and 9 cm from the longitudinal borders. Cages were located inside this zone. The strength of the ELF-EMF was 0.5 mT in the homogenous zone.

3.3. Testosterone Assay

About forty minutes after the injection, animals were anesthetized by chloroform, and blood samples were collected by cardiocentesis. Serum testosterone level was assayed using an RIA kit (Immunotech, France).

3.4. Data Analysis

All data was presented as mean±SEM. For multiple comparisons among different groups, one-way ANOVA method, and Tukey's multiple comparison test as the post hoc were used. For testing between-subjects effects, factorial ANOVA analysis of variance was performed. $P < 0.05$ was considered as the significance level.

4. Results


4.1. Comparisons Among Different Groups

Administration of muscimol hydrobromide at both low and high doses to sham exposed rats significantly decreased serum testosterone level as compared to sham exposed animals which received saline ($P = 0.012$ and $P = 0.032$ respectively). Although a slight increase was observed in rats treated with high dose bicuculline, administration of this agent without exposure to ELF-EMF, had no significant effect on testosterone level as compared to group 1 (Figure 1). Serum testosterone levels of rats in different groups, exposed to ELF-EMF were statistically the same. Results are summarized in Figure 2. Serum testosterone level of rats in group 2 (exposed to ELF-EMF and treated with saline) was only slightly lower than sham exposed animals which received saline (group 1). Moreover, serum testosterone of exposed and sham exposed rats in each drug treatment (bicuculline methiodide or muscimol hydrobromide) showed no significant difference ($P > 0.05$) (Figure 3).

4.2. Between-Subjects Effects


No interactive effects were observed in drug×ELF-EMF, dose×ELF-EMF or drug×dose×ELF-EMF ($P > 0.05$).

Figure 1. Serum Testosterone Level of Sham Exposed Rats (Mean and SEM) in Different Groups.




* The asterisk demonstrates significant difference with sham exposed + saline group ($P < 0.05$).

Figure 2. Serum Testosterone Level of Exposed Rats (Mean and SEM) in Different Groups



No significant difference was observed among groups ($P > 0.05$).

Figure 3. Serum Testosterone Level of Exposed and Sham Exposed Rats (Mean and SEM) With Different Treatments (Saline, Low Bicuculline, High Bicuculline, Low Muscimol Hydrobromide, and High Muscimol Hydrobromide).



No significant difference was observed in each treatment group ($P > 0.05$).

5. Discussion

In the present study, the plausible interaction between $GABA_A$ receptors function and ELF-EMF on serum testosterone levels of rats has been investigated.

GABA may affect testosterone level by both peripheral and central pathways. Adult Leydig cells possess GABA synthetic enzyme, as well as $GABA_A$ and $GABA_B$ receptors, which indicates that GABA may have a role in regulation of the functions of Leydig cells including testosterone production (8). Amikishieva et al. demonstrated that in unilateral hemicastrated rats, the maximal contribution of GABAergic mechanisms in the testosterone feedback regulation involves the GABA effect via the central $GABA_A$ receptors of the medio basal hypothalamus' serotonergic neurons. Consistent with these findings, we observed that intraperitoneal administration of muscimol hydrobromide as a $GABA_A$ receptor agonist significantly reduced testosterone level. Bicuculline methiodide did not appreciably affect testosterone level in our study, although a slight increase was observed when high dose of bicuculline was administered (23). Taherianfard and Ahamdi reported that peripheral administration of bicuculline to rats only at a very high dose (3 mg/kg) is able to lower serum testosterone level; this may describe the absence of an appreciable response to bicuculline methiodide in our study (24).

In the present study, exposure of animals to 50 Hz, 0.5 mT ELF-EMF for 30 days had no appreciable effect on testosterone level as compared to sham exposed animals, although a slight decrease was observed. Al-Akhras et al. exposed rats to 50 Hz, 25 μ T ELF-EMF for 18 weeks and observed a significant decrease in serum testosterone level of rats only after 6 and 12 weeks of exposure (17). Mostafa et al. assayed serum testosterone levels of rats exposed to 50 Hz, 10 mT ELF-EMF for a period of 1, 2 and 4 weeks. At the end of 1 week no significant change was observed in testosterone level; however, this parameter was decreased significantly at the end of the second week as compared to sham exposed rats. Interestingly a remarkable increase was observed in testosterone level at the end of 4th week as compared to the second week of exposure, although it was still significantly lower than the control group. It seems that the strength of the field and/or duration of exposure may play a role in the final outcome, and this explains the discrepancies observed in different studies. The exact mechanism by which ELF-EMFs may affect serum testosterone level has not been clarified yet (18). Some authors have reported a significant increase in serum LH level accompanied by a decrease in testosterone level of rats exposed to ELF-EMFs (17, 18). This indicates that the effect of these ELF-EMFs on testosterone level is more suspected to be peripherally rather than central inhibition of hypophysal hormones release.

We observed no interactive effects in drug \times ELF-EMF, dose \times ELF-EMF or drug \times dose \times ELF-EMF in rats treated

with agents acting on GABA_A receptors and exposed to ELF-EMF. This supports the speculation that both peripheral and central GABA_A receptors do not modulate the effect of ELF-EMF on testosterone level.

In conclusion, no interactivity is present in modulatory effects of GABA_A receptors and ELF-EMFs on serum testosterone of male rats.

Acknowledgements

This study was financially supported by the Vice-chancellery of Research of Shiraz University. Thanks are due to Dr. M. Ansari Lari for her kind help in data analysis.

Authors' Contribution

None declared.

Financial Disclosure

All the authors can confirm that there is no financial or other relationship that would cause a conflict of interest.

Funding/Support

None declared.

References

- Gamel-Didelon K, Corsi C, Pepeu G, Jung H, Gratzl M, Mayerhofer A. An autocrine role for pituitary GABA: activation of GABA-B receptors and regulation of growth hormone levels. *Neuroendocrinology*. 2002;**76**(3):170-7.
- Gamel-Didelon K, Kunz L, Fohr KJ, Gratzl M, Mayerhofer A. Molecular and physiological evidence for functional gamma-aminobutyric acid (GABA)-C receptors in growth hormone-secreting cells. *J Biol Chem*. 2003;**278**(22):20192-5.
- Mayerhofer A, Hohne-Zell B, Gamel-Didelon K, Jung H, Redecker P, Grube D, et al. Gamma-aminobutyric acid (GABA): a para- and/or autocrine hormone in the pituitary. *FASEB J*. 2001;**15**(6):1089-91.
- Gilon P, Bertrand G, Loubatieres-Mariani MM, Remacle C, Henquin JC. The influence of gamma-aminobutyric acid on hormone release by the mouse and rat endocrine pancreas. *Endocrinology*. 1991;**129**(5):2521-9.
- Rorsman P, Berggren PO, Bokvist K, Ericson H, Mohler H, Ostenson CG, et al. Glucose-inhibition of glucagon secretion involves activation of GABAA-receptor chloride channels. *Nature*. 1989;**341**(6239):233-6.
- Satin LS, Kinard TA. Neurotransmitters and their receptors in the islets of Langerhans of the pancreas: what messages do acetylcholine, glutamate, and GABA transmit? *Endocrine*. 1998;**8**(3):213-23.
- Ritta MN, Campos MB, Calandra RS. Coexistence of gamma-aminobutyric acid type A and type B receptors in testicular interstitial cells. *J Neurochem*. 1991;**56**(4):1236-40.
- Geigerseder C, Doepner R, Thalhammer A, Frungieri MB, Gamel-Didelon K, Calandra RS, et al. Evidence for a GABAergic system in rodent and human testis: local GABA production and GABA receptors. *Neuroendocrinology*. 2003;**77**(5):314-23.
- Naumenko EV, Zhukova AV, Serova LI. Participation of gamma-aminobutyric acid in the negative feedback mechanisms of the hypothalamohypophyseotesticular complex. *Neurosci Behav Physiol*. 1996;**26**(2):133-6.
- Karasek M, Woldanska-Okonska M. Electromagnetic fields and human endocrine system. *ScientificWorldJournal*. 2004;**4** Suppl 2:23-8.
- Kato M, Honma K, Shigemitsu T, Shiga Y. Circularly polarized, sinusoidal, 50 Hz magnetic field exposure does not influence plasma testosterone levels of rats. *Bioelectromagnetics*. 1994;**15**(6):513-8.
- Margonato V, Veicsteinas A, Conti R, Nicolini P, Cerretelli P. Biologic effects of prolonged exposure to ELF electromagnetic fields in rats. I. 50 Hz electric fields. *Bioelectromagnetics*. 1993;**14**(5):479-93.
- Marino AA, Wolcott RM, Chervenak R, Jourdeuil F, Nilsen E, Fritel C, 2nd, et al. Coincident nonlinear changes in the endocrine and immune systems due to low-frequency magnetic fields. *Neuroimmunomodulation*. 2001;**9**(2):65-77.
- Mostafa RM, Moustafa YM, Ali FM, Shafik A. Sex hormone status in male rats after exposure to 50 Hz, 5 mTesla magnetic field. *Arch Androl*. 2006;**52**(5):363-9.
- de Bruyn L, de Jager L. Effect of long-term exposure to a randomly varied 50 Hz power frequency magnetic field on the fertility of the mouse. *Electromagn Biol Med*. 2010;**29**(1-2):52-61.
- Kim YW, Kim HS, Lee JS, Kim YJ, Lee SK, Seo JN, et al. Effects of 60 Hz 14 microT magnetic field on the apoptosis of testicular germ cell in mice. *Bioelectromagnetics*. 2009;**30**(1):66-72.
- Al-Akhras MA, Darmani H, Elbetieha A. Influence of 50 Hz magnetic field on sex hormones and other fertility parameters of adult male rats. *Bioelectromagnetics*. 2006;**27**(2):127-31.
- Mostafa RM, El Hefnawi AL, Moustafa KM, Ali FM, Moustafa YM, Kamal S, et al. Effect of 50 Hz, 10 mTesla magnetic field on sex hormones level in male rats. *J Med Sci Res*. 2007;**1**:31-36.
- Farkhad SA, Zare S, Hayatgeibi H, Qadiri A. Effects of extremely low frequency electromagnetic fields on testes in guinea pig. *Pak J Biol Sci*. 2007;**10**(24):4519-22.
- Ochi R, Suemaru K, Watanabe S, Yamaguchi T, Takechi K, Kawasaka H, et al. Effect of heat exposure on aminophylline-induced convulsions in mice. *Biol Pharm Bull*. 2011;**34**(5):666-70.
- Zarrindast MR, Ghadimi M, Ramezani-Tehrani B, Sahebgharani M. Effect of GABA receptor agonists or antagonists on morphine-induced Straub tail in mice. *Int J Neurosci*. 2006;**116**(8):963-73.
- Eleuteri AM, Amici M, Bonfili L, Cecarini V, Cuccioloni M, Grimaldi S, et al. 50Hz Extremely Low Frequency Electromagnetic Fields Enhance Protein Carbonyl Groups Content in Cancer Cells: Effects on Proteasomal Systems. *J Biomed Biotechnol*. 2009;**2009**.
- Amikishieva AV. Testosterone and behavior: involvement of the hormone in psychotropic effects of baclofen. *Bull Exp Biol Med*. 2007;**143**(2):259-63.
- Taherianfard M, Ahmadi G. Role of GABAergic system on serum testosterone and K⁺ in male rat. *J School Vet Med Shahid Chamran Univ*. 2003;**8**:35-44.

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی

توجه: بررسی مقاله ای متون (مقدماتی)

کارگاه آنلاین
بررسی مقابله ای متون (مقدماتی)

PROPOSAL
پروپوزال

توجه: پروپوزال نویسی و پایان نامه نویسی

کارگاه آنلاین
پروپوزال نویسی و پایان نامه نویسی

ISI
Scopus

توجه: آشنایی با پایگاه های اطلاعات علمی بین المللی و ترند های جستجو

کارگاه آنلاین آشنایی با پایگاه های اطلاعات علمی بین المللی و ترند های جستجو