

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی

کارگاه آنلاین
بررسی مقابله ای متون (مقدماتی)

کارگاه آنلاین
پروپوزال نویسی و پایان نامه نویسی

کارگاه آنلاین آشنایی با پایگاه های اطلاعات علمی بین المللی و ترند های جستجو

Case Report

Cerebellar Malaria Due to *Plasmodium vivax* in a Child

* AM Taksande, KY Vilhekar

Dept. of Pediatrics, Mahatma Gandhi Institute of Medical Sciences, Sevagram, Wardha, Maharashtra, India

(Received 25 Nov 2007; accepted 27 Feb 2008-)

Abstract

Infection resulting from *Plasmodium vivax* is the commonest type found in India as well as other tropical regions. The clinical picture is variable depending on the nature, immunity of the child, and lives in an endemic area or not. Here, we report a child, who presented with cerebellar dysfunction with intact sensorium and *Plasmodium vivax* was positive on peripheral smear.

Keyword: Plasmodium vivax, Cerebellar, Malaria, India

Introduction

Malaria is an acute or chronic illness characterized by paroxysms of fever, chills, sweats, fatigue, anemia and splenomegaly. Cerebellar ataxias, extrapyramidal rigidity and cranial nerve palsies presenting with intact sensorium are rare focal neurological deficits reported in malaria (1). Here, we report a child, who presented with cerebellar dysfunction with intact sensorium; *Plasmodium vivax* was present on peripheral blood smear and responded very well to antimalarial treatment.

Case Report

A 10-year-old male child from a malaria endemic area was admitted with high grade, intermittent type of fever since 3 days earlier. The child had difficulty in walking, unsteadiness of gait and tremors from the Second day of illness. He had no history of convulsion, unconsciousness, & bladder or bowel incontinence. On examination, he was fully conscious and well oriented. His vital signs were stable. He had mild pallor with

no icterus or cyanosis. Central nervous system examination revealed ataxia and bilateral cerebellar signs. Cranial nerves were normal. On motor examination, he had normal tone with grade 5 power. Sensory system examination was unremarkable. Bilateral planter reflex was flexor. Per abdominal examination shows mild splenomegaly. Fundus was normal and no sign of meningeal irritation was seen. On investigation, hemoglobin: 9gm/dl, TLC: 6,000/cumm (52% polymorphs, 45% lymphocytes, 2% eosinophils, 1% monocytes), peripheral blood smear shows normocytic and normochromic RBC with ring stage of *P. vivax*. Rapid antigen test (Paracheck HRP-II Antigen detection kit) for *P. falciparum* was negative. Serum urea and electrolyte was normal. No abnormality was detected on chest x-ray. Computed tomography scan of the brain and cerebrospinal fluid examination were normal. The child was put on tab. chloroquine and he responded very well to treatment. Cerebellar signs disappeared by the 3rd day of treatment. On third day, child was put on primaquine tablet was cut for 7 days. After one week on follow-up, there was no fever and cerebellar signs.

Discussion

Malaria is one of the most common infectious diseases and an enormous public-health problem. The disease is caused by protozoan parasites of the genus *Plasmodium*. The most serious forms of the disease are caused by *P. falciparum* and *P. vivax*. Malaria afflicts more than 500 million people, causing from 1.7 million to 2.5 million deaths each year (2). *P. vivax* malaria generally is less severe than *P. falciparum* malaria but may cause death from ruptured spleen as in association with reticulocytosis after anemia. Cerebellar involvement is the most consistent neurological manifestation of complicated as well as of uncomplicated malaria. Purkinje cells are susceptible to damage due to hyperpyrexia. The patients of uncomplicated malaria can also develop cerebellar syndrome (3). Dominant cerebellar involvement could be part of cerebral malaria. As early as 1909, Deaderic reported cerebellar involvement in a case of malaria. Later Ringdon *et al.* reported definite involvement of cerebellum in patients who died of cerebral malaria as well as in experimental animals (3).

The diagnosis of cerebellar malaria can be made based on following features: a) isolated cerebellar symptoms of acute onset associated with fever b) presence of malarial parasite in the peripheral blood c) response to antimalarial drugs. Senanayake *et al* (4) reported the clinical features of delayed cerebellar ataxia following falciparum malaria in patients whereas Koibuchi (5) reported acute disseminated encephalomyelitis following *P. vivax* malaria. The high index of suspicion is rewarding most of the times because the disease has bizarre manifestations especially in young children, septicemia and encephalitis illnesses predominate. Immunopathologic events in patients with malaria include polyclonal activation resulting in hypergammaglobulinemia and formation of immune complexes, immunodepression, and release of cytokines. Cytoadherence of the infected erythrocyte to vascular endothelium occur in *P. falciparum* malaria (6). It may

lead to obstruction of blood flow and capillary damage with resultant vascular leakage of protein and fluid, edema, and tissue anoxia in the brain, heart, lung, intestine and kidney. Severe gait and truncal ataxia are striking features suggesting that the disease predominantly affects midline cerebellar structures (7, 8). The majority of patients have afebrile period before the onset of cerebellar symptoms. It is always associated with *P. falciparum* infection very rarely reported with *P. vivax* malaria (9).

Acknowledgements

The authors declare that they have no Conflict of Interests.

References

1. Kalita J, Dhanuka AK, Misra UK. Cerebellar ataxia in patients with cerebral malaria. *Neurol India*. 1996;44:227-228.
2. Elizabeth DB. Malaria In: *Textbook of Pediatric Infections Disease*. Ed. Feiqin RD, Demmler GJ, Gherry JD, Kaplan SL. Barnett ED. Saunders, Philadelphia 5th ed; 2004. p. 2714-2715.
3. Chitkara AJ, Anand NK, Saini L. Cerebellar Syndrome in malaria. *Indian Pediatr*. 1984; 21:908-919.
4. Senanayake N, de Silva HJ. Delayed cerebellar ataxia complicating falciparum malaria: A clinical study of 74 patients. *Journal of Neurology*. 1994;241:456-459.
5. Koibuchi T, Nakamura T, Miura T, Endo T, Nakamura H, Takahashi T, *et al.* Acute disseminated encephalomyelitis following *Plasmodium vivax* malaria. *J Infect Chemother*. 2003;9(3):254-6.
6. Krause PJ. Malaria (*Plasmodium*). In: *Nelson Textbook of pediatrics* 17th ed. Eds. Behrman RE, Kliegman RM, Nelson VC, Vaughan VC. Philadelphia, W. B. Saunders company; 2001. p. 1139-1142.

7. Phillips RE, Solomon T. Cerebral Malaria in children. *Lancet*. 1990;336:1355-1360.
8. Brewster DR, Kwiatkowski D, White NJ. Neurological sequelae of cerebral malaria in children. *Lancet*. 1990;336:1039-1043.
9. Taksande B, Jajoo U, Jajoo M. Cerebral malaria: A rare manifestation of *P. vivax*. *The internet J of Neurology*. 2001.

Archive of SID

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی

توجه: بررسی مقاله ای متون (مقدماتی)

کارگاه آنلاین
بررسی مقابله ای متون (مقدماتی)

PROPOSAL
پروپوزال

توجه: پروپوزال نویسی و پایان نامه نویسی

کارگاه آنلاین
پروپوزال نویسی و پایان نامه نویسی

ISI
Scopus

توجه: آشنایی با پایگاه های اطلاعات علمی بین المللی و ترند های جستجو

کارگاه آنلاین آشنایی با پایگاه های اطلاعات علمی بین المللی و ترند های جستجو