

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



عضویت در خبرنامه



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



مباحث پیشرفته یادگیری عمیق؛ شبکه های توجه گرافی (GAN)

مباحث پیشرفته یادگیری عمیق؛
شبکه های توجه گرافی
(Graph Attention Networks)



آموزش استفاده از وب آو ساینس

کارگاه آنلاین آموزش استفاده از
وب آو ساینس



کارگاه آنلاین مقاله روزمره انگلیسی

Letter

Posterior Scleral Thinning Accompanies Increased Vitreous Chamber Depth in Myopia

Abdullah Kaya¹, MD; Yildirim Yildirim², MD

¹Department of Ophthalmology, Anittepe Military Dispansery, Ankara, Turkey

²Department of Ophthalmology, GATA Haydarpaşa Training Hospital, Istanbul, Turkey

J Ophthalmic Vis Res 2016; 11 (2): 242.

Sir,

We read with great interest the article titled "Association between refractive errors and ocular biometry in Iranian adults" by Hashemi et al^[1] The authors aimed to evaluate the association between refractive errors and ocular biometry. We appreciate their valuable study and would like to make a comment.

In the mentioned study, a strong correlation was reported between vitreous chamber depth (VCD) and myopia. As the authors stated, this finding indicates the predominant role of the posterior segment in myopization. Similarly, a recent study conducted by Shen et al supported this idea and compared scleral thickness in secondary and primary high myopia.^[2] In high axial myopia secondary to congenital glaucoma, the sclera was found to be thinner both anterior and posterior to the equator; whereas in primary high axial myopia, scleral thinning was predominantly found posterior to the equator. Generalized scleral thinning secondary to congenital glaucoma shows the effect of increased intraocular pressure on the scleral wall. However, limited scleral thinning posterior to the equator in primary myopia indicates the predominant role of the posterior segment in myopization.

We congratulate authors for their findings regarding the relationship between VCD and myopia. Besides this

finding, the recently reported scleral thinning posterior to the equator may lead to new investigations.

Financial Support and Sponsorship

Nil.

Conflicts of Interest

There are no conflicts of interest.

REFERENCES

1. Hashemi H, Khabazkhoob M, Emamian MH, Shariati M, Miraftab M, Yekta A, et al. Association between refractive errors and ocular biometry in Iranian adults. *J Ophthalmic Vis Res* 2015;10:214-220.
2. Shen L, You QS, Xu X, Gao F, Zhang Z, Li B, et al. Scleral and choroidal thickness in secondary high axial myopia. *Retina* 2016. [Epub ahead of print]. doi: 10.1097/IAE.0000000000000947.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online

Quick Response Code:



Website:

www.jovr.org

DOI:

10.4103/2008-322X.183915

Correspondence to:

Abdullah Kaya, MD. Department of Ophthalmology, Anittepe Military Dispansery, Yucetepe Mh. Genclik Cad. 88. Sok. Çankaya, Ankara 06280, Turkey.
E-mail: abdullahkayamd@gmail.com

Received: 14-01-2016

Accepted: 13-05-2016

How to cite this article: Kaya A, Yildirim Y. Posterior scleral thinning accompanies increased vitreous chamber depth in myopia. *J Ophthalmic Vis Res* 2016;11:242.

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



عضویت در خبرنامه



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



مباحث پیشرفته یادگیری عمیق؛
شبکه های توجه گرافی
(Graph Attention Networks)



کارگاه آنلاین آموزش استفاده از
وب آوساینس



کارگاه آنلاین مقاله روزمره انگلیسی