

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی



مقاله نویسی علوم انسانی

مقاله نویسی علوم انسانی



اصول تنظیم قراردادها

اصول تنظیم قراردادها



آموزش مهارت های کاربردی در تدوین و چاپ مقاله

آموزش مهارت های کاربردی در تدوین و چاپ مقاله

A Potential Method for Remediation of Cadmium Pollution in Aquatic Medium by Hydrophyte, *Ceratophyllum demersum* L.

A. Parnian^{1*} - M. Chorom² - N. Jaafarzadeh³

Received: 13-08-2013

Accepted: 01-08-2016

Introduction: With increasing of population and the valuable water resource pollutions, a demand has been felt for new and inexpensive methods in order to remediation and improving of water quality. Cadmium is a trace element. In low concentration, this heavy metal is harmful to life, and considered as a dangerous pollutant. Cadmium leads to pollution and reduction of water quality; sometimes even toxicity through contaminated sources such as wastewater (Agricultural, municipal and industrial). Phytoremediation with aquatic macrophytes is an effective and inexpensive method for improving water quality and wastewater. The aim of this study was to investigate the cadmium phytoremediation by *Ceratophyllum demersum* L. as a potential method for remediation of cadmium pollution in aquatic medium.

Materials and Methods: In this study, the remediation of cadmium pollution in aquatic medium monitored, within 14 days cultivation of coontail (*Ceratophyllum demersum* L.). At first, for estimating the level of local wastewater cadmium pollutions, five-month cadmium concentration measurement of steel industrial wastewater and urban wastewater set. Then, plants collected from the irrigation channels of Shahid Chamran University of Ahvaz. After finding the best pH of nutrient solution for *Ceratophyllum demersum* L. growth by cultivating the plants in 2 liters pots filled by the solutions with three different pH (5.5, 7 and 9.5) within three weeks; 12 grams of plants cultivated in 2 liters of Hoagland nutrient solution contaminated by cadmium (pH = 7). The initial contamination levels were set as five different concentrations of cadmium (0, 1, 2, 4, and 6 mg l⁻¹) with three replications. The cadmium concentrations of the pots were measured every day and on the last day of cultivation, plants wet weight, plants dry weight and Cd concentration in plants were measured. Then, biomass production, Cd bioconcentration factor (BCF), Cd uptake index, and Cd uptake percentage of plants were calculated. Standard deviations calculation and correlation and regression analysis were performed using Microsoft Office Excel 2007 and SPSS 16. One-way ANOVA performed to identify significant differences in metal concentrations in the different treatments. Differences considered significant at $p < 0.05$.

Results and Discussion: Among three pH (5.5, 7 and 9.5) for plants cultivation, *C. demersum* L. grew better in pH = 7. In fact, the average amount of produced biomasses were 46.6 g (pH = 5.5), 79.6 g (pH = 7) and 68.4 g (pH = 9.5). Therefore, to investigate the Cd remediation, the pH of nutrient solution set equal to 7. The final Cd concentrations in nutrient solution for initial Cd concentrations of 1, 2, 4 and 6 mg l⁻¹ were 0.30, 0.36, 2.76 and 3.85 mg l⁻¹ respectively. Moreover, the Cd uptake percentage after 14 days cultivation of *C. demersum* L. in nutrient solution for initial Cd concentrations of 1, 2, 4 and 6 mg l⁻¹ were 70.00, 82.01, 31.00 and 35.83 % respectively. Cd uptake percentage of plants for initial concentrations of 4 and 6 mg l⁻¹ were significantly lesser than those of 1 and 2 mg l⁻¹. The decreased uptake efficiency percentage maybe caused by the effect of Cd toxicity on plant cell membrane permeability and efficiency. The average of BCF in plants for initial Cd concentrations of 1, 2, 4 and 6 mg l⁻¹ were 384.4, 707.9, 66.5 and 75.0 respectively. High reduction of BCF amounts with increasing the initial concentration of 2 to 4 and 6 mg l⁻¹, maybe caused by cadmium physiological adverse effects on plants. The averages of uptake index in plants were 1.26, 2.95, 2.24 and 3.92 mg for initial Cd concentrations of 1, 2, 4 and 6 mg l⁻¹ respectively. The results showed a reduction between 2 and 4 mg l⁻¹ concentrations that probably caused by Cd toxicity disruption on plants uptake mechanism and growth. Moreover, the increase of plants uptake index in initial concentration of 6 mg l⁻¹ could be explain by partial losing of the selective permeability of the plants cell membrane. The maximum (3.60 g/day) and minimum (1.62 g/day) of biomass production related to pollutant concentrations of 0 and 6 mg l⁻¹ respectively, and it shows a great effect of the Cd on *C. demersum* L. growth.

Conclusion: The plant accumulated cadmium efficiently, and the remediation efficiency was near to 82%. However, the pollutant removal was not complete in a short time. In total, phytoremediation of cadmium and other pollutants from wastewater or other aqueous solutions by *Ceratophyllum demersum*, as a native aquatic

1- Ph.D. of Soil Science, Young Researchers and Elite Club, Ahvaz Branch, Islamic Azad University, Ahvaz, Iran

(* - Corresponding Author Email: amir.parnian86@gmail.com)

2- Associate Professor, Faculty of Agriculture, Department of Soil Science, Shahid Chamran University, Ahvaz, Iran

3- Professor, Environmental Technologies Research Center, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran

plant of most of Iran's rivers, could be an efficient and appropriate method.

Keywords: Aquatic macrophyte, Cadmium, *Ceratophyllum demersum* L., Heavy metals, Wastewater phytoremediation

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی



مقاله نویسی علوم انسانی



اصول تنظیم قراردادها



آموزش مهارت های کاربردی در تدوین و چاپ مقاله