

Comparison of Berlin Questionnaire, STOP-Bang, and Epworth Sleepiness Scale for Diagnosing Obstructive Sleep Apnea in Persian Patients

Abstract

Background: Obstructive sleep apnea (OSA) is a common disorder characterized by snoring, daytime sleepiness, fatigue, and repeated termination of airflow (apnea). Standard polysomnography (PSG) is diagnostic gold standard for OSA. PSG is expensive and not available everywhere. To identify the best OSA questionnaire for screening in Persian population, we compared Berlin, STOP-BANG, and Epworth Sleepiness Scale (ESS). **Methods:** In a cross-sectional study conducted on 400 adult patients suspected of OSA in Bamdad Respiratory Research Center, patients completed three questionnaires. For each questionnaire, patients were divided into high risk and low risk. Then, PSG was performed for all patients. According to PSG, patients categorized into without OSA (apnea-hypopnea index [AHI] <5), mild OSA (15 > AHI ≥ 5), moderate OSA (30 > AHI ≥ 15), and severe OSA (AHI ≥ 30). Based on questionnaires and PSG results, predictive parameters for screening tests were calculated. **Results:** There were 234 (58.5%) males and 166 (41.5%) females in study population. Mean age of patients was 49.29 ± 9.75 standard deviation years. There was significant association between age with OSA ($P = 0.005$) and between body mass index (BMI) and neck circumference, with moderate and severe OSA ($P < 0.001$). Sensitivities of Berlin, STOP-BANG, and ESS were 86.42%, 81.46%, and 59%, respectively. Specificities of Berlin, STOP-BANG, and ESS were 52.94%, 82.35%, and 76.47%, respectively. **Conclusions:** This study suggested that Berlin and STOP-BANG are more sensitive and accurate than ESS for OSA screening in Iran.

Keywords: Diagnosis, obstructive sleep apnea, Persia, surveys and questionnaires

Introduction

Obstructive sleep apnea (OSA) is a common disorder that characterized by snoring, daytime sleepiness, fatigue, repeated termination of airflow (apnea), and hypoxemia.^[1] OSA is an independent risk factor for cardiovascular diseases including hypertension, coronary artery disease, stroke, and pulmonary hypertension.^[2] The prevalence of disease varies in different countries.^[1,3-6] The gold standard diagnostic method is polysomnography (PSG).^[2] PSG is a noninvasive technique that involves overnight monitoring of several physiological variables including electroencephalography, eye movements, muscle tone as well as respiratory effort, airflow, and oxygen saturation. An apnea is defined as the complete cessation of airflow for a minimum of 10 s. The definition of a hypopnea includes a reduction of airflow that is associated with either an oxygen desaturation (of at least 3% or 4%) or an arousal. OSA is diagnosed when the apnea-hypopnea index (AHI), i.e. the total number

of obstructive apneas and hypopneas per hour of sleep is >5. The severity of OSA is graded according to commonly used clinical criteria as mild (AHI ≥ 5 but <15), moderate (AHI ≥ 15 but <30), or severe (AHI ≥ 30).^[7] Considering high cost and inaccessibility of PSG, several screening questionnaires have been developed. The most popular questionnaires are Berlin questionnaire (BQ), STOP-BANG, and Epworth Sleepiness Scale (ESS). These questionnaires had wide range of sensitivity and specificity in different races.^[8-13] Hence, we conducted this study to compare the predictive values of these questionnaires in Persian population.

Methods

This was a cross-sectional study. We enrolled adult patients who visited Sleep Department of Bamdad Respiratory Research Center, for evaluating suspected OSA between May 2015 and June 2016. Inclusion criteria were as follows: age over 18 years, OSA

Babak Amra,
Maedeh Javani¹,
Forogh
Soltaninejad²,
Thomas Penzel³,
Ingo Fietze⁴,
Christoph Schoebel⁵,
Ziba Farajzadegan⁶

Department of Medicine, Pulmonary Unit, Isfahan University of Medical Sciences, Bamdad Respiratory Research Center, Isfahan, Iran, ¹Department of Internal Medicine, Isfahan University of Medical Sciences, Isfahan, Iran, ²Department of Medicine, Pulmonary Unit, Isfahan University of Medical Sciences, Isfahan, Iran, ³Center of Sleep Medicine, Charité – Universitätsmedizin Berlin, Berlin, Germany, ⁴Department of Cardiology and Pulmonology, Center of Sleep Medicine, Charité – Universitätsmedizin Berlin, Berlin, Germany, ⁵Department of Cardiology and Angiology, Center of Sleep Medicine, Charité – Universitätsmedizin Berlin, Berlin, Germany, ⁶Department of Community and Preventive Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Address for correspondence:

Dr. Forogh Soltaninejad,
Department of Medicine,
Pulmonary Unit, Isfahan
University of Medical Sciences,
Isfahan, Iran.
E-mail: soltaninejad.fg@gmail.com

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.ir

DOI:
10.4103/ijpvm.IJPVM_131_17

Quick Response Code:



This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Amra B, Javani M, Soltaninejad F, Penzel T, Fietze I, Schoebel C, et al. Comparison of Berlin questionnaire, STOP-Bang, and Epworth Sleepiness Scale for diagnosing obstructive sleep apnea in Persian patients. *Int J Prev Med* 2018;9:28.

symptoms (fatigue during the day, nocturnal snoring, and observed apnea by others), no previous treatment, completion of questionnaires, and personal consent to do PSG and participate in the study. Patients were excluded if there was a lack of consent to continue and incompletely finished questionnaire and with a history of psychiatric disorder. Ethical Committee of Isfahan University of Medical Sciences approved the study protocol. Informed written consent was obtained from all patients before inclusion in this study. All patients filled out STOP-BANG, Berlin, and ESS questionnaires.

The STOP-BANG questionnaire includes four subjective (STOP: snoring, tiredness, observed apnea, and high blood pressure) and four demographics items (Bang: body mass index [BMI], age, neck circumference, gender). Answering yes to three or more items is categorized as high risk for OSA.^[14] STOP-BANG questionnaire validated in meta-analysis for screening of OSA in sleep clinic and surgical population.^[10] BQ that was developed in 1999 includes three sections. The first section is about snoring, the second section is about daytime fatigue and sleepiness, and the last section is about medical history and anthropometric measures such as hypertension and BMI. If two or more categories were positive, the patient is considered high risk for OSA.^[12] The ESS is an eight-item questionnaire to measure daytime sleepiness. Questionnaire has a four-point Likert response format (0–3), and the score ranges from 0 to 24. ESS score ≥ 11 indicates excessive daytime sleepiness and high risk for OSA.^[15] We used valid Persian versions of three questionnaires.^[2,16,17]

Then, we performed standard attended overnight PSG by PSG device (SOMNOmedics GmbH, Randersacker, Germany). In standard PSG, electroencephalogram, electrocardiogram, electrooculogram, chin electromyogram, oronasal airflow, oxygen saturation by pulse oximetry, thoracic, abdominal, and leg movements were recorded. Scoring was done according to the American Association of Sleep Medicine guideline 2015.^[18] Patients were classified based on the AHI values without OSA ($AHI < 5$), mild OSA ($5 \leq AHI < 15$), moderate OSA ($15 \leq AHI < 30$), and severe OSA ($AHI \geq 30$).

Those who performed PSG were not aware of results of questionnaires. PSGs were interpreted without awareness of medical histories and questionnaires results.

SPSS version 17.0 (SPSS Science, Apache Software Foundation, Chicago, IL, USA) was used. Sensitivity, specificity, positive predictive values (PPV), negative predictive values (NPV), and positive and negative likelihood ratio (LR + and LR-) were calculated using K^2 . The Spearman's correlation coefficient was used for three questionnaires in patients. Tests were two-tailed and $P < 0.05$ was accepted as statistically significant. We evaluated discriminative ability of questionnaires with receiver operating characteristic (ROC) curves.

Results

From May 2015 to June 2016, we included 400 patients in our study. There were 234 (58.5%) males and 166 (41.5%) females and mean age was 49.29 ± 9.75 years. Demographic data for quantitative and qualitative parameters were presented in Table 1. There was significant association between age with OSA ($P = 0.005$) and between BMI and neck circumference, with moderate and severe OSA ($P < 0.001$). There was no significant association between genders with OSA.

Results of three questionnaires were presented as low risk and high risk in Table 2. In PSG, frequencies of without OSA, mild OSA, moderate OSA, and severe OSA were 17 (4.25%), 56 (14%), 84 (21%), and 243 (60.75%), respectively.

The BQ had the best sensitivity (86.42), but STOP-BANG had the best specificity (82.35%). Predictive parameters of questionnaires including sensitivity, specificity, PPV, NPV, and LR + and LR - were presented in Table 3.

Area under the ROC curve (95% confidence interval) for OSA ($AHI \geq 5$) in Berlin, STOP-BANG, and ESS questionnaires was 0.76 (0.69, 0.08), 0.89 (0.82, 0.96), and 0.69 (0.58, 0.81), respectively. ROC curves for three questionnaires were illustrated in Figure 1. The Spearman's correlation coefficient between STOP-BANG and ESS, STOP-BANG and Berlin, and Berlin and ESS were 0.20, 0.44, and 0.29, respectively ($P < 0.001$).

Discussion

When evaluating the predictive values of Berlin, STOP-BANG and ESS to identify patients at risk for OSA,

Table 1: Demographic characteristics of patients

Parameter	n (%)	Mean±SD	Minimum	Maximum
Sex				
Male/ female	235 (58.74)/ 165 (41.25)	-	-	-
Age	400	49.29 (9.72)	30	60
BMI	400	32.40 (7.43)	19	60
NC	400	40.83 (3.13)	32	54

SD=Standard deviation, NC=Neck circumference, BMI=Body mass index

Table 2: Results of Berlin, STOP-BANG, and Epworth Sleepiness Scale screening questionnaires for obstructive sleep apnea (apnea-hypopnea index ≥ 5)

Questionnaire	Result	AHI	
		<5 (n)	≥ 5 (n)
Berlin	Low risk	9	52
	High risk	8	331
STOP-BANG	Low risk	14	71
	High risk	3	312
ESS	Low risk	13	157
	High risk	4	226

ESS=Epworth Sleepiness Scale, AHI=Apnea-hypopnea index

Berlin had the highest sensitivity, but STOP-BANG had the highest specificity, area under the curve (AUC), and PPV. Sensitivity and specificity of ESS were between the others. The sensitivity of ESS was especially low.

In the study of Pataka *et al.*, ESS, Berlin, STOP, STOP-BANG, and 4-V were evaluated in the general population. The STOP-BANG had the highest sensitivity, NPV, and AUC, but the least specificity. 4-V had the

highest specificity. In that study, sensitivity of STOP-BANG and Berlin was 96.2% and 84.4%, respectively. Both tests had low specificity. The sensitivity of Berlin was similar to our study. The low specificity seems to be caused by difference of populations under the study. Their sample was representative of general population.^[19]

In a systematic review performed by Abrishami *et al.*, Berlin was the second and STOP-BANG was the third most sensitive questionnaire after Wisconsin in people without a history of sleep disorders. They concluded that STOP-BANG had high quality methodological and reasonably accurate results. Similar to our study, sensitivity of Berlin was more than STOP-BANG.^[20] Du *et al.* assessed predictive value of Berlin, STOP, STOP-BANG, and ESS. STOP-BANG had the highest sensitivity, but Berlin had the highest AUC. AUC for Berlin was similar to our study.^[21]

In the study of Kim *et al.*, three questionnaires including sleep apnea of sleep disorder questionnaire, Berlin, and STOP-BANG were compared. Sensitivity of STOP-BANG was more than Berlin. They excluded people with comorbidities such as psychiatric, neurologic, and severe

Table 3: Predictive parameters for Berlin, STOP-BANG, and Epworth Sleepiness Scale screening questionnaires for obstructive sleep apnea (apnea-hypopnea index ≥ 5)

Parameter	Berlin (%)	STOP-BANG (%)	ESS (%)
Sensitivity	86.42	81.46	59
Specificity	52.94	82.35	76.47
PPV	97.64	99	98.26
NPV	14.75	16.47	7.64
LR positive	1.79	4.50	2.45
LR negative	0.01	0.23	0.53

ESS=Epworth Sleepiness Scale, PPV=Positive predictive value, NPV=Negative predictive value, LR=Likelihood ratio




Figure 1: Receiver operator characteristic curves of questionnaires (a) Epworth Sleepiness Scale, (b) Berlin, and (c) STOP-BANG for obstructive sleep apnea (apnea-hypopnea index ≥ 5)

gastrointestinal, cardiac, and malignancy. Therefore, their results cannot be generalized to the general population.^[22] In another studies of Berlin and STOP-Bang in general population, both had good predictive value for screening moderate and severe OSA.^[23,24]

In multiple studies, different predictive parameters were reported for OSA screening questionnaires. Discrepancies were caused by different sampling (community vs. sleep clinic), sample size, sample characteristics (gender, age, BMI, and culture), and AHI cutoff (mild vs. moderate to severe).

We tried to identify the best questionnaire for screening OSA in Persian population. Patients were referred to sleep clinic due to symptoms. We did not perform the study in general population regardless of symptoms. Although we did not exclude patients with comorbidities to simulate general population, our population was not an exact sample of general population. Patients without OSA (AHI <5) were few. This preselected status of population probably affected our findings. Furthermore, all patients completed Berlin first and ESS last. Because of the effect of time on accuracy of answers, this fixed sequence could be a confounding factor.

Therefore, we recommend the future study on Persian population with sampling of community and random sequences for answering to questionnaires. Accuracy of answers decreases with increasing questionnaires probably. Therefore, it is necessary to use valid questionnaires such as Berlin and STOP-BANG in one session.

Conclusions

There is a need for highly sensitive and accurate tests for screening and avoiding misses. Considering this, our study showed that BQ and STOP-Bang are more reliable and preferable than ESS for screening of OSA in Persian population.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Received: 03 Mar 17 **Accepted:** 10 Jun 17

Published: 09 Mar 18

References

- Amra B, Farajzadegan Z, Golshan M, Fietze I, Penzel T. Prevalence of sleep apnea-related symptoms in a Persian population. *Sleep Breath* 2011;15:425-9.
- Amra B, Nouranian E, Golshan M, Fietze I, Penzel T. Validation of the persian version of berlin sleep questionnaire for diagnosing obstructive sleep apnea. *Int J Prev Med* 2013;4:334-9.
- Hiestand DM, Britz P, Goldman M, Phillips B. Prevalence of symptoms and risk of sleep apnea in the US population: Results from the national sleep foundation sleep in America 2005 poll. *Chest* 2006;130:780-6.
- Kim J, In K, Kim J, You S, Kang K, Shim J, *et al.* Prevalence of sleep-disordered breathing in middle-aged Korean men and women. *Am J Respir Crit Care Med* 2004;170:1108-13.
- Peppard PE, Young T, Barnett JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol* 2013;177:1006-14.
- Udwadia ZF, Doshi AV, Lonkar SG, Singh CI. Prevalence of sleep-disordered breathing and sleep apnea in middle-aged urban Indian men. *Am J Respir Crit Care Med* 2004;169:168-73.
- Pamidi S, Aronsohn RS, Tasali E. Obstructive sleep apnea: Role in the risk and severity of diabetes. *Best Pract Res Clin Endocrinol Metab* 2010;24:703-15.
- Chiu HY, Chen PY, Chuang LP, Chen NH, Tu YK, Hsieh YJ, *et al.* Diagnostic accuracy of the Berlin questionnaire, STOP-BANG, STOP, and Epworth sleepiness scale in detecting obstructive sleep apnea: A bivariate meta-analysis. *Sleep Med Rev* 2016. pii: S1087-079230127-7.
- Khaledi-Paveh B, Khazaie H, Nasouri M, Ghadami MR, Tahmasian M. Evaluation of berlin questionnaire validity for sleep apnea risk in sleep clinic populations. *Basic Clin Neurosci* 2016;7:43-8.
- Nagappa M, Liao P, Wong J, Auckley D, Ramachandran SK, Memsoudis S, *et al.* Validation of the STOP-bang questionnaire as a screening tool for obstructive sleep apnea among different populations: A systematic review and meta-analysis. *PLoS One* 2015;10:e0143697.
- Nagappa M, Wong J, Singh M, Wong DT, Chung F. An update on the various practical applications of the STOP-Bang questionnaire in anesthesia, surgery, and perioperative medicine. *Curr Opin Anaesthesiol* 2017;30:118-25.
- Netzer NC, Stoohs RA, Netzer CM, Clark K, Strohl KP. Using the Berlin questionnaire to identify patients at risk for the sleep apnea syndrome. *Ann Intern Med* 1999;131:485-91.
- Sharma A, Molano J, Moseley BD. The STOP-BANG questionnaire improves the detection of epilepsy patients at risk for obstructive sleep apnea. *Epilepsy Res* 2017;129:37-40.
- Chung F, Yegneswaran B, Liao P, Chung SA, Vairavanathan S, Islam S, *et al.* STOP questionnaire: A tool to screen patients for obstructive sleep apnea. *Anesthesiology* 2008;108:812-21.
- Johns MW. A new method for measuring daytime sleepiness: The Epworth sleepiness scale. *Sleep* 1991;14:540-5.
- Sadeghniaat Haghghi K, Montazeri A, Khajeh Mehrizi A, Aminian O, Rahimi Golkhandan A, Saraei M, *et al.* The Epworth Sleepiness Scale: Translation and validation study of the Iranian version. *Sleep Breath* 2013;17:419-26.
- Sadeghniaat-Haghghi K, Montazeri A, Khajeh-Mehrizi A, Ghajarzadeh M, Alemohammad ZB, Aminian O, *et al.* The STOP-BANG questionnaire: Reliability and validity of the Persian version in sleep clinic population. *Qual Life Res* 2015;24:2025-30.
- Berry RB, Gamaldo CE, Harding SM, Brooks R, Lloyd RM, Vaughn BV, *et al.* AASM Scoring Manual Version 2.2 Updates: New Chapters for Scoring Infant Sleep Staging and Home Sleep Apnea Testing. *J Clin Sleep Med* 2015;11:1253-4.
- Pataka A, Daskalopoulou E, Kalamaras G, Fekete Passa K, Argyropoulou P. Evaluation of five different questionnaires for assessing sleep apnea syndrome in a sleep clinic. *Sleep Med* 2014;15:776-81.
- Abrishami A, Khajehdehi A, Chung F. A systematic review of screening questionnaires for obstructive sleep apnea. *Can J Anaesth* 2010;57:423-38.
- Du L, Li Z, Tang X. Application value of four different

- questionnaires in the screening of patients with obstructive sleep apnea. *Zhonghua Yi Xue Za Zhi* 2015;95:3407-10.
22. Kim B, Lee EM, Chung YS, Kim WS, Lee SA. The utility of three screening questionnaires for obstructive sleep apnea in a sleep clinic setting. *Yonsei Med J* 2015;56:684-90.
 23. Tan A, Yin JD, Tan LW, van Dam RM, Cheung YY, *et al.* Using the berlin questionnaire to predict obstructive sleep apnea in the general population. *J Clin Sleep Med* 2017;13:427-32.
 24. Tan A, Yin JD, Tan LW, van Dam RM, Cheung YY, Lee CH. Predicting obstructive sleep apnea using the STOP-Bang questionnaire in the general population. *Sleep Med* 2016;27-28:66-71.

Archive of SID