

Phylogenetic study based on the phosphoprotein gene of Iranian Newcastle disease viruses (NDV) isolates, 2010 -2012

Ghalyanchi Langroudi, A.^{1*}, Hosseini, H.², Karimi, V.³, Madadgar, O.¹, Hashemzadeh, M.⁴, Ghafouri, S.A.⁵, Bagheri Ghadikolaei, S.S.⁶, Vahedi, S.M.⁶

¹Department of Microbiology and Immunology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

²Department of Clinical Sciences, Faculty of Veterinary Medicine, Islamic Azad University, Karaj Branch, Karaj, Iran

³Department of Poultry Diseases, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

⁴Department of Research & Production of Veterinary and Poultry Vaccines, Razi Vaccine and Serum Research Institute, Karaj, Iran

⁵Department of Poultry Health and Diseases, Iranian Veterinary Organization, Tehran, Iran

⁶Students' Scientific Association of Microbiology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran

Key words:

Newcastle disease, phosphoprotein, phylogenetic study

Correspondence

Ghalyanchi Langeroudi, A.
Department of Microbiology and Immunology, Faculty of Veterinary Medicine, University of Tehran, Tehran, Iran
Tel: +98(21) 61117053
Fax: +98(21) 66933222
Email: ghalyana@ut.ac.ir

Received: 3 March 2014

Accepted: 12 May 2014

Introduction

Newcastle disease (ND), caused by Newcastle disease virus (NDV), is a highly contagious respiratory, enteric or neurological viral disease and an OIE notifiable devastating disease that is found in most avian species, especially in chickens (Miller et al., 2009).

The NDV is a member of avian paramyxovirus serotype-1 (APMV-1), which belongs to the genus Avulavirus and Paramyxoviridae family. The NDV has RNA genome (a negative-sense single stranded,

Abstract:

BACKGROUND: Newcastle disease virus (NDV) is the causative agent of the Newcastle disease (ND), a highly contagious disease in birds that causes significant economic losses to the poultry industry worldwide. ND is endemic in Iran and outbreaks are reported regularly in commercial poultry flocks and different species of birds. **OBJECTIVES:** The current study was carried out to characterize NDV based on phosphoprotein (P) gene from recent outbreaks in Iran, 2010-2012. **METHODS:** The P gene fragment of NDV isolates of five chickens, 1 ostrich, and 1 Pigeon paramyxovirus-1 was obtained by RT-PCR and sequenced. **RESULTS:** Phylogenetic analysis of sequences revealed that chicken and ostrich NDV isolates were closely related and placed in the genotype VII and Pigeon Paramyxovirus-1 was located in the genotype V. **CONCLUSIONS:** This is the first report of Phosphoprotein gene sequences of NDV strains isolated in Iran. This study will help us to understand the epidemiology and molecular characteristics of Newcastle disease virus in Iran.

15 Kb), which codes for an RNA-directed RNA polymerase (L), hemagglutinin-neuraminidase protein (HN), fusion protein (F), matrix protein (M), phosphoprotein (P), and nucleoprotein (NP) (Alexander et al., 1997).

All NDV isolates are classified as three pathotypes based on the disease severity: mesogenic, velogenic, and lentogenic. Genetic and antigenic analyses of NDV isolates have determined the existence of two main classes I and II, which could be divided into ten genotypes (1-10 in class I and I-X in class II): The genotypes VI and VII being further

divided into seven (VIa-VIg) and five (VIIa-VIIe) sub-genotypes respectively (Miller et al., 2010).

The phosphoprotein, or P protein, of NDV, has the critical role in replication and transcription and has multiple functions. For example, the P protein also acts as a chaperone to prevent uncontrolled encapsidation of non-viral RNA by the NP protein (Jahanshiri et al., 2005; Locke et al., 2000; Smith and Hightower, 1981). ND is endemic in Iran and we have some reports for incidence of ND in every year. In the past few decades, implementations of extensive vaccination programs in commercial poultry farms, and to some extent in small rural poultry flocks have reduced the number of epizootics outbreaks of ND in Iran (Bozorgmehri-Fard and Keyvanfar, 1979; Rezaeianzadeh et al., 2011). No data is available about the P gene of Iranian NDV isolates till now. The study is the first report on P gene characterization of Iranian NDV isolates and provides a suggestion on NDV data collection to increase future virus evolutionary analysis.

Materials and Methods

Samples: We worked on six velogenic NDV isolates and one Pigeon Paramyxovirus-1 (approved by analyses of sequences of Fusion gene) that were isolated from different outbreaks in broiler and breeder farms, ostrich and pigeon case. Ten-day-old SPF embryonated chicken eggs (allantoic cavity) were inoculated. These eggs were incubated at 37°C for up to 2 days, embryonic death was monitored, and then allantoic fluid was collected under routine conditions. All data of mentioned isolates are available in Table 1.

RNA extraction and RT-PCR: RNA is extracted from allantoic fluid through QIAamp virus spin kit (QIAGEN, USA) and stored at -70°C. Reverse transcription was done by using random hexamer with revert aid first strand cDNA synthesis Kit (Fermentas Co, Canada) (OIE, 2012). Amplification of the P gene was carried out by PCR as described by using one pair of specific primers of phosphoprotein gene (NDF: ACCAGYGGRACTGTCATHGAC, NDR: CGGACAGTGTCCYTCTCYAC). The PCR amplification was carried out in a 50 µL reaction mixture containing 0.2 mM dNTP, 0.5 mM MgCl₂, 0.2 mM each primer, 19 PCR buffer and 1 U of Taq

polymerase. Reactions were performed according to the following protocol: 95°C for 5 min, followed by 35 cycles (95°C for 45 s, 58°C for 45 s, 72°C for 1 min) and a final extension step of 10 min at 72°C.

Sequencing and bioinformatics study: The PCR products were purified with the AccuPrep[®] PCR Purification kit (Bioneer Co, South Korea) and sent to Source BioScience Company (UK) for DNA sequencing. The primary sequence analyses and edition were carried out in the CLC sequence viewer (Ver. 6.0.2). Multiple amino acid alignments were performed on P genes representative viruses using Clustal W (MEGA5) (Tamura et al., 2011) on current study isolates and other NDV isolates from Gene bank (Table 2). Phylogenetic trees were drawn from nucleotide sequences based on the sequence of phosphoprotein gene using Neighbor-Joining method with MEGA5 model with 1000 bootstrapping replications in molecular evolutionary genetics analysis (MEGA5). The structure of the NDV that were sequenced and analyzed in this study has been submitted to GeneBank under the accession number KF824513-KF824519 (Table 1).

Results

The partial sequences of phosphoprotein gene of six Iranian NDV isolates were amplified and sequenced. MEGA5 program were used to determine the sequence similarity and homology of the p gene among them. Results indicate that nucleotide homology among six NDVs isolates with P genes is between 99.6% and 99.9%. The percent identify based on amino acid is between 99.3%- 99.8%. In BLAST result NDVs isolates have high nucleotide identities (97%) with JSG0210 (JF340367), chicken/TC/9/2011 (KC461214), JSD0812 (GQ849007). Pigeon Paramyxovirus-1 has 94% nucleotide identities with Pigeon paramyxovirus-1 strain AV324/96 (GQ429292). A phylogenetic tree (Figure 1) was constructed based on the amino acid sequences of the P gene in all isolates and one and the corresponding region of the other NDV strains retrieved from Gene Bank (Table 2). All six NDV field isolates, isolated in 2010-2012 outbreaks, were classified into genotype VII and one Pigeon Paramyxovirus-1 was located in the genotype V.




Figure 1. Nucleic acid Phylogenetic relationships of phosphoprotein gene of Newcastle disease genome isolated from broilers and pigeon, Iran. The Phylogenetic tree was generated using Neibor Joining model with MEGA (version 5.1 beta). Numbers below branches indicate bootstrap value from 1000 replicates, bootstrap values. Horizontal distances are proportional to the minimum number of nucleic acid differences required to join nodes. The vertical lines are for spacing branches and labels. Analysis was based on complete open reading frames of all gene segments. The scale bar represents the distance unit between sequence pairs. The virus genome of broiler characterized in this report is indicated as Black Circle. Black triangle is indicated for pigeon isolate. The sequences obtained from Gene Bank.

Discussion

Newcastle disease (ND) has a destructive economic

effect on commercial farms in Iran; thus, it is necessary to identify the ecology and transmission dynamics in order to improve disease control measures. This study was conducted to characterize

Table 1. The characteristic of Iranian Newcastle disease virus isolates (2010-2012).

No	Strain Name	Accession Number (Fusion Protein)	Accession Number (Phospho protein)	Isolation from Species	Clinical sign	Year	Province/Iran	Sample
1	IR-HGT2010.1	JX131352	KF824516	Broiler	Neurological	2010	Gilan	Brain
2	IR-HGT2011.1	JX131355	KF824517	Broiler	Neurological	2011	Mazandaran	Brain
3	IR-HGT2012.1	JX131357	KF824513	Broiler	Digestive	2012	Alborz	Cecal tonsil
4	IR-HGT2012.2	JX131358	KF824514	Broiler	Neurological	2012	Isfahan	Brain
5	IR-HGT2012.4	JX131360	KF824515	Breeder	Neurological	2012	Mazandaran	Brain
6	IR-OS-HGT2012	JX131350.1	KF824518	ostrich	Neurological	2012	Tehran	Brain
7	IR-PHG-2012	No submission	KF824519	Pigeon	Neurological	2012	Tehran	Brain

Table2. List of Newcastle disease viruses' characteristics that included in the phylogenetic.

Accession Number	Strain Name	Class	Sub-Genotype
AY935499	I-2	II	I
AY935498	99-1435	II	I
EU289028	VG/GA	II	II
NC_002617	B1	II	II
JF950510	Lasota	II	II
FJ430160	JS/9/05/Go	II	III
EF201805	Mukteswar	II	III
AY741404	Herts/33	II	IV
EU293914	Italien	II	IV
AY562986	Anhinga/U.S.(FI)/44083/93	II	V
AY562987	Gamefowl/U.S.(CA)/211472/02	II	V
JX518532	Pigeon paramyxovirus	II	V
GQ429292	AV324/96	II	V
JX901109	PMV-1/Belgium/98-238/1998	II	V
AJ415883	Pigeon paramyxovirus-1	II	V
AY562985	Cockatoo/Indonesia/14698/90	II	VI
AJ880277	Pigeon paramyxovirus 1	II	VI
FJ872531	Muscovy duck/China(Fujian)/FP1/00	II	VII
DQ659677	NA-1(Goose)	II	VII
AF473851	SF02/Goose	II	VII
FJ754272	WF01/D	II	VII
JF340367	JSG02/00	II	VII
KC461214	chicken/US/9/2011	II	VII
JSD0812	CGJ849007	II	VII
DQ097393	FE-R49/99	I	--
FJ794269	NDV08-004	I	--

NDVs isolated from five chicken, one ostrich and one Pigeon paramyxovirus-1. Phylogenetic analysis of sequences revealed that chicken and ostrich origin NDV isolates were closely related and placed in the genotype VII and Pigeon Paramyxovirus-1 was located in the genotype V. It is similar to previous results of researches of Iranians on F (Kianizadeh et al., 2002) and M gene (Langeroudi et al., 2012). Ebrahimi et al. (2012), based on F gene of Iranian NDV isolates, reported that they were representing

sub-genotype VIIb (Ebrahimi et al., 2012). According to the results of our study, it is the first report that mentioned Iranian NDV strains were located in genotype VIIb. Aboshah (2012), working on NDV isolates from Iranian commercial farms, been revealed that these isolates were located in VIIb subgenotype (Aboshah, 2012). Esmaealzadeh and co-investigators showed that the 6 Iranian isolates examined share significant similarity with 2 Russian isolates, Stern/18str/2755/2001 and VOL95 (Esmaeilzad et al., 2012). The results also indicate that genotype VII was the most prevalent isolate in Iran during the recent years. Genotype VII of NDV in Asia has been traced back to 1984 in Taiwan and to 1985 in Japan (Yang et al., 1999). According to Munir et al. (2012) the P gene was the most variable gene among the six NDV genes when compared with representatives of each genotype. Therefore, it is believed that the P gene is an evolutionary strategy of the virus that increases the coding capacity of the genome (Munir et al., 2012). It is the first report for characterization and phylogenetic analysis of Pigeon Paramyxovirus-1 in Iran. The data revealed that Iranian pigeon paramyxovirus-1 belongs to Genotype V beside other related pigeon Paramyxovirus-1. The presence of multiple NDV strains in Iran, the Far East and highly transmissible nature of the virus can complicate and increase the cost of attempts to prevent the spread of infection to the other parts of the world. In accordance with high occurrence of ND in pigeon population in Iran, more detailed studies should be carried out on their isolates. Finally, we suggest that researchers perform full length characterization on Iranian NDV isolates in recent years to reach more details of molecular epidemiology. In conclusion, our results reveal that NDV surveillance would be helpful to expand the understanding of NDV epidemiology in endemic regions.

Acknowledgments

The authors would like to acknowledge Mr. Ahmad Vahedi for his technical assistance.

References

1. Abdoshah, M., Pourbakhsh, S., Peighambari, S., Shojadoost, B., Momayez, R., Mojahedi, Z. (2012) Pathogenicity indices of Newcastle disease viruses isolated from Iranian poultry flocks in Iran. *J Vet Res.* 67: 159-164.
2. Alexander, D.J., Manvell, R.J., Lowings, J.P., Frost, K.M., Collins, M.S., Russell, P.H., Smith, J.E. (1997) Antigenic diversity and similarities detected in avian paramyxovirus type 1 (Newcastle disease virus) isolates using monoclonal antibodies. *Avian pathol.* 26: 399-418.
3. Bozorgmehri-Fard, M.H., Keyvanfar, H. (1979) Isolation of Newcastle disease virus from teals (*Anas crecca*) in Iran. *J Wildlife Dis.* 15: 335-337.
4. Ebrahimi, M.M., Shahsavandi, S., Moazenijula, G., Shamsara, M. (2012) Phylogeny and evolution of Newcastle disease virus genotypes isolated in Asia during 2008-2011. *Virus genes.* 45: 63-68.
5. Esmaelizad, M., Ashtiani, M.P., Jelokhani-Niaraki, S., Hashemnejad, K. (2012) Identification of 13 specific nucleotide patterns in the HN gene of Newcastle disease viruses isolated in Iran. *Turk J Biol.* 36: 135-142.
6. Jahanshiri, F., Eshaghi, M., Yusoff, K. (2005) Identification of phosphoprotein-phosphoprotein and phosphoprotein-nucleocapsid protein interaction domains of the Newcastle disease virus. *Arch Virol.* 150: 611-618.
7. Kianizadeh, M., Aini, M., Omar, A.R., Yusoff, K., Sahrabadi, M., Kargar, R. (2002) Sequence and phylogenetic analysis of the fusion protein cleavage site of Newcastle disease virus field isolates from Iran. *Acta Virologica.* 46: 247-251.
8. Kianizadeh, M., Ideris, A., Shahrabadi, M.S., Kargar, R., Pourbakhsh, S.A., Omar, S., Yusoff, A. (1999) Biological and Molecular characterization of Newcastle disease virus Isolated from Iran. *Arch Razi Ins.* 50: 1-10.
9. Langeroudi, A.G., Hosseini, H., Karimi, V., Hashemzadeh, M., Estabragh, A.S., Madadgar, O., (2012) Phylogenetic study base on matrix gene of Iranian Newcastle disease virus isolates, 2011-2012. *Comp Clin Pathol.* p.1-5.
10. Locke, D.P., Sellers, H.S., Crawford, J.M., Schultz-Cherry, S., King, D.J., Meinersmann, R.J., Seal, B.S., (2000) Newcastle disease virus phosphoprotein gene analysis and transcriptional editing in avian cells. *Virus Res.* 69: 55-68.
11. Miller, P.J., Decanini, E.L., Afonso, C.L. (2010) Newcastle disease: evolution of genotypes and the related diagnostic challenges. *Infection, genetics and evolution.* *J Mol Epidemiol Evol Genet Infect Dis.* 10: 26-35.
12. Miller, P.J., Kim, L.M., Ip, H.S., Afonso, C.L. (2009) Evolutionary dynamics of Newcastle disease virus. *Virology.* 391: 64-72.
13. Munir, M., Tonari, S., Abbas, M., Berg, M. (2012) Sequencing and analysis of the complete genome of Newcastle disease virus isolated from a commercial poultry farm in 2010. *Arch Virol.* 157: 765-768.
14. OIE, A. (2012) Manual of Diagnostic Tests and Vaccines for Terrestrial Animals. Office international des epizooties, Paris, France.
15. Rezaelanzadeh, G., Dadras, H., Ali, A.S.M., Nazemshirazi, M. (2011) Serological and molecular study of Newcastle disease virus circulating in village chickens of Fars province, Iran. *J Vet Med Anim Health.* 3: 105-111.
16. Smith, G.W., Hightower, L.E. (1981) Identification of the P proteins and other disulfide-linked and phosphorylated proteins of Newcastle disease virus. *J Virol.* 37: 256-267.
17. Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M., Kumar, S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol Biol Evol.* 28: 2731-2739.
18. Yang, C.Y., Shieh, H.K., Lin, Y.L., Chang, P.C. (1999) Newcastle disease virus isolated from recent outbreaks in Taiwan phylogenetically related to viruses (genotype VII) from recent outbreaks in western Europe. *Avian Dis.* 43: 125-130.

مطالعه شجره‌شناسی بر اساس ژن فسفو پروتئین ویروس‌های نیوکاسل جدا شده در ایران، ۱۳۸۹-۱۳۹۱

آرش قلیان چی لنگرودی^{۱*} حسین حسینی^۲ وحید کریمی^۳ امید مددگار^۱ مسعود هاشم زاده^۴ سید علی غفوری^۵
سید سینا باقری قادیکلایی^۶ سید میلاد واحدی

(۱) گروه میکروبیولوژی و ایمنی شناسی، دانشکده دامپزشکی دانشگاه تهران، تهران، ایران

(۲) گروه علوم درمانگاهی، دانشگاه آزاد اسلامی واحد کرج، کرج، ایران

(۳) گروه بیماری‌های طیور، دانشکده دامپزشکی دانشگاه تهران، تهران، ایران

(۴) موسسه واکسن و سرم‌سازی رازی، کرج، ایران

(۵) سازمان دامپزشکی کشور، تهران، ایران

(۶) انجمن علمی دانشجویی میکروبیولوژی، دانشکده دامپزشکی دانشگاه تهران، تهران، ایران

(دریافت مقاله: ۱۲ اسفند ماه ۱۳۹۲، پذیرش نهایی: ۲۲ اردیبهشت ماه ۱۳۹۳)

چکیده

زمینه مطالعه: ویروس نیوکاسل (NDV) عامل بیماری نیوکاسل (بیماری بسیار واگیر در پرندگان) سبب وارد آمدن خسارات اقتصادی قابل توجه به صنعت طیور سراسر جهان می‌گردد. بیماری نیوکاسل در ایران اندمیک است و از رخداد بیماری در طیور صنعتی و سایر پرندگان در کشور گزارش‌های متعددی داده شده است. **هدف:** مطالعه حاضر برای توصیف خصوصیات ویروس بیماری نیوکاسل (NDV) بر اساس ژن فسفو پروتئین در واگیری اخیر این بیماری در سال‌های ۸۹ تا ۹۱ انجام شد. **روش کار:** قطعه ژن فسفو پروتئین ویروس بیماری نیوکاسل از روی جدایه‌های بدست آمده از پنج قطعه مرغ، یک شتر مرغ و یک پارامیکسو ویروس تیپ اکبوتر باروش نسخه برداری معکوس - واکنش زنجیره‌های پلی‌مرز (RT-PCR) بدست آمد و تسلسل‌یابی شد. **نتایج:** تحلیل شجره‌شناسی نشان داد جدایه‌های حاصل از ماکیان و شتر مرغ بسیار به یکدیگر نزدیک بودند و در ژنوتیپ هفت و هشتاد و یک حاصل از پارامیکسو ویروس تیپ اکبوتر در ژنوتیپ پنج قرار دارند. **نتیجه‌گیری نهایی:** مطالعه حاضر اولین گزارش از توالی‌یابی ژن فسفو پروتئین حاصل از جدایه‌های ویروس بیماری نیوکاسل در ایران است. این مطالعه در درک همه‌گیرشناسی و خصوصیات ملکولی ویروس بیماری نیوکاسل بسیار کمک‌کننده خواهد بود.

واژه‌های کلیدی: بیماری نیوکاسل، فسفو پروتئین، مطالعه شجره‌شناسی

* نویسنده مسؤول: تلفن: +۹۸(۲۱)۶۱۱۱۷۰۵۳، نمابر: +۹۸(۲۱)۶۶۹۳۳۲۲۲، Email: ghanjyana@ut.ac.ir

Archiving
www.SID.ir

Surf and download all data from SID.ir: www.SID.ir

Translate via STRS.ir: www.STRS.ir

Follow our scientific posts via our Blog: www.sid.ir/blog

Use our educational service (Courses, Workshops, Videos and etc.) via Workshop: www.sid.ir/workshop