

**Original article****Inhibitory and killing activities of black tea (*Camellia sinensis*) extract against *Salmonella enterica* serovar Typhi and *Vibrio cholerae* O1 biotype El Tor serotype Ogawa isolates**

Shyamapada Mandal, PhD<sup>1\*</sup>, Manisha DebMandal, PhD<sup>2</sup>, Nishith Kumar Pal, MD<sup>1</sup>, Krishnendu Saha, MSc<sup>1</sup>

<sup>1</sup>Department of Microbiology, Bacteriology and Serology Unit, Calcutta School of Tropical Medicine, C. R. Avenue, Kolkata-700 073, India

<sup>2</sup>Department of Physiology and Biophysics, KPC Medical College and Hospital, 1F Raja S C Mallick Road, Jadavpur, Kolkata-700 032, India

**How to cite this article:**

Mandal S, DebMandal M, Pal NK, Saha K. Inhibitory and killing activities of black tea (*Camellia sinensis*) extract against *Salmonella enterica* serovar Typhi and *Vibrio cholerae* O1 biotype El Tor serotype Ogawa isolates. Jundishapur J Microbiol. 2011; 4(2): 115-121.

**Received:** August 2010

**Accepted:** October 2010

**Abstract**

**Introduction and objective:** Previous studies have shown the potential antibacterial activity of black tea, *Camellia sinensis*, extract against many pathogenic bacteria including Gram-positives (*Listeria monocytogenes* and *Staphylococcus aureus*) and Gram-negatives (*Vibrio cholerae* and *Salmonella enterica* serovar Typhi). The aim of the current study was to assess the *in vitro* antibacterial activity of *C. sinensis* ethanolic extract (CAS) against multidrug-resistant (MDR) clinical isolates of *S. Typhi* and *V. cholerae* O1 biotype El Tor serotype Ogawa (*V. cholerae* Ogawa).

**Materials and methods:** The zone diameter of inhibition (ZDI) and minimum inhibitory concentration (MIC) values of CAS for *S. Typhi* and *V. cholerae* Ogawa isolates were determined by agar diffusion and agar dilution techniques. Bacterial killing studies were carried out in order to assess the bactericidal activity of CAS against *S. Typhi* and *V. cholerae* Ogawa, from Kolkata, India.

**Results:** For the *S. Typhi* and *V. cholerae* Ogawa isolates, the ZDIs of CAS ranged 12-17mm and 13-21mm, respectively, and the MICs of the extract were 400-600µg/ml and 200-600µg/ml, respectively. The CAS had bactericidal action at concentrations 512µg/ml and 256µg/ml, respectively for *S. Typhi* and *V. cholerae* Ogawa.

**Conclusion:** The black tea (*C. sinensis*) extract could be useful in combating emerging drug-resistance among enteropathogens including *S. Typhi* and *V. cholerae* Ogawa.

**Keywords:** Black tea; Bactericidal activity; Zone diameter of inhibition; Minimum inhibitory concentration; *Salmonella* serovar Typhi; *Vibrio cholerae* Ogawa

**\*Address for correspondence:**

Dr. Shyamapada Mandal, Department of Zoology, Gurudas College, Narkeldanga, Kolkata-700 054, India, Tel: +9133 65211163; E-mail: samtropmed@gmail.com

Jundishapur Journal of Microbiology, School of Medicine, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, Tel: +98611 3330074; Fax: +98611 3332036; URL: <http://jjm.ajums.ac.ir>; E-mail: editorial office: [jjm@ajums.ac.ir](mailto:jjm@ajums.ac.ir)

JJM. (2011); 4(2): 115-21.

## Introduction

Typhoid and cholera, which are caused due to the infection of *Salmonella enterica* serovar Typhi (*S. Typhi*) and *Vibrio cholerae*, respectively, are endemic in many developing countries like India, and the rapid emergence of resistance to multiple antibiotics among these two strains makes the situation alarming [1-4]. This phenomenon of multi-drug resistance of human pathogenic microorganisms has necessitated the search for new antimicrobial substances from other sources including plants, which have long been utilized as the source of therapeutic agents worldwide, and to treat many life threatening diseases due to bacterial infections [5,6].

The antimicrobial activities of medicinal plants have been reported from different countries against different pathogens [7]. Black tea (*Camellia sinensis*) has been shown to have a wide range of beneficial physiological and pharmacological effects. Toda *et al.* [8] found that extracts of tea inhibited and killed a large number of Gram-positive and Gram-negative bacteria including *S. Typhi* and *V. cholerae*. The inhibitory as well as bactericidal activity of tea extracts against various pathogenic bacteria causing several infections has been reported [8-11].

Tiwari *et al.* [12] reported antibacterial activity of *C. sinensis* against different bacterial genera including *S. Typhi*. But, from our region, no scientific study has been documented on antibacterial activity of *C. sinensis* against any bacteria including *S. Typhi* and *V. cholerae* causing, respectively, typhoid and cholera. Thus, it seems reasonable to explore the possibility of using *C. sinensis* extract for eradication of pathogenic bacteria. Therefore, in this study, we investigated the antibacterial activity of the ethanolic extract of black tea (*C. sinensis*) against clinical isolates of *S.*

*Typhi* and *V. cholerae* O1 biotype El Tor serotype Ogawa (*V. cholerae* Ogawa) in order to explore the potential inhibitory and killing activities of *C. sinensis* against the typhoidal and choleraogenic bacteria, which are life-threatening diseases in Kolkata, India.

## Materials and methods

### Bacterial isolates

The multidrug-resistant bacterial isolates of *S. Typhi* (n=12) and *V. cholerae* Ogawa (n=9), were obtained, from blood samples of typhoid patients and rectal swabs of cholera patients, respectively at the Calcutta School of Tropical Medicine, India. The control strain was *Escherichia coli* ATCC 25922. For positive control, the all *S. Typhi* and *V. cholerae* Ogawa isolates were treated against chloramphenicol, by agar dilution, and found susceptible to the antibiotic.

### Plant materials and extract preparation

The dried leaves of Indian black tea (*C. sinensis*), in the form of granules, was purchased from market in Kolkata, India, and the sample was stored in the plastic bag at 4°C before extract preparation. The crude extract of black tea was prepared following the method described earlier [13], with slight modification. Briefly, 50g of the sample was soaked in 50ml of 95% ethanol for 48h with vigorous shaking at 2h interval. The mixture was then filtered, and the filtrate was vaporized to dryness, and weighed. The stock solution of the crude ethanolic extract of *C. sinensis* (CAS) was prepared by dissolving the dried extract with 50% ethanol in sterile double distilled water to obtain the final concentration of 10mg/ml.

### Media and bacterial inocula

In the present study, two media (Hi-Media, India) were used. Mueller-Hinton broth

(MHB) for subculturing the strains, inocula preparation and performing bacterial killing experiments, and Mueller-Hinton agar (MHA) to determine the zone diameter of inhibition (ZDI) and minimum inhibitory concentration (MIC). The bacterial inocula were prepared as  $10^4$ cfu/spot,  $10^8$ cfu/ml and  $5 \times 10^5$ cfu/ml, respectively for agar dilution, agar diffusion and kill-kinetic studies, following the previously described methodology [13].

#### *Antibacterial activity of tea extract*

The ZDIs of the isolates were determined with agar diffusion technique, as described by Nas [14], using an inoculum of  $10^8$ cfu spread on MHA plates. Each of the inoculated plates was divided into two equal sectors, and the extract, 25 $\mu$ l (equivalent to 250 $\mu$ g), was dropped on to a sector marked previously as CAS; on to the other sector (which was used as the control) 25 $\mu$ l of 50% ethanol was placed. After allowing the diffusion of the extract in to the agar for about 45mins at room temperature, the plates were incubated at 35°C for 24h, and ZDIs produced due to the action of the extract were measured for all the isolates.

The control sectors had no ZDI to ZDI of 6mm for *S. Typhi* and *V. cholerae* Ogawa isolates, and thus, the sensitivity of CAS for the isolates were considered with  $ZDI \geq 7$ mm [15]. The agar dilution technique was used to determine the MICs of CAS for all the bacterial isolates, following spot inoculation, and the extract concentrations utilized in the experiment ranged 100-600  $\mu$ g/ml; the other details are described elsewhere [13]. Briefly, each of the tea extract mixed agar plates was divided into 22 equal sectors, and inoculated with approximately  $10^4$ cfu per sector; thus for 22 isolates, six agar plates were prepared for six different concentrations of the extract (100-600 $\mu$ g/ml). The plates were then incubated at 35°C for 24h. The MICs were

defined as the lowest concentration of the extract at which no visible growth was found; hazy growth and one or two colonies on the spot were ignored.

Bacterial killing studies, in duplicate, were carried out for both *S. Typhi* 2K/2 strain (CAS MIC 600 $\mu$ g/ml) and *V. cholerae* Ogawa S11 strain (CAS MIC 600 $\mu$ g/ml), in MHB using approximately  $5 \times 10^5$ cfu/ml of initial inoculum, following the protocol published earlier [13]. In this method, in order to determine the effect of different concentrations (16-512 $\mu$ g/ml) of the extract on bacterial density (cfu/ml), the bacterial suspension ( $5 \times 10^5$ cfu/ml) in MHB was incubated for 24h at 35°C, and viable cells were counted for each of the extract concentrations. The  $\geq 3$  log<sub>10</sub> decrease in the inoculum after 24h of incubation was considered as the bactericidal activity of the extract [16].

#### *Statistical analysis*


The  $\chi^2$ -test was performed to compare the killing activities of CAS against *S. enterica* serovar Typhi and that of CAS against *V. cholerae* O1 biotype El Tor serotype Ogawa based on their growths (in the terms of cfu/ml) at different concentrations. P value of  $\leq 0.05$  was considered significant. The differences in the inhibitory activities of the extract against the two bacterial strains considered significant at  $p \leq 0.05$  following t-test.

#### **Results**


Figure 1 represents the ZDI obtained due to the action of CAS against the test isolates. The *S. Typhi* isolates had ZDIs of 12-17mm, and the ZDI ranged 13-21mm for *V. cholerae* Ogawa isolates, and thus all the bacterial isolates were sensitive to CAS.

The MICs of CAS for the test bacteria are shown in figure 2. All the *S. Typhi* isolates (12; 100%) showed CAS MICs 400-600 $\mu$ g/ml, and most of them (7;

58.33%) had MIC of 500µg/ml. Among nine *V. cholerae* Ogawa isolates, 6(66.67%) had CAS MICs of 400-600µg/ml, and the rest 3 (33.33%) isolates had 200-300µg/ml.



**Fig. 1:** Zone diameter of inhibition (ZDI) due to the action of black tea (*C. sinensis*) extracts against *S. enterica* serovar Typhi and *V. cholerae* O1 biotype El Tor serotype Ogawa isolates




**Fig. 2:** Minimum inhibitory concentration (MIC) values of black tea (*C. sinensis*) extracts for *S. enterica* serovar Typhi and *V. cholerae* O1 biotype El Tor serotype Ogawa isolates

The ZDI and MIC of the *E. coli* ATCC 25922 control strain was 16mm and 300µg/ml, respectively (not shown in the figures).

Figure 3 illustrates the killing activities of various concentrations (16-512µg/ml) of CAS against *S. Typhi* and *V. cholerae* Ogawa. At 16µg/ml (lowest concentration used in the experiment) of CAS the bacterial density increased up to 6.34 and 6.41 log<sub>10</sub> cfu/ml, respectively for *S. Typhi*

and *V. cholerae* Ogawa strains. The CAS, at 32µg/ml, started to exhibit killing effect on *S. Typhi*, while the extract had similar effect on *V. cholerae* Ogawa at 64µg/ml. The CAS showed bactericidal activity at concentrations of 512µg/ml and 256µg/ml, respectively, against *S. Typhi* and *V. cholerae* Ogawa, leaving 2.19 log<sub>10</sub> cfu/ml and 2.58 log<sub>10</sub> cfu/ml, respectively, after 24h.



**Fig. 3:** Killing activity of black tea (*C. sinensis*) extracts on *S. enterica* serovar Typhi and *V. cholerae* O1 biotype El Tor serotype Ogawa

The difference between the inhibitory activities of CAS, both in the terms of ZDI and MIC, for *S. Typhi* and *V. cholerae* was significant ( $p < 0.05$ ) as determined with t-test. The significant differences ( $p < 0.001$ ) were observed between the number of viable cells of the two bacterial strains, at their bactericidal concentrations of 512µg/ml (for *S. Typhi*) and 256µg/ml (for *V. cholerae* Ogawa), using  $\chi^2$  test.

### Discussion

The development of resistance to multiple antibiotics among disease causing bacteria [17,18] prompted many researchers [19,20] to find the new sources of non-antibiotic drugs mainly among the plant extracts in order to overcome the disadvantages of antibiotic resistance. The present study was conducted to evaluate the *in vitro* antibacterial activity of leaf granules extracts of black tea, *C. sinensis*, against *S.*

Typhi and *V. cholerae* Ogawa. Michalczyk and Zawislak [21] reported that the black tea extract had strongest inhibition activity than any other (pu-erh and green) tea against human intestinal bacteria, and the *S. enteritidis* was inhibited to a larger extent by black tea than by green tea. Shetty *et al.* [22] showed that the black tea, Japanese green tea and China tea inhibited the growth of various bacteria, causing diarrhoeal diseases, including *V. cholerae* and *S. Typhi*.

Mbata *et al.* [23] reported earlier the antibacterial activity of *C. sinensis* leaf against *Listeria monocytogenes* by disc diffusion method, and showed that the methanolic extract had greater antibacterial property (ZDI 20.1mm) compared to the water extract (ZDI 10 mm); by agar diffusion method the water extract did not show inhibitory effect, while the methanolic extract had ZDI of 15mm. The CAS, in the present study, exhibited potent antibacterial activities against all bacterial isolates tested, and the extract had stronger activity against *V. cholerae* Ogawa compared to *S. Typhi* showing ZDIs of 13-21mm (Mean of  $17.11 \pm 2.934$  SD) and 12-17mm (Mean of  $14.67 \pm 2.015$  SD), respectively, and with respect to the bacteria used, the difference between inhibitory activity of the CAS was significant ( $p < 0.05$ ).

The degree of antibacterial activity of the extract in the terms of MIC values against the bacterial strains also supports the findings of the agar diffusion method ( $p < 0.05$ ), as represented in this communication. The isolates of *V. cholerae* Ogawa had CAS MICs of 200-600 $\mu$ g/ml (Mean of  $411.11 \pm 126.93$  SD), and the MICs of *S. Typhi* isolates were between 400  $\mu$ g/ml and 600 $\mu$ g/ml (Mean of  $508.33 \pm 66.855$  SD). Tiwari *et al.* [12] showed wide differences in the MIC (9.089-94.61mg/ml) of tea extract against different bacterial strains including *S. Typhi* with

MICs of 79.56-91.98mg/ml. Mbata *et al.* [23] recorded MICs of 0.26 and 0.68mg/ml, respectively of methanolic and aqueous extracts of *C. sinensis* leaf against *L. monocytogenes*. The differences in antibacterial activities of *C. sinensis* extract might be related to the degrees of susceptibility of cell wall of test bacterial strains, kinds of solvent used in the extract preparation, and different methodology adopted in the determination of antibacterial activity [24].

Mandal *et al.* [13] reported earlier the findings of time-kill studies with *Azadirachta indica* seed extract against *S. enterica* serovar Typhi. Shetty *et al.* [22] showed the bactericidal activity of *C. sinensis* against *V. cholerae* and *S. Typhi*. The CAS, in the present study, showed bactericidal activity against *S. Typhi* and *V. cholerae* Ogawa, at 512 $\mu$ g/ml and 256 $\mu$ g/ml, respectively. There were significant decreases in the number of viable *S. Typhi* and *V. cholerae* Ogawa cells at the bactericidal concentrations, comparing to the initial inoculum used ( $p < 0.001$ ).

The CAS, both at 256 and 512 $\mu$ g/ml was found more effective against *V. cholerae* Ogawa than against *S. Typhi*, after 24h of incubation, and significant differences were found between the number of viable cells of the two bacterial strains at each concentration ( $p < 0.001$ ). These values may be used as guide for treatment of bacterial infections, and thus may form the basis in the determination of optimal dosage selection. Such bactericidal action of the CAS might be due to the presence of several active components singly or in combination against *V. cholerae* Ogawa and *S. Typhi* strains.

The polyphenolic compounds including several catechins, mainly epigallocatechin gallate, and the theaflavins are reported to be the microbiologically active molecules in

black tea extracts [25-27], and the combination of flavor compounds, especially indole with some of the sesquiterpenes, are known to display marked bactericidal synergy [28].

In the present study, these compounds could be responsible for the inhibition of *S. Typhi* and *V. cholerae* Ogawa. The antibacterial activities displayed by the CAS following different methods, as represented in the present communication, suggest the ethnopharmacological use of black tea (*C. sinensis*) extract as a remedy to treat *S. enterica* serovar Typhi and *V. cholerae* O1 biotype El Tor serotype Ogawa infections, which in some situation result in outbreaks of the diseases (typhoid and cholera, respectively) caused by them [2,29].

### Conclusion

It is concluded that black tea (*C. sinensis*), which is one of the most popular beverages worldwide, has medicinal and health-promoting properties. Thus, the antimicrobial agents of it could serve as viable supplements of the present range of antibiotics. Thus, it is implied that *C. sinensis* extract will be useful as an effective antibacterial agent against both *S. enterica* serovar Typhi and *V. cholerae* O1 biotype El Tor serotype Ogawa. However, considerable research is required to improve the bioavailability of the active principles and to determine dose as well as toxicity before those can be used as therapeutic agents.

### References

- 1) Capoor MR, Nair D, Aggarwal P, Mathys V, Dehem M, Bifani PJ. *Salmonella enterica* Serovar Typhi: Molecular analysis of strains with decreased susceptibility and resistant to ciprofloxacin in India from 2001-2003. *Braz J Infect Dis.* 2007; 1: 423-5.
- 2) Mandal S, Mandal MD, Pal NK. Antimicrobial resistance pattern of *Salmonella typhi* isolates in Kolkata, India during 1991-2001: a retrospective study. *Jpn J Infect Dis.* 2002; 55: 58-9.
- 3) Sur D, Dutta S, Sarkar BL, Manna B. Occurrence, significance and molecular epidemiology of cholera outbreaks in West Bengal. *Indian J Med Res.* 2007; 125: 772-6.
- 4) Taneja N, Kaur J, Sharma K, et al. A recent outbreak of cholera due to *Vibrio cholerae* O1 Ogawa in and around Chandigarh, North India. *Indian J Med Res.* 2003; 117: 243-6.
- 5) Taran M, Ghasempour HR, Shirinpour E. Antimicrobial activity of essential oils of *Ferulago angulata* subsp. *carduchorum*. *Jundishapur J Microbiol.* 2010; 3: 10-4.
- 6) Amin M, Segatoleslami S, Hashemzadeh M. Antimycobacterial activity of partial purified extract of *Allium ascalonicum*. *Jundishapur J Microbiol.* 2009; 2: 144-7.
- 7) Rios JL, Recio MC. Medicinal plants and antimicrobial activity. *J Ethnopharmacol.* 2005; 100: 80-4.
- 8) Toda M, Okubo S, Hiyoshi R, Shimamura T. The bactericidal activity of tea and coffee. *Lett Appl Microbiol.* 1989; 8: 123-5.
- 9) Ciraj AM, Sulaim J, Mamatha B, Gopalkrishna, BK, Shivananda PG. Antibacterial activity of black tea (*Camelia sinensis*) extract against *Salmonella* serotypes causing enteric fever. *Indian J Med Sci.* 2001; 7: 376-81.
- 10) Diker KS, Akan M, Hascelik G, Yurdakok M. The bactericidal activity of tea against *Campylobacter jejuni* and *Campylobacter coli*. *Lett Appl Microbiol.* 1991; 12: 34-5.
- 11) Hara Y, Ishigami T. Antibacterial activities of tea polyphenols against foodborne pathogenic bacteria. *J Jpn Soc Food Sci Technol.* 1989; 36: 996-9.
- 12) Tiwari RP, Bharti SK, Kaur HD, Kaur RP, Dikshit, Hoondal GS. Synergistic antimicrobial activity of tea and antibiotics. *Indian J Med Res.* 2005; 122: 80-4.
- 13) Mandal S, Mandal M, Pal NK. Antibacterial potential of *Azadirachta indica* seed and *Bacopa monniera* leaf extracts against multidrug resistant *Salmonella enterica*

- serovar Typhi isolates. *Arch Med Sci.* 2007; 3: 14-8.
- 14) Nas MN. *In vitro* studies on some natural beverages as botanical pesticides against *Erwinia amylovora* and *Curtobacterium flaccumfaciensis* subsp. *Poinsettiae*. *Turk J Agric For.* 2004; 28: 57-61.
  - 15) Nascimento GGF, Locatelli J, Freitas PC, Silva GL. Antibacterial activity of plant extracts and phytochemicals on antibiotic-resistant bacteria. *Braz J Microbiol.* 2000; 31: 247-56.
  - 16) Leclercq R, Bingen E, Su QH, Lambert-Zechovski N, Courvalin P, Duval J. Effects of combinations of  $\beta$ -lactams, daptomycin, gentamycin and glycopeptides against glycopeptide resistant Enterococci. *Antimicrob Agents Chemother.* 1991; 35: 92-8.
  - 17) Davies J. Inactivation of antibiotics and the dissemination of resistance genes. *Science.* 1994; 264: 375-82.
  - 18) Marchese A, Schito GC. Resistance patterns of lower respiratory tract pathogens in Europe. *Int J Antimicrob Agents.* 2001; 16: 25-9.
  - 19) Mansouri S, Foroumadi A, Ghanei T, Gholamhosseinian NA. Antibacterial activity of the crude extracts and fractionated constituents of *Myrtus communis*. *Pharmaceut Biol.* 2001; 39: 399-401.
  - 20) Srinivasan D, Nathan S, Suresh T, Perumalsamy O. Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine. *J Ethnopharmacol.* 2001; 74: 217-20.
  - 21) Michalczyk M, Zawislak A. The effect of tea infusions on the proliferation of selected bacteria important for the human intestinal tract. *Acta Sci Pol Technol Aliment.* 2008; 7: 59-65.
  - 22) Shetty M, Subbannayya K, Shivananda PG. Antibacterial activity of tea (*Camellia sinensis*) and coffee (*Coffea arabica*) with special reference to *Salmonella typhimurium*. *J Commun Dis.* 1994; 26: 147-50.
  - 23) Mbata TI, Debiao LU, Saikia A. Antibacterial activity of the crude extract of Chinese green tea (*Camellia sinensis*) on *Listeria monocytogenes*. *African J Biotechnol.* 2008; 7: 1571-3.
  - 24) Turkmen N, Velioglu YS, Sari F, Polat, G. Effect of extraction conditions on measured total polyphenol contents and antioxidant and antibacterial activities of black tea. *Molecules.* 2007; 12: 484-96.
  - 25) Hamilton-Miller JMT. Antimicrobial properties of tea (*Camellia sinensis* L.). *Antimicrob Agents Chemother.* 1995; 39: 2375-7.
  - 26) Kawamura J, Takeo T. Antibacterial activity of tea catechin to *Streptococcus mutans*. *J Jpn Soc Food Sci Technol.* 1989; 36: 463-7.
  - 27) Yam TS, Shah S, Hamilton-Miller JMT. Microbiological activity of whole and fractionated crude extracts of tea (*Camellia sinensis*), and of tea components. *FEMS Microbiol Lett.* 1997; 152: 169-74.
  - 28) Muroi H, Kubo I. Combination effects of antibacterial compounds in green tea flavor against *Streptococcus mutans*. *J Agric Food Chem.* 1993; 41: 1102-5.
  - 29) Mandal S, Mandal MD, Pal NK. Plasmid mediated antibiotic resistance of *Vibrio cholerae* O1 biotype El Tor serotype Ogawa associated with an outbreak in Kolkata, India. *Asian Pacific J Trop Med.* 2010; 3: 637-41.