

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



عضویت در خبرنامه



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



مباحث پیشرفته یادگیری عمیق؛
شبکه های توجه گرافی
(Graph Attention Networks)



کارگاه آنلاین آموزش استفاده از
وب آو ساینس



کارگاه آنلاین مقاله روزمره انگلیسی



Antibiotic Susceptibility of Bacterial Strains Isolated From Urinary Tract Infections in Karaj, Iran

Rahem Khoshbakht^{1*}, Ayub Salimi², Hesamaddin Shirzad Aski¹, Hale Keshavarzi³

¹ Department of pathobiology, school of veterinary medicine, Shiraz University, Shiraz, IR Iran

² Razi Vaccine and Serum Research Institute, Karaj, IR Iran

³ School of veterinary medicine, Shiraz University, Shiraz, IR Iran

ARTICLE INFO

Article type:
Original Article

Article history:
Received: 13 Mar 2012
Revised: 07 May 2012
Accepted: 21 May 2012

Keywords:
Urinary Tract Infections
Antibiotic Susceptibility

ABSTRACT

Backgrounds: Urinary Tract Infection (UTI) is one of the most common bacterial diseases worldwide that can be presented asymptomatic or symptomatic, characterized by a wide range of symptoms from mild irritative voiding to bacteremia, sepsis, or even death. In latter decades, widespread utilization of antibiotics has resulted in increased incidence of antibiotic resistance among urinary tract pathogens all over the world.

Objectives: The aim of this study was to determine patterns of antibiotic resistance bacteria isolated from urinary tract infections in the city of Karaj, Iran.

Materials and Methods: This descriptive study was conducted in the city of Karaj from November 2009 to August 2010. Bacterial isolates were identified by standard biochemical tests. Antibacterial susceptibility test was performed by Kirby-Bauer method against some common antibiotics for Gram negative and Gram positive bacteria and the data were analyzed by SPSS software, version 12.0.1.

Results: In this study, 230 (8.06%) out of 2852 patients showed positive urine cultures of which there were 204 (88.69%) females and 26 (11.3%) males. 180 (83.17%) cases of isolated bacteria were Gram negative bacilli while 50 (21.73%) cases were Gram positive cocci. *E. coli* isolates, as the most frequent Gram negative bacteria, showed high resistance to cephalothin (88.16%) and high susceptibility to nitrofurantoin (90.78%). Also *Staphylococcus saprophyticus* isolates, as the most frequent Gram positive bacteria, exhibited high resistance to ampicillin, tetracycline, and erythromycin (92.31%) and high susceptibility to nitrofurantoin and vancomycin (92.3%).

Conclusions: considering the results of this study, *E. coli* isolates were the predominant pathogens in UTI. Our data demonstrated that antibiotics with the highest activity included ciprofloxacin and nitrofurantoin and those with the least activity included ampicillin and sulfamethoxazole-trimethoprim. Our findings reinforce the need for ongoing investigations to show the trends of antibiotic resistance which can help antibiotic prescription in the clinics.

Published by Kowsar Corp, 2013. cc 3.0.

► Implication for health policy/practice/research/medical education:

The article will help Clinicians in predicting and planning correct strategies for UTI therapy.

► Please cite this paper as:

Khoshbakht R, Salimi A, Shirzad Aski H, Keshavarzi H. Antibiotic Susceptibility of Bacterial Strains Isolated From Urinary Tract Infections in Karaj, Iran. *Jundishapur J Microbiol.* 2013;6(1):86-90. DOI: 10.5812/jjm.4830

* Corresponding author: Rahem Khoshbakht, Department of pathobiology, school of veterinary medicine, Shiraz University, Shiraz, IR Iran. Tel: +98-7116138697, Fax: +98-9113919059, E-mail: khoshbakht_r@shirazu.ac.ir

DOI: 10.5812/jjm.4830

© 2013 Ahvaz Jundishapur University of Medical Sciences; Published by Kowsar Corp.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

1. Background

Urinary tract infections (UTIs) are among the most common bacterial diseases worldwide (1), and the commonest genitourinary disease in children (2). UTI can be asymptomatic or symptomatic, characterized by a wide spectrum of symptoms ranging from mild irritative voiding to bacteremia, sepsis, or even death (3). UTIs account for approximately 11.3 million office visits with an overall cost of 1.6 billion dollars annually in the US health budget (4). Infection may occur at any part of the genitourinary tract, including urethra, bladder, ureter, renal pelvis, or renal parenchyma (5, 6). Most infections are caused by retrograde ascent of bacteria from fecal flora to bladder and kidney via urethra, especially in females whom the urethra is shorter and wider (7).

Anatomical structure of female urethra and vagina make them susceptible to trauma during sexual intercourse and bacterial passage through urethra up to the bladder during pregnancy and delivery (8, 9). The infection may involve only lower or both upper and lower parts of urinary tract. The term "cystitis" has been used to describe a syndrome including dysuria, frequency, and occasionally supra pubic tenderness. "Acute pyelonephritis" describes a clinical syndrome characterized by flank pain or tenderness, or both, and fever, often accompanied by dysuria, urgency, and frequency (10). Majority of UTIs are not life threatening and do not cause any irreversible damage. However, when the kidneys are involved, there is a risk of irreparable tissue damage with an increased risk of bacteremia (11).

Bacteria are the major causative organism and are responsible for more than 95% of UTI cases (12). *Escherichia coli* is the most frequent infecting organism in acute infections (13, 14). Area-specific monitoring studies aimed to understanding the types of pathogens responsible for UTIs and their resistance patterns may help clinicians to select proper empirical treatment (15).

2. Objectives

The aims of this study were to investigate microbial species isolated from patients with UTI and evaluate their in vitro susceptibility patterns to commonly used antimicrobial agents in Karaj, Iran.

3. Materials and Methods

3.1. Specimen Collection and Bacterial Isolates

This study was conducted from November 2009 to August 2010 on 2852 untreated patients with clinical symptoms of UTI referred to a central diagnostic lab in Karaj, Iran. Clean catch midstream urine samples (MSU) were collected in sterile disposable containers (4-5 ml) and transported immediately to the laboratory. Urine specimens were subjected to general urine

examinations using direct microscopy for white blood cell (WBC) counting. Urine samples were cultured on 5% blood agar, MacConkey, and Eosin-Methylene blue (EMB) agar (Oxoid Ltd, Basingstoke, Hampshire, UK) using calibrated loops for semi-quantitative method (16) and incubated in both aerobic and anaerobic conditions for 24 hours at 37°C. Cultures without any colony at the end of 24h incubation were further incubated for 48h. Samples with colony count equal or more than 10^5 Cfu/ml were considered positive. The isolates were identified and confirmed using standard microbiological methods including Gram staining, colonial morphology on media, growth on selective media, lactose and mannitol fermentation, H₂S production, catalase, oxidase, coagulase, indole, and citrate utilization, and urease test (16-18).

3.2. Antimicrobial Susceptibility Testing

Antimicrobial susceptibility testing was performed on Mueller-Hinton agar (Merck, Germany) using disk diffusion (Kirby Bauer's) technique according to Clinical and Laboratory Standards Institute (CLSI) guidelines. The antibiotic discs and their concentrations consisted of Cefotaxime (CTX, 30µg, Polfa, Tarchomin, Poland), Cephalothin (CF, 30µg, Polfa, Tarchomin, Poland), Imipenem (IPM, 10µg, Zeneca, Macclesfield, UK), Ciprofloxacin (CP, 5µg), Ceftazidime (CAZ, 30µg), Ceftizoxime (CT, 30µg), Gentamicin (GM, 10µg), Ampicillin (AM, 10µg), Trimethoprim-sulfamethoxazole (SXT, 30µg), Nitrofurantoin (FM, 50µg) and Nalidixic acid (NA, 30µg) for Gram negative isolates, and Chloramphenicol (C, 30µg, Bayer, Wuppertal, Germany), Ampicillin (AM, 10µg), Erythromycin (E, 15µg), Norfloxacin (NOR, 10µg), Gentamicin (GM, 10µg), Vancomycin (V, 30µg), Nitrofurantoin (FM, 50µg), Tetracycline (TE, 30µg), and Trimethoprim-sulfamethoxazole (SXT, 30µg) for Gram positive isolates all obtained from Hi media, India.

E. coli (ATCC 25922), *Staphylococcus aureus* (ATCC 25923), and *Pseudomonas aeruginosa* (ATCC 27853) were used as quality control strains.

Data were analyzed using SPSS version 12.0.1. Discrete variables were expressed as percentages and proportions were compared using the Chi-square test. Statistical significant difference was considered at value of $P \leq 0.05$.

4. Results

In this study, 230 (8.06%) out of 2852 patients showed positive urine cultures of which there were 204 (88.69%) females and 26 (11.3%) males. 180 (83.17%) cases of the isolated bacteria were Gram negative bacilli while 50 (21.73%) cases were Gram positive cocci. The frequency of isolated microorganisms and their relations to sex is given in Table 1.

Patients' age ranged from 10 days to 90 years with an average age of 26.17 years. The age distribution was

Table 1. : Frequency of Bacterial Agents Isolated From Urine Specimens and Their Relation to Sex in This Study

Isolated Bacteria	No. (%)	Female, No. (%)	Male, No. (%)
Gram-Negative bacilli			
<i>Escherichia coli</i>	152 (66.08)	136 (89.5)	16 (10.5)
<i>Klebsiella spp.</i>	22 (9.6)	18 (81.8)	4 (18.2)
<i>Proteus spp.</i>	4 (1.7)	4 (100)	0 (0)
<i>Enterobacter spp.</i>	2 (0.86)	2 (100)	0 (0)
Gram-Positive cocci			
<i>Staphylococcus saprophyticus</i>	26 (11.3)	22 (84.6)	4 (15.4)
<i>Staphylococcus aureus</i>	4 (1.7)	4 (100)	0 (0)
<i>Streptococcus group B</i>	10 (8.69)	10 (100)	0 (0)
<i>Enterococcus spp.</i>	10 (8.69)	8 (80)	2 (20)
Total	230 (100)	204 (88.7)	26 (11.3)

22.6% for patients aged ≤ 2 year, 9.56% for 2 to 14 years, 57.39% for 15 to 60 years, and 9.56% for ≥ 60 years. *E. coli* was the most common microorganism isolated in all age-groups and both sexes. No significant difference was found between isolated bacteria and age of the patients. The isolation percentages of bacterial species were found as *E. coli* (66.08%); *S. saprophyticus* (11.3%); *Klebsiella spp.* (9.6%); *S. group B* (4.34%); *Enterococcus spp.* (4.34%); *S. aureus* (1.7%); *Proteus spp.* (1.7%), and *Enterobacter spp.* (0.86%). *E. coli* isolates, the predominant cause of UTIs, showed high resistance to gentamicin, ampicillin, SXT, and cephalothin, and higher sensitivity to imipenem, ciprofloxacin, nitrofurantoin, and ceftizoxime.

Klebsiella strains displayed a similar resistance pattern as for *E. coli* and showed high susceptibility to ciprofloxacin, ceftizoxime, nalidixic acid, and imipenem and high resistance to gentamicin, cephalothin, and ampicillin. On *Proteus* strains, ampicillin, nitrofurantoin, and SXT showed the least (0%, 0%, and 25%,

respectively) and ciprofloxacin, ceftizoxime, and imipenem showed the highest susceptibility (100%, 100% and, 75%, respectively) that were almost similar to the results with other Gram positive bacteria. The results were slightly different in *Enterobacter* as it can be seen the utmost resistance of *Enterobacter* was recorded to ampicillin and nalidixic acid while 100% susceptibility to other antibiotics was recorded.

In this study, staphylococci were responsible for about 13% of UTIs cases; among these, *S. saprophyticus* isolates as most frequent Gram positive bacteria showed high resistance to ampicillin, tetracycline, and erythromycin, and high sensitivity to nitrofurantoin, gentamicin, norfloxacin, and vancomycin. The rates of resistance to 11 selected antimicrobial agents against Gram negative bacilli and to 9 selected antibiotics against Gram positive cocci are summarized in Table 2 and Table 3, respectively.

Table 2. Prevalence and Antimicrobial Resistance of Gram Negative Uropathogens Isolated in this Study

	No.	AM	CP	CT	IPM	GM	SXT	NA	CTX	FM	CAZ	CF
<i>E. coli</i>	152	73.69	26.32	15.79	15.79	73.69	69.74	60.53	38.16	9.22	55.27	88.16
<i>Klebsiella spp.</i>	22	63.64	27.28	27.28	18.9	63.64	63.64	27.28	45.46	54.55	54.55	45.46
<i>Proteus spp.</i>	4	100	0	0	25	50	75	50	50	100	50	25
<i>Enterobacter spp.</i>	2	100	0	0	0	50	0	100	0	0	0	0

Abbreviations: AM, Ampicillin; CP, Ciprofloxacin; CT, Ceftizoxime; IPM, Imipenem; GM, Gentamicin; SXT, Sulfamethoxazole-trimethoprim; NA, Nalidixic acid; CTX, Cefotaxime; FM, Nitrofurantoin; CAZ, Ceftazidime; CF, Cephalothin

Table 3. Prevalence and Antimicrobial Resistance of Gram Positive Uropathogens Isolated in this Study

	No.	AM	C	E	NOR	GM	V	FM	TE	SXT
<i>Staphylococcus saprophyticus</i>	26	92.31	61.54	92.31	30.77	30.77	7.7	7.7	92.31	77
<i>Staphylococcus aureus</i>	4	100	50	50	0	50	50	0	50	0
<i>Streptococcus group B</i>	10	30	20	30	40	70	80	0	70	80
<i>Enterococcus spp.</i>	10	40	40	70	100	100	0	0	100	100

Abbreviations: AM, Ampicillin; C, Chloramphenicol; E, Erythromycin; NOR, Norfloxacin; GM, Gentamicin; V, Vancomycin; FM, Nitrofurantoin; TE, Tetracycline; SXT, Sulfamethoxazole-trimethoprim

5. Discussion

This study shows the distribution of microbial species isolated from patients with UTI at a diagnostic laboratory center in Karaj, Iran and their susceptibility pattern to antimicrobial agents. Furthermore, we described the relationship between sex and isolated bacterial agents of UTI. Microbial infection of the UTI is one of the most common infectious diseases worldwide (19). Approximately 1 in 3 women will require antimicrobial treatment for a UTI before age 24, and 40% to 50% of women will suffer from UTI during their lifetime (20). The sex distribution of patients in our study is consistent with those in other studies (2, 16, 21) showing a statistically predominance of females (88.69% of the positive cultures). It is assumed that the short urethra in girls predisposes them to ascending infection. Furthermore, management of micturition in women is important. Management mistakes made by women include cleaning perineum forward from the anus to the vulva (2) that can cause urinary tract infection.

Sexual activity has been reported to influence higher prevalence of UTI in females (22). Considering the fact that most of infecting organisms are commensals of perianal and vaginal regions, emphasis on personal hygiene especially in females may be important in reducing the incidence of UTI (18). Males are less prone to UTIs possibly because of their longer urethra and the presence of antimicrobial substances in prostatic fluid (23). Our study indicates that *E. coli* is still the most common cause of UTI in Iran. This corresponds with the data obtained by other investigators (3, 15, 17). In addition, coagulase negative *Staphylococcus spp.* was the most common cause of UTI among Gram positive bacteria. Recent studies have revealed the importance of *coagulase negative Staphylococcus spp.* in urinary tract infections (24).

Our findings demonstrated that these pathogens play an important role in UTI as 11.3% of cases in our study were affected by these bacteria. *Klebsiella spp.* with the rate of 9.6% was the third species that caused UTI. In Our study, as with previous studies, *E. coli* demonstrated a very high microbial resistance to antibiotics. The analyzed results of antibiotic susceptibility test showed that *E. coli* was highly resistant to gentamicin (73.69%), ampicillin (73.69%), SXT (69.74%), and cephalothin (88.16%). This is similar to previous studies in USA (25) and Iran (17). On the other hand, very low resistance was detected to antibiotics such as ciprofloxacin (26.32%), imipenem (15.79%), nitrofurantoin (9.22%), and ceftizoxime (15.79%).

In the present study, the *coagulase negative Staphylococcus* showed 92.31% resistance to ampicillin, tetracycline, and erythromycin, respectively. Results showed that the best activity against these bacteria was achieved by nitrofurantoin, gentamicin, norfloxacin, and vancomycin that corresponds with other study in our region and other parts of the world (26, 27). *Klebsiella spp.* exhibited vary-

ing antibiotic resistance and showed higher resistance to gentamicin (63.64%) and ceftazidime (54.55%). Also *Klebsiella spp.* showed the lowest resistance to ciprofloxacin, ceftizoxime, and nalidixic acid (27.28%). Surprisingly, in the present study, the susceptibility of *Enterobacteriaceae* to gentamicin was low while other studies in Iran reported high sensitivity to this antibiotic (21, 23) which can be due to uncontrolled administration of the drug.

A significant increase in resistance of pathogenic strains to SXT, ampicillin, and cephalothin has been found worldwide (28). On the other hand, the effective drugs for UTI are ciprofloxacin, nitrofurantoin, and imipenem in our region. Low resistance to these drugs was observed because they were not easily available and are relatively expensive compared to others. Thus, these drugs could be considered as alternative options in empirical treatment of UTIs (16). Ciprofloxacin as an option for remedy to UTIs has been considered since its multiple mechanisms of action seem to have enabled it to retain potent activity against *E. coli*. Ciprofloxacin shows high level of activity against *E. coli* in UTI compared to other commonly used agents such as ampicillin and SXT (29). The data suggest that these drugs should be used cautiously to ensure that resistance rates to these substantial antibiotics for UTIs do not increase.

In the last decades, the number of reports about appearance of bacteria with antibiotic resistance has increased all over the world (3). This study, like others, shows clearly that there are significant geographic differences in the susceptibility of commonly used antimicrobials against UTIs (25, 30). As a result, accurate knowledge on local epidemiology and patterns of antimicrobial resistance in uropathogens are essential to design a clinically effective therapy for UTI. However in the current study we observed a high level and generalized resistance to commonly used antibiotics particularly among Gram negative isolates. This rate of resistance, especially to gentamicin is higher in our region than that observed in other countries such as Poland and Canada. Reducing the number of prescription for a particular antibiotic can lead to a decrease in resistance rates (31, 32). Studies like the present study are useful in determining local trends and risk factors for antimicrobial resistance and furthermore it is important to urge physicians and other health workers on performing antibiotic susceptibility test before blind antibiotic therapy.

With regard to continuous changing in causative bacteria isolated from patients with urinary tract infection and antibiotic sensitivity pattern, we recommend annual determination of bacterial sensitivity in populations as a guideline to be respected by physicians. Furthermore, considering susceptibility pattern of antibiotic agents for UTI therapy, particularly caused by *E. coli* as the most significant uropathogen, it seems that the drugs like ampicillin and SXT should be prescribed cautiously especially against

to those, demonstrate appropriate respond to other antibiotics such as ciprofloxacin and nitrofurantoin.

Acknowledgements

The authors would like to express their appreciations to Dr. hi and Dr. M. Basaki for review of the manuscript and suggestions. We also thank Mr. A. Shahed for his technical support.

Financial Disclosure

All authors declare that they have no conflict of interest.

Funding/Support

None declared.

Authors' Contribution

None declared.

References

- Hooton TM. A simplified approach to urinary tract infection. *Hosp Pract (Off Ed)*. 1995;**30**(2):23-30.
- Modarres s, Nassiri Oskoi N. Bacterial etiologic agents of urinary tract infection in children in the Islamic Republic of Iran. *East Mediterranean Health J*. 1997;**3**(2):290-5.
- Ranjbar R, Haghi-Ashtiani M, Jafari NJ, Abedini M. The prevalence and antimicrobial susceptibility of bacterial uropathogens isolated from pediatric patients. *Iranian J Pub Health*. 2009;**38**(2).
- Vasquez Y, Hand W. Antibiotic susceptibility patterns of community-acquired urinary tract infection isolates from female patients on the US (Texas)-Mexico border. *J Appl Res*. 2004;**4**(2):321-6.
- Behrman R, Kliegman R, Jenson H. Nelson Textbook of Pediatrics. Philadelphia: WB Saunders Company. Lippincott Williams & Wilkins; 1989.
- Feld LG, Mattoo TK. Urinary Tract Infections and Vesicoureteral Reflux in Infants and Children. *Pediatr Rev*. 2010;**31**(11):451-63.
- Inabo H, Obanibi W. Antimicrobial susceptibility of some urinary tract clinical isolates to commonly used antibiotics. *Afr J Biotechnol*. 2006;**5**(5):487-9.
- Al Sweih N, Jamal W, Rotimi VO. Spectrum and antibiotic resistance of uropathogens isolated from hospital and community patients with urinary tract infections in two large hospitals in Kuwait. *Med Princ Pract*. 2005;**14**(6):401-7.
- Kolawole A, Kolawole O, Kandaki-Olukemi Y, Babatunde S, Durawade K, Kolawole C. Prevalence of urinary tract infections (UTI) among patients attending Dalhatu Araf Specialist Hospital, Lafia, Nasarawa State, Nigeria. *Int J Medicinal Med Sci*. 2009;**1**(5):163-7.
- Mandell GL, Bennett JE, Dolin R. Principles and practice of infectious diseases. *Churchill Livingstone*. 2005:881-2.
- Hvidberg H, Struve C, Krogfelt KA, Christensen N, Rasmussen SN, Frimodt-Moller N. Development of a long-term ascending urinary tract infection mouse model for antibiotic treatment studies. *Antimicrob Agents Chemother*. 2000;**44**(1):156-63.
- Ramesh N, Sumathi CS, Kannan VR. Urinary Tract Infection and Antimicrobial Susceptibility Pattern of Extended Spectrum of Beta Lactamase Producing Clinical Isolates. *Adv Biol Res*. 2008;**2**(5-6):78-82.
- Jellheden B, Norrby RS, Sandberg T. Symptomatic urinary tract infection in women in primary health care. Bacteriological, clinical and diagnostic aspects in relation to host response to infection. *Scand J Prim Health Care*. 1996;**14**(2):122-8.
- Ronald A. The etiology of urinary tract infection: traditional and emerging pathogens. *Am J Med*. 2002;**113** Suppl 1A:14S-9S.
- Hryniewicz K, Szczypa K, Sulikowska A, Jankowski K, Betlejewska K, Hryniewicz W. Antibiotic susceptibility of bacterial strains isolated from urinary tract infections in Poland. *J Antimicrob Chemother*. 2001;**47**(6):773-80.
- Beyene G, Tsegaye W. Bacterial uropathogens in urinary tract infection and antibiotic susceptibility pattern in jimma university specialized hospital, southwest ethiopia. *Ethiop J Health Sci*. 2011;**21**(2):141-6.
- Amin M, Mehdinejad M, Pourdangchi Z. Study of bacteria isolated from urinary tract infections and determination of their susceptibility to antibiotics. *Jundishapur J Microbiol*. 2011;**2**(3):118-23.
- Oluremi B, Idowu A, Olaniyi J. Antibiotic susceptibility of common bacterial pathogens in urinary tract infections in a Teaching hospital in South-western Nigeria. *Afr J Microbiol Res*. 2011;**5**(22):3658-63.
- Ghotaslou R, Yaghoubi A, Sharify S. Urinary Tract Infections in Hospitalized Patients during 2006 to 2009 in. *J Cardiovasc Thorac Res*. 2010;**2**(1):39-42.
- Foxman B. Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. *Dis Mon*. 2003;**49**(2):53-70.
- Khameneh ZR, Afshar AT. Antimicrobial susceptibility pattern of urinary tract pathogens. *Saudi J Kidney Dis Transpl*. 2009;**20**(2):251-3.
- Adedeji B, Abdulkadir O. Etiology and antimicrobial resistance pattern of bacterial agents of urinary tract Infections in students of tertiary Institutions in Yola Metropolis. *Biol Res*. 2009;**3**(3-4):67-70.
- Farajnia S, Alikhani MY, Ghotaslou R, Naghili B, Nakhband A. Causative agents and antimicrobial susceptibilities of urinary tract infections in the northwest of Iran. *Int J Infect Dis*. 2009;**13**(2):140-4.
- Huebner J, Goldmann DA. Coagulase-negative staphylococci: role as pathogens. *Annu Rev Med*. 1999;**50**:223-36.
- Daza R, Gutierrez J, Piedrola G. Antibiotic susceptibility of bacterial strains isolated from patients with community-acquired urinary tract infections. *Int J Antimicrob Agents*. 2001;**18**(3):211-5.
- Khorshidi A, Moniri R, Shajari G. Antimicrobial Resistance in Gram-negative Bacilli Isolated from Urinary Tracts Infections. *Arch Razi Ins*. 2003;**55**.
- Sheikholeslami Z, Hassanshahi G. The frequency of coagulase negative staphylococci urinary infections with antimicrobial resistance pattern in Rafsanjan. *Pak J Med Sci*. 2010;**26**(1):107-10.
- Gupta K, Hooton TM, Stamm WE. Increasing antimicrobial resistance and the management of uncomplicated community-acquired urinary tract infections. *Ann Intern Med*. 2001;**135**(1):41-50.
- Abelson Storby K, Osterlund A, Kahlmeter G. Antimicrobial resistance in Escherichia coli in urine samples from children and adults: a 12 year analysis. *Acta Paediatr*. 2004;**93**(4):487-91.
- Karlowsky JA, Kelly LJ, Thornsberry C, Jones ME, Sahm DF. Trends in antimicrobial resistance among urinary tract infection isolates of Escherichia coli from female outpatients in the United States. *Antimicrob Agents Chemother*. 2002;**46**(8):2540-5.
- Garau J, Xercavins M, Rodriguez-Carballeira M, Gomez-Vera JR, Coll I, Vidal D, et al. Emergence and dissemination of quinolone-resistant Escherichia coli in the community. *Antimicrob Agents Chemother*. 1999;**43**(11):2736-41.
- Natsch S, Conrad C, Hartmeier C, Schmid B. Use of amoxicillin-clavulanate and resistance in Escherichia coli over a 4-year period. *Infect Control Hosp Epidemiol*. 1998;**19**(9):653-6.

SID



سرویس های
ویژه



سرویس ترجمه
تخصصی



کارگاه های
آموزشی



بلاگ
مرکز اطلاعات علمی



عضویت در
خبرنامه



فیلم های
آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



مباحث پیشرفته یادگیری عمیق؛
شبکه های توجه گرافی
(Graph Attention Networks)



کارگاه آنلاین آموزش استفاده از
وب آوساینس



کارگاه آنلاین مقاله روزمره انگلیسی