

QUANTITATIVE SIZE ASSESSMENT OF THE LUMBAR SPINAL CANAL BY COMPUTED TOMOGRAPHY

M. Midia¹ and Z. Miabi^{2*}

1) Department of Radiology, McMaster University, Ontario, Canada

2) Department of Radiology, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran

Abstract- By determining normal ranges of spinal canal diameters we can make early diagnosis in persons who have lower diameters of spinal canal. These persons are predisposed to spinal canal stenosis that is a major cause of spinal radiculopathies. In different studies performed in several countries, minimum and maximum ranges of spinal canal diameters were different for each population. In this study, we tried to determine the mean values of normal spinal canal diameters and areas in Tabriz and its suburb. 39 healthy, young to mid-age cases were selected. Our study was focused on L3-L4 and L4-L5. The following parameters were measured: the area of cross-section of the vertebral body, the area of cross-section of the dural sac, interarticular diameter, interligamentous diameter, antero-posterior diameter of the lumbar canal, inter-pedicular diameter, and the area of cross-section of the vertebral canal. A correlation between the parameters studied and the height of subjects was significant for interligamentous diameter (for L3/L4 and L4/L5) and interarticular diameter (only at L3/L4), cross-section area of the vertebrae (both L3 and L4), cross-section area of vertebral canal (only at L5 level), area of dural sac (at L3/L4 and L4). It was suggested that these diameters and areas should be interpreted as a function of height of the subject. Most of diameters studied had smaller means than those in previous studies. This can be attributed to differences between populations and it can be interpreted as predisposition to spinal canal stenosis in our population.

© 2007 Tehran University of Medical Sciences. All rights reserved.

Acta Medica Iranica, 45(5): 377-382; 2007

Key words: Spinal canal, radiculopathy, canal stenosis

INTRODUCTION

In different studies performed in several counties, minimum and maximum ranges of spinal canal diameters were different for each population. By determining normal ranges of spinal canal diameters we can make early diagnosis in persons who have lower diameters of spinal canal. These persons are predisposed to spinal canal stenosis, which is a major cause of spinal radiculopathies.

Low back pain is a major cause of disability, poor quality of life and societal costs. For example, in the United States: (1) the annual societal cost of back pain is estimated to be between \$20 and \$50 billion, (2) back symptoms are the most common cause of disability in patients under 45 years of age, (3) approximately 1 percent of the U.S. population is chronically disabled because of back pain.

The structure of the lumbar spine is complex. To diagnose and treat this area effectively, one must have a clear knowledge of the normal anatomy. Following is an introduction of anatomy of lumbar spine.

Spinal canal stenosis may involve the central spinal canal, its lateral recess (neural canal), the intervertebral foramina, or all three components.

Received: 14 Feb. 2007, Revised: 16 May 2007, Accepted: 10 Sep. 2007

*** Corresponding Author:**

Zinat Miabi, Department of Radiology, School of Medicine, Medical Sciences/Tabriz University, Tabriz, Iran

Tel: +98 411 3011001

Fax: +98 411 3011001

E-mail: miabiz7@yahoo.com

Spinal canal stenosis may be developmental or, as is more common, an acquired disease from numerous causes (3, 4)

Congenital lumbar spinal canal stenosis is more often seen in males than in females, usually in their second or third decade (5). Spinal stenosis syndrome affects mainly patients at their 5th-6th decades of life. There is a strong debate in the literature whether patients with spinal stenosis should be operated or treated conservatively (6).

Differences in dimensions of male and female specimens were not found to be statistically significant (7). At all levels (L1- L5) the transverse diameters of the lumbar spinal canal were approximately 1- 1.5 mm higher in males than in females (8).

Pathological changes can occur in the diameters of the lumbar spinal canal. Therefore, assessing the canal size an important diagnostic procedure (7).

The patients without lumbar symptoms had wider foramina and sagittal diameters in S1 than those with lumbar symptoms (8).

In this study, we tried to determine the mean values of normal spinal canal diameters and areas in a healthy, young to mid-age population from Tabriz and its suburb. There are many studies (using different modalities) about measurement of spinal canal diameters in foreign in foreign populations (1, 5, 6, 8, 10). To our knowledge, there is no report of normal spinal canal diameters in Iranian populations.

MATERIALS AND METHODS

The subjects

This study was performed on 39 subjects (16 males and 23 female) with age ranging from 18 to 40. Study cases were selected from patients referred to perform CT scan of other parts of the body and had no low back pain or other problem attributable to lumbar spine.

Technique

Examinations were made using a GE CT-MAX II scanner. Our study was focused on L3-L4 and L4-L5 levels because these levels are studied most frequently in CT-scan examinations. Meanwhile,

examination of L5-S1 level is difficult because of technical limitations (angle of gantry in our scanner was limited to 20 degree). For each, we took a lateral scout view of lumbar spinal canal. Then cuts were made perpendicular to the posterior wall of the vertebral body at each level. 2-mm thickness was used. For each level (L3-L4 and L4-L5) we acquired a cut through highest part of intervertebral foramen, a cut through the disc, and a cut through the middle third of the lower vertebral body. Images were reconstructed in high resolution and printed on film with a window for bony structures (level: 300, width: 1500) and a window for soft tissues (level: 80, width: 1000) and transferred to a PC using an HP transparent flatbed scanner. Measurements were made in adobe Photoshop and converted to actual sizes (mm) using the scale printed with each image.

Parameters studied

Cuts below the pedicles were made through the highest part of the intervertebral foramina. The following parameters were measured: the area of cross-section of the vertebral body (SCV-L3 and SCV-L4); the area of cross-section of the dural sac (SF-L3 and SF-L4).

These cuts were made at the level of the middle of the disc and are concerned with the intervertebral articulation. The following parameters were measured: interarticular diameter (DIA-L3/L4 and DIA-L4/L5); interligamentous diameter (DIL-L3/L4 and DIL-L4/L5); area of cross-section of the dural sac (SE-L3/L4 and SF-L4/L5). Measurements were made as greatest diameter between the internal limits of two articulations for DIA, and the distance between internal borders of the soft parts of the articulations (capsules and ligamentum flavum) on the line joining the articulations for DIL. These cuts were made through the third of the vertebral body and show a complete vertebral body ring. The following parameters were measured: antero-posterior diameter of the lumbar canal (DAP-L4 and DAP-L5); inter-pedicular diameter (DIP-L4 and DIP-L5); the area of cross-section of the vertebral canal (SC-L4 and SC-L5). Measurements were made as the distance from the mid-point of the posterior wall of the vertebral body to the anterior border of the point of union of the two laminae DAP, and the

greatest distance between the inner borders of the two pedicles for DIP. Measurements of SCV, DAP, DIP, SC, and DIA were made on images printed using bony structures window and level, whereas those for SF and DIL were made on images with soft tissue window.

Statistical method

The presence of any significant correlation between the parameters that were measured and the height of the subjects was sought. Coefficients of correlation and p-values corresponding to these coefficients were calculated using SPSS Windows software. A coefficient of correlation was considered significant (not occurring by chance) when the corresponding *P* value is less than or equal to 0.050 (5%).

RESULTS

Mean age and height of patients were 30 ± 6 year and 167 ± 9.15 cm, respectively. Mean area of vertebral body of L3 and L4 was 1515 ± 254.6 mm² and 1470 ± 255.4 mm², respectively. Mean area of dural sac of L3 and L4 was 142 ± 30.7 mm² and 128 ± 36.4 mm², respectively (Table 2).

Mean of interarticular diameter (DIA), interligamentous diameter (DIL), antero-posterior diameter (DSP), interpedicular diameter (DIP) and area of spinal of canal (SC) were showed in table 2.

Significant liner correlation was found between height and area of vertebral body of L3 (*P* = 0.008) and L4 (*P* = 0.007), area of spinal canal of L5 (*P* = 0.012), interarticular diameter of L3/L4 (*P* = 0.011), area of dural sac of L3/L4 (*P* = 0.033), interligamentous diameter of L3/L4 (*P* = 0.001) and L4/L5 (*P* = 0.046), respectively (Table 3).

DISCUSSION

The diameters

A correlation between the diameters studied and the height of subjects was significant for interligamentous diameter (DIL-L3/L4 and DIL-L4/L5) and interarticular diameter only at L3/L4 level (DIA-L3/L4) (Table 3). This is in contrary with previous studies. Gouzien *et al.* concluded that there

Table 1. The significant correlation between the measured parameters and the age of subject

	Mean	SD	Minimum value	Maximum value
Age (year)	30	6.0	18	40
Height (cm)	167	9.15	155	187
Subpedicular cut:				
SCV-L3 (mm ²)	1515	254.6	1133	2164
SCV-L4 (mm ²)	1470	255.4	1123	2139
SF-L3 (mm ²)	142	30.7	87	218
SF-L4 (mm ²)	128	36.4	39	221
Cut through disc:				
DIA-L3/L4 (mm)	20	3.2	11	28
DIA-L4/L5 (mm)	22	3.7	12	28
DIL-L3/L4 (mm)	12	2.9	4	17
DIL-L4/L5 (mm)	13	3.3	7	20
SF-L3/L4 (mm ²)	133	39.2	68	247
SF-L4/L5 (mm ²)	133	41.6	49	256
Cut through pedicles and lamina:				
DAP-L4 (mm)	14	2.7	8	23
DAP-L5 (mm)	15	2.5	10	23
DIP-L4 (mm)	23	2.1	20	27
DIP-L5 (mm)	30	4.7	22	44
SC-L4 (mm ²)	229	45.8	120	322
SC-L5 (mm ²)	281	70.0	137	490

Abbreviations: SCV: area of vertebral body; SF: area of dural sac; DIA: interarticular diameter; DIL: interligamentous diameter; DSP: antero-posterior diameter; DIP: interpedicular diameter; SC: area of spinal of canal.

Table 3. The significant correlation between the measured parameters and the height of subject

Parameter	Mean	SD	P-value	Regression line $y=ax+b$
SCV-L3 (mm ²)	1515	254.6	0.008	$y=10.76x-280.2$
SCV-L4 (mm ²)	1470	255.4	0.007	$y=10.82x-336.2$
SC-L5v	281	70.0	0.012	$y=2.79x-184.7$
DIA-L3/L4 (mm)	20	3.2	0.011	$y=0.13x-1.097$
SF-L4 (mm ²)	133	39.2	0.033	$y=1.28x-80.1$
SF-L4 (mm ²)	128	36.4	0.050	$y=106x-49.36$
DIL-L3/L4 (mm)	12	2.9	0.001	$y=0.15x-13.36$
DIL-L4/L5 (mm)	13	3.3	0.046	$y=0.1x-3.90$

Abbreviations: SCV: area of vertebral body; SC: area of spinal canal; DIA: interarticular diameter; SF: area of dural sac; DIL: interligamentous diameter.

Table 4. Comparison between mean measurements from the present study and previous reports

Parameter	Our study 19988-	Gouzien <i>et al.</i> ⁵	Hwan-Mp Lee ⁶	Wilmink <i>et al.</i> ¹⁰ 1988-
	Iran	1990-France	1995-Korea	Netherlands
SCV-L4 (mm ²)	1470	1318	-	-
SF-L4 (mm ²)	142	169	-	-
SF-L4 (mm ²)	128	168	-	-
DIA-L3/L4 (mm)	20	22	-	-
DIA-L4/L5 (mm)	22	24	-	-
DIL-L3/L4 (mm)	12	12	-	10
DIL-L4/L5 (mm)	13	15	-	14
SF-L3/L4 (mm ²)	133	155	-	-
SF-L4/L5 (mm ²)	133	157	-	-
DAP-L4 (mm)	14	17	14	14
DAP-L5 (mm)	15	18	14	14
DIP-L4 (mm)	23	26	23	22
DIP-L5 (mm)	30	31	26	25
SC-L4 (mm ²)	229	288	-	-
SC-L5 (mm ²)	281	350	-	-

Abbreviations: SCV, area of vertebral body; SFC, area of dural sac; DIA, interarticular diameter; DIL, interligamentous diameter; DAP, antero-posterior diameter; DIP, interpedicular diameter; SC, area of spinal canal.

is a significant correlation between height and only interpedicular diameter (Table 5) (10). In our study there was not a significant correlation between interpedicular diameter and height of subjects.

Although it is stated in previous literature that measurement of interligamentous diameter is important in the study of a canal narrowed by degenerative disease, previous studies were not able

to demonstrate a significant correlation between height and DIL (10). In our study this significant correlation was present for both levels (DIL-L3/L4 and DIL-L4/L5).

Most of diameters studied had smaller means than those in previous studies (Table 4) (1, 10, 11, 15).

Table 5. Significance of correlation between the measured parameter and the height of subjects

Parameter	Our series (1998)	Gouzien <i>et al.</i> (1990) ⁵
SCV-L3 (mm ²)	Significant ($P = 0.008$)	-
SCV-L4 (mm ²)	Significant ($P = 0.007$)	Significant ($P = 0.010$)
SF-L3 (mm ²)	Non-significant ($P = 0.139$)	-
SF-L4 (mm ²)	Significant ($P = 0.050$)	-
DIA-L3/L4 (mm)	Significant ($P = 0.011$)	-
DIA-L4/L5 (mm)	Non-significant ($P = 0.069$)	-
DIL-L3/L4 (mm)	Significant ($P = 0.001$)	Non-significant ($P = 0.100$)
DIL-L4/L5 (mm)	Significant ($P = 0.046$)	Non-significant ($P = 0.500$)
SF-L3/L4 (mm ²)	Significant ($P = 0.033$)	Significant ($P = 0.010$)
SF-L4/L5 (mm ²)	Non-significant ($P = 0.347$)	-
DAP-L4 (mm)	Non-significant ($P = 0.494$)	Non-significant ($P = 0.670$)
DAP-L5 (mm)	Non-significant ($P = 0.241$)	Non-significant ($P = 0.100$)
DIP-L4 (mm)	Non-significant ($P = 0.133$)	Non-significant ($P = 0.001$)
DIP-L5 (mm)	Non-significant ($P = 0.127$)	Non-significant ($P = 0.001$)
SC-L4 (mm ²)	Non-significant ($P = 0.155$)	Non-significant ($P = 0.001$)
SC-L5 (mm ²)	Significant ($P = 0.012$)	Significant ($P = 0.010$)

Abbreviations: SCV: area of vertebral body; SF: area of dural sac; DIA: interarticular diameter; Dil: interligamentous diameter; DAP: antero-posterior diameter; DIP: interligamentous diameter; SC: area of spinal canal.

The anteroposterior diameter (DAP) which is essential in diagnosis of canal stenosis, has greater mean in some of previous studies (10). It is stated that the lowest possible normal limit for DAP appears to be 12 mm (9, 10), but we had a minimum value of 8 mm for DAP-L4, 10 mm for DAP-L5, and 6 cases had DAP value less than 12. This can be due to differences between populations in our study and previous studies, or differences between techniques of measurement (less probable particularly for study performed by Gouzien *et al.*).

The areas

A significant correlation was present between cross-section area of the vertebrae (both SCV-L3 and SCV-L4) and height of subjection. For cross-section area of vertebral canal there was a significant correlation only at L5 level (SC-L5) (Table 3).

Relationship between area of canal and that of vertebral body have been established in previous literature. Jones and Thomson have found a ratio with mean value of 0.16 and SD of 0.033 (by conventional radiology) (12). Gouzien *et al.* have found a mean value of 0.22 and SD of 0.05 (by CT scan) (10). In our series we found a mean value of 0.158 and a SD of 0.033 ($P = 0.033$) (closer to Hones and Thomson).

The area of dural sac was significantly correlated with height at L3/L4 and L4 levels (SF-L3/L4 and SF-L4) (Table 3). It has a fundamental importance in the examination of spinal canal. It is more logical to consider the area of dural sac in relation to the size of the canal, because a small sac within a canal that is narrow may be asymptomatic but a large sac in a normal size canal may be pathological.

The ratio between area of dural sac and area of canal has been studied previously. Gouzien *et al.* has found a mean value of 0.58 and SD 0.10.⁵ We found a mean value of 0.558 and a SD of 0.103 ($P < 0.001$). It must be mentioned that measurements made by CT scan are static and do not take into account any changes that results from alteration in position and posture. In addition there is other soft tissue structures within spinal canal such as epidural fat and ligamentum flavum and the joint capsules. In conclusion, a study of lumbar spinal canal at L3 to L5 level was performed on 39 cases. A significant

correlation was found between height of subjects and cross-sectional areas of vertebral body, spinal canal, and dural sac and interligamentous and interarticular diameters.

Some diameters that are important in examination of spinal canal stenosis was not significantly correlated with height of subjects (DAP-L4 and DAP-L5). It is important to note that in our study antero-posterior diameter of spinal canal had a mean value slightly lesser than previous studies in other populations (10). Meanwhile, (and maybe more important) in our series we had a considerable number of subjects with DAP value lesser than accepted minimum value in other populations (10, 11, 15). This can be attributed to differences between populations and it can be interpreted as predisposition to spinal canal stenosis in our population. Most of the other parameters had a lesser mean value than other population. This can also be attributed to differences between population. It is suggested that there can be significant correlation between diameters and areas of spinal canal at upper levels and height of subjects.

Conflict of interests

The authors declare that they have no competing interests.

REFERENCES

1. Amonoo-Kuofi HS, Patel PJ, Fatani JA. Transverse diameter of the lumbar spinal canal in normal adult Saudis. *Acta Anat (Basel)*. 1990;137(2):124-128.
2. Borenstein, DG; Wiesel, SW: Low back pain. Philadelphia PA, WB Saunders, 1989; P. 3-9.
3. Chafets, N; Rothman, SLG; Genant, HK; Kaiser JA: Computed tomography and magnetic resonance imaging of the spine. In: Moss, A; Gamsu, G; Genant HK (ed): *Computed tomography of the body*. 2nd Ed. Philadelphia PA, WB Saunders, 1992; P. 496-500.
4. Shabat S, Folman Y, Leitner Y, Fredman B, Gepstein R. Failure of conservative treatment for lumbar spinal stenosis in elderly patients. *Arch Gerontol Geriatr*. 2007 May-Jun; 44(3):235-241.

Assessment of size of lumbar spinal canal by CT

5. Ebraheim NA, An HS, Xu R, Ahmad M, Yeasting RA. The quantitative anatomy of the cervical nerve root groove and the intervertebral foramen. *Spine*. 1996 Jul 15; 21(14):1619-1623.
6. Tacar O, Demirant A, Nas K, Altındağ O. Morphology of the lumbar spinal canal in normal adult Turks. *Yonsei Med J*. 2003 Aug 30; 44(4):679-685.
7. Santiago FR, Milena GL, Herrera RO, Romero PA, Plazas PG. Morphometry of the lower lumbar vertebrae in patients with and without low back pain. *Eur Spine J*. 2001 Jun; 10(3):228-233.
8. Gepstein R, Folman Y, Sagiv P, Ben David Y, Hallel T. Does the anteroposterior diameter of the bony spinal canal reflect its size? An anatomical study. *Surg Radiol Anat*. 1991; 13(4):289-291.
9. Gouzien P, Cazalbou C, Boyer B, Darodes de Tailly P, Guenec Y, Sénécaïl B. Measurements of the normal lumbar spinal canal by computed tomography. Segmental study of L3-L4 and L4-L5 related to the height of the subject. *Surg Radiol Anat*. 1990; 12(2):143-148.
10. Jones, RA; Thomson, JL: The narrow lumbar canal. A clinical and radiological review. *Journal of Bone and Joint Surgery (British)* 50:595-605; 1968.
11. Lee HM, Kim NH, Kim HJ, Chung IH. Morphometric study of the lumbar spinal canal in the Korean population. *Spine*. 1995 Aug 1; 20(15):1679-1684.
12. Lee, B.C.P; Kazam, E; Newman, A.D.: Computed tomography of the spine and spinal cord. *Radiology* 128:95-102; 1978.
13. Rao, K.C.V.G.; Williams L.P.: MRI and CT of the spine. Baltimore MA, Williams & Wilkins, 1994; P. 188-198.
14. Wilmink JT, Korte JH, Penning L. Dimensions of the spinal canal in individuals symptomatic and non-symptomatic for sciatica: a CT study. *Neuroradiology*. 1988; 30(6):547-550.
15. Shabat S, Folman Y, Leitner Y, Fredman B, Gepstein R. Failure of conservative treatment for lumbar spinal stenosis in elderly patients. *Arch Gerontol Geriatr* 2006 Jul 24.
16. Ebraheim NA, An HS, Xu R, Ahmad M. Yeasting RA. The quantitative anatomy of the cervical nerve root groove and the intervertebral foramen. *Spine* 1996 Jul 15; 21(14): 1619-1623.
17. Tacar O, Demirant A, Nas K, Altındağ O. Morphology of the lumbar spinal canal in normal adult Turks. *Yonsei Med J* 2003; 44(4): 679-685.
18. Santiago FR, Milena GL, Herrera RO, Romero PA, Plazas PG. Morphometry of the lower lumbar vertebrae in patients with and without low back pain. *Eur Spine J* 2001; 10(3): 228-233.

Surf and download all data from SID.ir: www.SID.ir

Translate via STRS.ir: www.STRS.ir

Follow our scientific posts via our Blog: www.sid.ir/blog

Use our educational service (Courses, Workshops, Videos and etc.) via Workshop: www.sid.ir/workshop