

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی




عضویت در خبرنامه



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



تازه آموزش
مباحث پیشرفته یادگیری عمیق؛ شبکه های توجه گرافی (GAN)

مباحث پیشرفته یادگیری عمیق؛
شبکه های توجه گرافی
(Graph Attention Networks)



تازه آموزش
آموزش استفاده از وب آو ساینس

کارگاه آنلاین آموزش استفاده از
وب آو ساینس



کارگاه آنلاین مقاله روزمره انگلیسی

Evaluation of SD-208, a TGF- β -RI Kinase Inhibitor, as an Anticancer Agent in Retinoblastoma

Puran Fadakar¹, Abolfazl Akbari², Fariba Ghassemi³, Gholam Reza Mobini⁴, Masoumeh Mohebi³, Manzar Bolhassani⁵, Hoda Abed Khojasteh³, and Mansour Heidari^{5,6}

¹ Department of Biochemistry and Genetics, Molecular and Medicine Research Center, Arak University of Medical Sciences, Arak, Iran

² Colorectal Research Center, Iran University of Medical Sciences, Tehran, Iran

³ Eye Research Center, Farabi Eye Hospital, Tehran University of Medical Sciences, Tehran, Iran

⁴ Medical Plants Research Center, Shahrekord University of Medical Sciences, Shahrekord, Iran

⁵ Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

⁶ Experimental Medicine Research Center, Tehran University of Medical Sciences, Tehran, Iran

Received: 25 Nov. 2015; Accepted: 14 Jan. 2016

Abstract- Retinoblastoma is the most common intraocular tumor in children resulting from genetic alterations and transformation of mature retinal cells. The objective of this study was to investigate the effects of SD-208, TGF- β -RI kinase inhibitor, on the expression of some miRNAs including a miR-17/92 cluster in retinoblastoma cells. Prior to initiate this work, the cell proliferation was studied by Methyl Thiazolyl Tetrazolium (MTT) and bromo-2'-deoxyuridine (BrdU) assays. Then, the expression patterns of four miRNAs (18a, 20a, 22, and 34a) were investigated in the treated SD-208 (0.0, 1, 2 and 3 μ M) and untreated Y-79 cells. A remarkable inhibition of the cell proliferation was found in Y-79 cells treated with SD-208 versus untreated cells. Also, the expression changes were observed in miRNAs 18a, 20a, 22 and 34a in response to SD-208 treatment ($P < 0.05$). The findings of the present study suggest that the anti-cancer effect of SD-208 may be exerted due to the regulation of specific miRNAs, at least in this particular retinoblastoma cell line. To the best of the researchers' knowledge, this is the first report demonstrating that the SD-208 could alter the expression of tumor suppressive miRNAs as well as oncomiRs *in vitro*. In conclusion, the present data suggest that SD-208 could be an alternative agent in retinoblastoma treatment.

© 2016 Tehran University of Medical Sciences. All rights reserved.

Acta Med Iran, 2016;54(6):352-358.

Keywords: miRNA; Retinoblastoma; SD-208; Y-79 cell line

Introduction

Retinoblastoma (Rb), the most common pediatric cancer of the eye, exists in sporadic and heritable forms (1). Two kinds of this cancer, bilateral (15% inheritable) and unilateral (85% non-heritable) have been reported. This malignancy occurs as part of a familial cancer syndrome; however, the main ratio of retinoblastoma arises with no preceding family history (2). Retinoblastoma is first cancer confirmed to have a genetic origin (2,3). Although genetic changes in Rb gene have been frequently reported in retinoblastoma, the genetic alteration in other biomolecules and genes such as MYCN oncogene appears to be important in its pathogenesis (4-8). Studies have shown that

inappropriate expression or loss of RB function is associated with other human cancers (9-13).

Today, it is well known that numerous biomolecules such as micro RNAs (miRNAs) and long non-coding RNAs (lncRNAs) are implicated in the regulation of various human cancer genes (14-19). For instance, these molecules play a vital role in tumor suppressor networks such as RB gene (20). The expression profile and biological significance of miRNAs depend on the target genes and kind of malignancy (tumor microenvironment). Many researchers determined that a specific group of miRNAs is highly expressed in multiple tumor types and turn on some oncogenes while overexpression of another group leads to the activation of some tumor suppressor genes. It is quite

Corresponding Author: M. Heidari

Department of Medical Genetics, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran
Tel: +98 21 88953005, Fax: +98 21 88953005, E-mail address: mheidari@sina.tums.ac.ir

understandable, therefore, that these molecules could be used as biomarkers and targets for the cancer diagnosis and therapy, respectively.

In spite of the fact that the survival rate of retinoblastoma is about 98%, chemotherapeutic resistance and various side effects of current therapies are considered the main challenges for this disease. Studies have shown that the pretreatment evaluation of retinoblastoma cases is individualized and directed by the chosen mode of therapy. The present study, for the first time, investigated the effect of a TGβRI kinase inhibitor, SD-208, on the regulation of four miRNAs 18a, 20a, 22 and 34a in retinoblastoma

Materials and Methods

Cell culture

The human retinoblastoma Y-79 cell line was obtained from National Cell Bank of Iran (NCBI) affiliated to Pasteur Institute (Tehran, Iran). The cells were grown in suspension at a concentration of 105-106 cells/ml in RPMI-1640 medium (Gibco; Germany) containing 10% fetal bovine serum (FBS) (Gibco; Germany), 2 mM L-glutamine, 1 mM sodium pyruvate, and antibiotics (100 U/ml of penicillin and 100 µg/ml of streptomycin). Cell culture condition designed at 37°C in a humidified atmosphere of 5% CO₂-95% air.

SD-208 treatment

The treatment of cells with SD-208 (Sigma-Aldrich, Belgium) was conducted as previously described (21). Briefly, SD-208 was dissolved in dimethyl sulfoxide (DMSO) with a concentration of 20 µM and stored at -20°C. The Y-79 cells in each experiment were treated using a different range of SD-208 concentrations (as explain in the below sections).

MTT assay

The cytotoxicity effect of SD-208 was measured by the MTT assay as previously described (21,22). Briefly, Y-79 cells were seeded in 96-well flat-bottom microtitration plates (SPL Life Sciences; South Korea) at a density of 4×10⁴ cells/well (200 µl media/well). After reaching to ~85% confluency, the cells were treated with various concentrations of SD-208 (0.0, 1, 2, 3 µM). In all in vitro experiments, control cells were incubated with dimethyl sulfoxide (DMSO) alone (with a final concentration 0.2%).

BrdU assay

The BrdU assay was performed using BrdU ELISA

kit as previously described (21). Briefly, the Y-79 cells were seeded in 96-well flat-bottom microtitration plates at a density of 4×10⁴ cells/well (100 µl media/well). Then, the cells were treated with SD-208 (0.0, 1, 2, 3 µM) and the cell apoptosis calculated as previously described (21).

Total RNA extraction

The isolation of total RNA from cells with a cell density of 80% confluent was conducted using Trizol® total RNA isolation reagent (Gibco BRL, Life Technologies, Gaithersburg, MD, USA) according to the manufacturer's instructions. To remove any genomic contamination, the total RNA was treated with RNase-free DNase I and stored at -70°C for further tests.

miRNA expression analysis by Real-time q-PCR

Four miRNAs (including miR-18a, 20a, 22, 34a) genes involving in retinoblastoma were selected from the Sanger Center miRNA Registry at <http://www.sanger.ac.uk/Software/Rfam/mirna/index.shtm>.

The microRNAs expression was analyzed by real-time quantitative polymerase chain reactions (qPCR) using the SYBR Green method by the specific primers for mature forms of aforementioned micro RNAs as per the manufacturer's instructions (Parsgenom, Iran). The real-time q-PCR carried out by poly (A) tailing method relying on polyadenylation, followed by cDNA synthesis using a universal oligo(dT) primer. RNU6B was used as an endogenous (internal) control, and the data were normalized to the expression level of this housekeeping gene.

Our protocol consisted of two steps: reverse transcription (RT) and real-time PCR. First, RNA was transcribed to complementary DNA (cDNA) using the poly-adenylation procedure. Briefly, a 20 µl reaction including 2 µg total RNA, 2 µl of 10X reaction buffer, 2 µl of 10 mM ATP and 1 unit of Poly (A) polymerase was incubated at 38 °C for 30 min, followed by enzyme inactivation at 65 °C for 3 min. After polyadenylation, the cDNA was synthesized using 2 µg of polyadenylated RNA (RNA poly A tail) and miR-cDNA Syn specific primers in a reverse transcription reaction. This reaction was performed in a volume of 10 µl containing 1 µl of the polyadenylation reaction product, 1 µl of 0.5 mM universal RT primer, 1 µl of 10 mMdNTP, 0.5 µl of RT enzyme in. The reaction was incubated at 42 °C for 60 min and then terminated by heating at 85 °C for 5 min.

Diluted cDNAs then were amplified in a 20µl reaction containing SYBER Green Mastermix, forward

and reverse miR-specific primers (each 1 μ l) and DEPC-treated distilled water by 35 cycles of PCR amplification under the following conditions: denaturing at 95 °C for 1 min, annealing at 60 °C for 1 min, and extension at 72 °C for 1 min. Real-time PCR was performed on a Bio-Rad CFX96 Real-Time PCR System.

All the reactions were performed in triplicate. Specificity of primers was verified by observing a single peak dissociation curve for each run. The threshold cycle (CT) is defined as the fractional cycle number at which the fluorescence passes the fixed threshold. CT values were converted into total copy numbers using a standard curve. The absence of contamination was verified using non-template controls.

Statistical analysis

Statistical analysis for multiple comparisons of differences between treatment groups was performed with SPSS software. The standard error of means was computed, and analysis of variance (ANOVA, Tukey post tests) completed via GraphPad Prism 5.0 software. Gene expression analysis was performed by $2^{-\Delta\Delta Ct}$ method via GraphPad Prism 5.0 software. $P < 0.05$ were considered as statistically significant.

Results

SD-208 decreases proliferation of and induces apoptosis in Y-79 cells

To determine the inhibitory effect of SD-208 on cell proliferation in Y-79 cells, the cell growth was analyzed by performing MTT assay with various conditions. The data resulted from the present study clearly showed that SD-208 treatment significantly reduces the viability of Y-79 cells in a dose- and time-dependent manner. Figure 1 shows the cell response to different drug concentrations (0.0, 1, 2, 3 μ M) for a time course of 24, 48, and 72 h. The results of cell function assays revealed a significant change in the cell viability in the treated versus untreated Y-79 cells ($P < 0.05$), especially at 48 h and 72 h.

Moreover, to examine whether SD-208 can promote apoptosis in Y-79 cells, we analyzed apoptotic index using a BrdU experiment in the SD-208 treated cells compared to controls (Figure 2). The apoptotic findings revealed a significant change in cell death in the treated versus control cells ($P < 0.05$) at 48 h and 72 h.

The obtained results show that the cytotoxicity and apoptosis effects of SD-208 on retinoblastoma cell line are time and dose-dependent. Besides, its apoptotic effect like its toxicity seems to follow a time and dose-

dependent manner.




Figure 1. Effect of SD-208 on the Y-79 cell viability. Y-79 cells were treated by 0.0, 1, 2 and 3 μ M for different time courses. The cell toxicity was examined by MTT assay and data was reported as the percentage change in comparison with the controls. Comparison of the cell viability of Y-79 cells in different concentrations of SD-208 to controls was analyzed by one-way ANOVA test. Results were expressed as mean \pm SEM from three independent experiments. $P < 0.05$ was considered as statistically significant




Figure 2. Impact of SD-208 on the Y-79 cell proliferation. The Y-79 cells were treated by 0.0, 1, 2 and 3 μ M for different time courses. Cell apoptosis was examined by BrdU assay and analysis of one-way ANOVA was used to compare the cell apoptosis. Results were expressed as mean \pm SEM from three independent experiments. $P < 0.05$ was considered as statistically significant

SD-208 regulates miRNA expression in retinoblastoma

To study the ability of the SD-208 as a potential anticancer agent via changing oncogene/tumor suppressor miRNAs, the researchers partially evaluated its effects on some miRNA expression. As Figure 3

shows, SD-208 could significantly upregulate miR-18a, 22a, and 34a ($P < 0.05$) while downregulating miR-20a ($P < 0.05$) expression in retinoblastoma cells.




Figure 3. Effect of SD-208 on miRNA expression. Evaluation of the effects of SD-208 on miRNA expression was completed by Real-time q-PCR. SD-208 could significantly upregulate miR-18a, 22a, and 34a ($P < 0.05$) while downregulating miR-20a ($P < 0.05$) expression

Discussion

It is well established that TGF β signaling pathway plays a critical role in the normal and abnormal cellular processes. In this study, we focused on the involvement of miRNAs in the anticancerous effect of SD-208. We found that the kinase inhibitor, SD-208, was able to inhibit TGF- β -evoked proliferation in retinoblastoma via regulation of cancer-related miRNAs. A few reports suggested that SD-208 could not be an effective chemotherapeutic agent in at least some human cancers (21,23). Although in a study it was shown that SD-208 could decrease the tumor angiogenesis in athymic nude mice, it was not able to inhibit mammary tumor growth and metastasis. Most recently, Akbari *et al.*, reported no significant reduction in cell proliferation and angiogenesis in human colon adenocarcinoma cells treated with SD-208 (21).

In contrast, to work described above, accumulated data have demonstrated the potential anti-cancer activities of SD-208 (24-26) in distinct malignancies such as pancreatic (27,28), glioma (25) and melanoma (24). Mohammad KS defined SD-208 as promising novel agents for the treatment of human malignant glioma and other conditions associated with pathological TGF β activity. Also, Uhl M demonstrated that the targeting of TGF- β signaling pathway by SD-208 may prevent the development of melanoma bone metastases. However, these reports do not address exactly how SD-208 regulates miRNA expression and the genes which

are implicated in the cancer therapy.

Our study identified that miR18a to be modulated by SD-208. The miR18a overexpression seems to be an important factor in cancer treatment. Krutilina *et al.* reported that miR18a expression suppressed distant metastasis via the hypoxia-inducible factor 1-alpha pathway (29). Interestingly, it is well known that neglected or untreated retinoblastoma cases can demonstrate extraocular spread, primarily through the optic nerve (30,31). Based on these observations, it might be possible that the overexpression of miR18a could be implicated in repressing the metastatic behavior of intraocular retinoblastoma. In addition, miR-34 and miR-22 in our study following real time RT-PCR validation demonstrated a significant upregulation in treated Y-79 with SD-208 cells as compared to the controls. The miR-34 family has three members, miR-34 a,b,c. It has been thought that miR-34a is an important regulator of tumor suppression. This molecule controls different aspects of cell cycle processes such as differentiation and apoptosis; Its differential expression has been detected in different cancer such as breast, melanoma, pancreatic, and prostate cancer cell lines (32). However, the exact molecular mechanisms by which it acts as an anti-oncogenic activity is not well known. Various studies demonstrated that miR-34a is regulated by diverse approaches such as p53-dependent (33-35) and p53-independent mechanisms (36-38). TGF β signaling pathway as a molecular target for the miR-34a could also be another proposed model of function. More recently, a study suggested the capability of miR-34a to suppress epithelial-mesenchymal transition (EMT) in human cancer through TGF β signaling pathway (39). In addition, a cross talk between miR-34a, as a complex network interaction could interpret the miR-34a-mediated apoptosis in cancer. In this regard, a study using Y79 cells demonstrated that the expression of miR-34 was elevated after p53 activation (40). Our results, consistent with previous studies, recommend that miR-34a-triggered cell apoptosis in response to the SD-208 might open a new window for better understanding and application of the kinase inhibitor-based therapy.

Moreover, miR-20a was significantly downregulated in treated retinoblastoma cells with SD-208 compared to controls in this study. miR-20 belongs to the miR-17/92 cluster and acts as an important oncogene in several cancers such as retinoblastoma (41) and targets genes important in regulating cell proliferation and cancer development. Previous studies have reported that this miRNA is involved in cellular proliferation (42) cell

cycle (43) and invasion activity (41). As it mentioned, Y-79 cells do not functionally express TGF β RI, a key component for TGF β signal transduction. Altogether, these observations postulate another alternative system by which miR-20a exerts its normal and abnormal biological function.

The miR-17-92 cluster comprises six miRNAs: miR-17, miR-18a, miR-19a, miR-19b, miR-20a, and miR-92a (44). It has been well characterized that the miR-20a acts as an oncogene in various malignancies cells by targeting different cell signaling pathways (44-48). Evidence that the miRNA exhibits oncogenic effects in retinoblastoma came from distinct studies (49-53). Our results in agreement with these studies approved the miR-20a could potentially act as an oncogene in retinoblastoma.

The most obvious reason to account for the mechanism(s) by which SD-208 could regulate the miRNAs in retinoblastoma could be due to different scenarios. Studies have reported the absence of TGF β RI in some cancerous cells such as Y-79 (54) and SW48 cells (55). As mentioned before, SD-208 can block the biological effects of TGF β cascades by inhibiting of the kinase activity of TGF β RI. In the present study also, we analyzed the TGF β RI and TGF β RII mRNA expression in the cells treated with and without the SD-208. We could not detect their mRNA in all tested samples (data not shown). Considering this phenomenon; it would be useful to explain an appropriate relationship between SD-208 and miRNA regulation under this circumstance. Our previous results showed that SD-208 could regulate miRNA-135b in colorectal cancer (55). Taken together, our findings support the notion that SD-208 could potentially regulate the miRNA expression by indirectly affecting the TGF β signaling (also we called a non-canonical pathway).

Acknowledgment

This study financially was supported by Tehran University of Medical Sciences and Arak University of Medical Sciences.

References

- Gurney JG, Ross JA, Wall DA, Bleyer WA, Severson RK, Robison LL. Infant cancer in the U.S.: histology-specific incidence and trends, 1973 to 1992. *J Pediatr Hematol/Oncol* 1997;19:428-32.
- Hanahan D, Weinberg RA. The Hallmarks of Cancer. *Cell* 2000;100:57-70.
- Knudson AG, Jr. Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci U S A*. 1971;68:820-3.
- Rushlow DE, Mol BM, Kennett JY, Yee S, Pajovic S, Theriault BL, et al. Characterisation of retinoblastomas without RB1 mutations: genomic, gene expression, and clinical studies. *Lancet Oncol* 2013;14:327-34.
- Zhou Y, Zhang X, Klibanski A. Genetic and epigenetic mutations of tumor suppressive genes in sporadic pituitary adenoma. *Mol Cell Endocrinol* 2014;386:16-33.
- MacGrogan D, Bookstein R. Tumour suppressor genes in prostate cancer. *Seminars in cancer biology*. 1997;8:11-9.
- Akbari A, Mobini GR, Maghsoudi R, Akhtari J, Faghiloo E, Farahnejad Z. Modulation of transforming growth factor- β signaling transducers in colon adenocarcinoma cells induced by staphylococcal enterotoxin B. *Molr Med Rep* 2016;13:909-91.
- Sabado Alvarez C. Molecular biology of retinoblastoma. *Clin Transl Oncol* 2008;10:389-94.
- Felsher DW. Role of MYCN in retinoblastoma. *Lancet Oncol* 2013;14:270-1.
- Parisi T, Bronson RT, Lees JA. Inactivation of the retinoblastoma gene yields a mouse model of malignant colorectal cancer. *Oncogene* 2015;34:5890-9.
- Sahi H, Savola S, Sihto H, Koljonen V, Bohling T, Knuutila S. RB1 gene in Merkel cell carcinoma: hypermethylation in all tumors and concurrent heterozygous deletions in the polyomavirus-negative subgroup. *APMIS* 2014;122:1157-66.
- Tan HL, Sood A, Rahimi HA, Wang W, Gupta N, Hicks J, et al. Rb loss is characteristic of prostatic small cell neuroendocrine carcinoma. *Clin Cancer Res* 2014;20:890-903.
- Karimi A, Majidzadeh-A K, Madjd Z, Akbari A, Habibi L, Akrami SM. Effect of Copper Sulfate on Expression of Endogenous L1 Retrotransposons in HepG2 Cells (Hepatocellular Carcinoma). *Biol Trace Element Res* 2015;165:131-4.
- Toki H, Inoue M, Minowa O, Motegi H, Saiki Y, Wakana S, et al. Novel retinoblastoma mutation abrogating the interaction to E2F2/3, but not E2F1, led to selective suppression of thyroid tumors. *Cancer Sci* 2014;105:1360-8.
- Varghese AM, Zakowski MF, Yu HA, Won HH, Riely GJ, Krug LM, et al. Small-cell lung cancers in patients who never smoked cigarettes. *J Thorac Oncol* 2014;9:892-6.
- Deng L, Yang SB, Xu FF, Zhang JH. Long noncoding RNA CCAT1 promotes hepatocellular carcinoma progression by functioning as let-7 sponge. *J Exp Clin Cancer Res* 2015;34:18.

17. Hunten S, Kaller M, Drepper F, Oeljeklaus S, Bonfert T, Erhard F, et al. p53-regulated networks of protein, mRNA, miRNA and lncRNA expression revealed by integrated pSILAC and NGS analyses. *Mol Cell Proteomics* 2015;14:2609-29.
18. Sheikhzade M, Akbari A, Karkhaneh R, Ghassemi F, Sadeghi AR, Shahrzad Kavakani M, Mobini GR, et al. Digoxin Inhibits Retinoblastoma through Suppressing a Non-canonical TGF β Signaling Pathway. *BJMMR* 2016;15:1-10.
19. Jin K, Luo G, Xiao Z, Liu Z, Liu C, Ji S, et al. Noncoding RNAs as potential biomarkers to predict the outcome in pancreatic cancer. *Drug Des Devel Ther* 2015;9:1247-55.
20. Liang WC, Fu WM, Wong CW, Wang Y, Wang WM, Hu GX, et al. The lncRNA H19 promotes epithelial to mesenchymal transition by functioning as miRNA sponges in colorectal cancer. *Oncotarget* 2015;6:22513-25.
21. Liz J, Esteller M. lncRNAs and microRNAs with a role in cancer development. *Biochim Biophys Acta* 2016;1859:169-76.
22. Mirakholi M, Mahmoudi T, Heidari M. MicroRNAs horizon in retinoblastoma. *Acta Med Iran* 2013;51:823-9.
23. Zhou X, Ye F, Yin C, Zhuang Y, Yue G, Zhang G. The Interaction Between miR-141 and lncRNA-H19 in Regulating Cell Proliferation and Migration in Gastric Cancer. *Cell Physiol Biochem* 2015;36:1440-52.
24. Mirakholi M, Mahmoudi T, Heidari M. MicroRNAs horizon in retinoblastoma. *Acta Med Iran* 2013;51:823-9.
25. Akbari A, Amanpour S, Muhammadnejad S, Ghahremani MH, Ghaffari SH, Dehpour AR, et al. Evaluation of antitumor activity of a TGF-beta receptor I inhibitor (SD-208) on human colon adenocarcinoma. *Daru* 2014;22:47.
26. Lin SY, Chang HH, Lai YH, Lin CH, Chen MH, Chang GC, et al. Digoxin Suppresses Tumor Malignancy through Inhibiting Multiple Src-Related Signaling Pathways in Non-Small Cell Lung Cancer. *PloS One* 2015;10:e0123305.
27. Ge R, Rajeev V, Ray P, Lattime E, Rittling S, Medicherla S, et al. Inhibition of growth and metastasis of mouse mammary carcinoma by selective inhibitor of transforming growth factor-beta type I receptor kinase in vivo. *Clin Cancer Res* 2006;12:4315-30.
28. Mohammad KS, Javelaud D, Fournier PG, Niewolna M, McKenna CR, Peng XH, et al. TGF-beta-RI kinase inhibitor SD-208 reduces the development and progression of melanoma bone metastases. *Cancer Res* 2011;71:175-84.
29. Uhl M, Aulwurm S, Wischhusen J, Weiler M, Ma JY, Almirez R, et al. SD-208, a novel transforming growth factor beta receptor I kinase inhibitor, inhibits growth and invasiveness and enhances immunogenicity of murine and human glioma cells in vitro and in vivo. *Cancer Res* 2004;64:7954-61.
30. Yin JJ, Selander K, Chirgwin JM, Dallas M, Grubbs BG, Wieser R, et al. TGF-beta signaling blockade inhibits PTHrP secretion by breast cancer cells and bone metastases development. *J Clin Invest* 1999;103:197-206.
31. Gaspar NJ, Li L, Kapoun AM, Medicherla S, Reddy M, Li G, et al. Inhibition of transforming growth factor beta signaling reduces pancreatic adenocarcinoma growth and invasiveness. *Mol Pharmacol* 2007;72:152-61.
32. Medicherla S, Li L, Ma JY, Kapoun AM, Gaspar NJ, Liu YW, et al. Antitumor activity of TGF-beta inhibitor is dependent on the microenvironment. *Anticancer Res* 2007;27:4149-57.
33. Krutilina R, Sun W, Sethuraman A, Brown M, Seagroves TN, Pfeffer LM, et al. MicroRNA-18a inhibits hypoxia-inducible factor 1-alpha activity and lung metastasis in basal breast cancers. *Breast Cancer Res* 2014;16:R78.
34. Jacquemin C, Karcioğlu ZA. Detection of optic nerve involvement in retinoblastoma with enhanced computed tomography. *Eye* 1998;12:179-83.
35. Magrann I, Abramson DH, Ellsworth RM. Optic nerve involvement in retinoblastoma. *Ophthalmology* 1989;96:217-22.
36. Lodygin D, Tarasov V, Epanchintsev A, Berking C, Knyazeva T, Korner H, et al. Inactivation of miR-34a by aberrant CpG methylation in multiple types of cancer. *Cell Cycle* 2008;7:2591-600.
37. Kovacs B, Lumayag S, Cowan C, Xu S. MicroRNAs in early diabetic retinopathy in streptozotocin-induced diabetic rats. *Invest Ophthalmol Vis Sci* 2011;52:4402-9.
38. He L, He X, Lowe SW, Hannon GJ. microRNAs join the p53 network--another piece in the tumour-suppression puzzle. *Nature Rev Cancer* 2007;7:819-22.
39. Raver-Shapira N, Marciano E, Meiri E, Spector Y, Rosenfeld N, Moskovits N, et al. Transcriptional activation of miR-34a contributes to p53-mediated apoptosis. *Mol Cell* 2007;26:731-43.
40. Christoffersen NR, Shalgi R, Frankel LB, Leucci E, Lees M, Klausen M, et al. p53-independent upregulation of miR-34a during oncogene-induced senescence represses MYC. *Cell Death Differ* 2010;17:236-45.
41. Schirmer U, Doberstein K, Rupp AK, Bretz NP, Wuttig D, Kiefel H, et al. Role of miR-34a as a suppressor of L1CAM in endometrial carcinoma. *Oncotarget* 2014;5:462-72.
42. Genovese G, Ergun A, Shukla SA, Campos B, Hanna J, Ghosh P, et al. microRNA regulatory network inference identifies miR-34a as a novel regulator of TGF-beta signaling in glioblastoma. *Cancer Discov* 2012;2:736-49.

43. Qiao P, Li G, Bi W, Yang L, Yao L, Wu D. microRNA-34a inhibits epithelial mesenchymal transition in human cholangiocarcinoma by targeting Smad4 through transforming growth factor-beta/Smad pathway. *BMC Cancer* 2015;15:469.
44. Dalgard CL, Gonzalez M, deNiro JE, O'Brien JM. Differential microRNA-34a expression and tumor suppressor function in retinoblastoma cells. *Invest Ophthalmol Vis Sci* 2009;50:4542-51.
45. Fan X, Liu Y, Jiang J, Ma Z, Wu H, Liu T, et al. miR-20a promotes proliferation and invasion by targeting APP in human ovarian cancer cells. *Acta Biochim Biophys Sin* 2010;42:318-24.
46. O'Donnell KA, Wentzel EA, Zeller KI, Dang CV, Mendell JT. c-Myc-regulated microRNAs modulate E2F1 expression. *Nature* 2005;435:839-43.
47. Pickering MT, Stadler BM, Kowalik TF. miR-17 and miR-20a temper an E2F1-induced G1 checkpoint to regulate cell cycle progression. *Oncogene* 2009;28:140-5.
48. Reddycherla AV, Meinert I, Reinhold A, Reinhold D, Schraven B, Simeoni L. miR-20a Inhibits TCR-Mediated Signaling and Cytokine Production in Human Naïve CD4+ T Cells. *PLoS One* 2015;10:e0125311.
49. He FC, Meng WW, Qu YH, Zhou MX, He J, Lv P, et al. Expression of circulating microRNA-20a and let-7a in esophageal squamous cell carcinoma. *World J Gastroenterol* 2015;21:4660-5.
50. Li X, Zhang Z, Yu M, Li L, Du G, Xiao W, et al. Involvement of miR-20a in Promoting Gastric Cancer Progression by Targeting Early Growth Response 2 (EGR2). *Int J Mol Sci* 2013;14:16226-39.
51. Xiong Y, Zhang L, Kebebew E. MiR-20a Is Upregulated in Anaplastic Thyroid Cancer and Targets LIMK1. *PLoS One* 2014;9:e96103.
52. Zhao S, Yao D, Chen J, Ding N, Ren F. MiR-20a Promotes Cervical Cancer Proliferation and Metastasis in Vitro and In Vivo. *PLoS One* 2015;10:e0120905.
53. Beta M, Venkatesan N, Vasudevan M, Vetrivel U, Khetan V, Krishnakumar S. Identification and Insilico Analysis of Retinoblastoma Serum microRNA Profile and Gene Targets Towards Prediction of Novel Serum Biomarkers. *Bioinform Biol Insights* 2013;7:21-34.
54. Ebi H, Sato T, Sugito N, Hosono Y, Yatabe Y, Matsuyama Y, et al. Counterbalance between RB inactivation and miR-17-92 overexpression in reactive oxygen species and DNA damage induction in lung cancers. *Oncogene* 2009;28:3371-9.
55. Guo J, Miao Y, Xiao B, Huan R, Jiang Z, Meng D, et al. Differential expression of microRNA species in human gastric cancer versus non-tumorous tissues. *J Gastroenterol Hepatol* 2009;24:652-7.
56. Sylvestre Y, De Guire V, Querido E, Mukhopadhyay UK, Bourdeau V, Major F, et al. An E2F/miR-20a autoregulatory feedback loop. *J Biol Chem* 2007;282:2135-43.
57. Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, et al. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Natl Acad Sci U S A*. 2006;103:2257-61.
58. Kimchi A, Wang XF, Weinberg RA, Cheifetz S, Massague J. Absence of TGF-beta receptors and growth inhibitory responses in retinoblastoma cells. *Science (New York, NY)* 1988;240:196-9.
59. MacKay SL, Yaswen LR, Tarnuzzer RW, Moldawer LL, Bland KI, Copeland EM, 3rd, et al. Colon cancer cells that are not growth inhibited by TGF-beta lack functional type I and type II TGF-beta receptors. *Ann Surg* 1995;221:767-76.
60. Akbari A, Ghahremani MH, Mobini GR, et al. Down-regulation of miR-135b in colon adenocarcinoma induced by a TGF-β receptor I kinase inhibitor (SD-208). *Iran J Basic Med Sci* 2015;18:856-61.

SID



سرویس های
ویژه



سرویس ترجمه
تخصصی



کارگاه های
آموزشی



بلاگ
مرکز اطلاعات علمی




عضویت در
خبرنامه



فیلم های
آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی جهاد دانشگاهی



مباحث پیشرفته یادگیری عمیق؛
شبکه های توجه گرافی
(Graph Attention Networks)



کارگاه آنلاین آموزش استفاده از
وب آوساینس



کارگاه آنلاین مقاله روزمره انگلیسی