

Original Article

Effect of Psychological Distress on Weight Concern and Weight Control Behaviors

Hamidreza Roohafza MD¹, Ali Kabir MD MPH PhD Candidate^{2,3}, Masoumeh Sadeghi MD⁴, Pedram Shokouh MD⁵, Seyed Hossein Aalaei-Andabili MD⁶, Yadollah Mehrabi PhD⁷, Nizal Sarrafzadegan MD⁸

Abstract

Background: Obesity is associated with chronic disorders like coronary artery diseases, metabolic syndrome, cancers, and psychiatric disorders. Stress may contribute to weight gain by disrupting weight concern, and lead to uncontrolled eating behavior. This study aimed to investigate the effects of stress on weight concern and control behaviors in normal weight and obese adults.

Methods: A total of 9544 subjects were selected by multi-stage random sampling from three provinces in central Iran. Information related to weight concern and control behavior was registered in normal weight and obese participants. Psychological distress was measured by a 12-item General Health Questionnaire (GHQ-12) and subjects were divided into high and low stress groups. Logistic regression was used for analysis.

Results: The mean age of participants was 38.7 ± 15.5 years and 50% (4772) of them were males. The adjusted odds ratio (OR) for age, sex and education of high stress to low stress level for weight concern, weight control behavior and acceptable physical activity behavior was more than 1; but the OR was less than 1 for waist circumference, obesity and healthy diet behavior. Among obese participants, higher levels of stress were associated with lower weight concern with OR, 95%CI: 0.821, (0.682 – 0.988), lower acceptable physical activity with OR = 0.833, 95%CI: (0.624 – 0.912), but higher rates of healthy diet behavior with OR = 1.360, 95% CI: (1.040 – 1.780).

Conclusion: Individuals with high stress level have lower weight concern and lower physical activity; therefore, they are prone to weight gain and obesity. It could be concluded that stress management should be considered as a crucial component of obesity prevention and control programs.

Keywords: Iran, obesity, psychological distress, weight concern, weight control behavior

Cite this article as: Roohafza H, Kabir A, Sadeghi M, Shokouh P, Aalaei-Andabili SH, Mehrabi Y, et al. Psychological distress effect on weight concern and weight control behaviors. *Arch Iran Med.* 2014; **17**(9): 608 – 612.

Introduction

Obesity is a rising problem in modern societies; it leads to an increase in the prevalence of chronic disorders such as type 2 diabetes mellitus, cardiovascular diseases, metabolic syndrome, some types of cancers, and psychiatric disorders.¹ Chronic stress has attracted attention as a promoting factor for the development of obesity. Epidemiological studies have reported an association between stress, obesity and its consequences. Stress

can influence human eating behaviors by increasing the appetite for highly palatable and high energy foods. Moreover, when exposed to a stressor, people prefer to eat or overeat some foods that they were avoiding before.² It seems that eating in the absence of hunger is affected by stress among adults.³ In support of this notion, high depression and anxiety levels have been reported among both overweight males and females.⁴ On the other hand, one study did not find any significant difference in food intake between high stress and low stress females.⁵ Moreover, female gender and older age have been found as effective factors for stress level.⁶ Also, it has been suggested that obesity is an important predictor of psychological distress.⁷

Physical activity and exercise are known as weight control behaviors. A review suggested that physical activity and exercise could contribute to favorable adaptation with stress and attenuate its negative effects.⁸ It has been reported that stress significantly impairs physical activity,⁹ and encourages overeating and oversleeping.¹⁰

Some studies have already reported the prevalence of obesity and its association with stress in Iran,^{11,12} but none were designed to investigate the factors affecting stress level or the effects of stress on weight concern and weight control behaviors. In order to fill this gap, we performed this study with main goals of determining the effective factors on stress level, and association between stress levels and weight concern and control behaviors in normal and obese individuals of a representative sample of the central parts of Iran.

Authors' affiliations: ¹Mental Health Department, Isfahan Cardiovascular Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran, ²Minimally Invasive Surgery Research Center; Iran University of Medical Sciences, Tehran, Iran, ³Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran, ⁴Cardiac Rehabilitation Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran, ⁵Hypertension Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran, ⁶Nikan Health Researchers Institute, Tehran, Iran, ⁷Department of Epidemiology, School of Public Health, Shahid Beheshti University of Medical Sciences, Tehran, Iran, ⁸Isfahan Cardiovascular Research Center, Isfahan Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran.

Corresponding author and reprints: Masoumeh Sadeghi MD, Cardiac Rehabilitation Research Center, Isfahan Cardiovascular Research Institute, WHO Collaborating Center for Research and Training in Cardiovascular Diseases Control Isfahan University of Medical Sciences, Isfahan, Iran. P. O. Box: 81465-1148, Tel: +98-311-3359797, Fax: +98-311-3373435, E-mail: m_sadeghi@crc.mui.ac.ir, aikabir@yahoo.com.

Accepted for publication: 20 June 2014

Materials and Methods

Data were obtained from part of the “Isfahan Healthy Heart Program” (IHHP), which is a community-based program designed to prevent and control cardiovascular disease and promote healthy lifestyle behaviors. The IHHP aims and design are described in detail elsewhere.^{13,14} The protocol of the study was approved by the ethics committee of the Isfahan University of Medical Sciences.

Study population

A total of 9553 men and women aged ≥ 19 years from three cities in central parts of Iran (Isfahan, Najafabad and Arak) were selected according to the latest national census in 2007. In each urban or rural community, a sample of adults was selected through multi-stage random sampling. Clusters were defined according to the local health services divisions. Then, they were stratified based on their age and sex. To achieve adequate sample size, those who declined to participate in the study were replaced by their neighbors. Pregnant, mentally retarded or physically disabled individuals were not eligible for enrolment. According to the 2006 National Census, the population of Isfahan and Najafabad (a cities neighboring Isfahan) was 1,986,542 and 282,430, respectively. Arak, with a population of 615,702 is located in northwest of Isfahan.^{14,15}

Demographic and anthropometric measurements

After obtaining written informed consent, data on demographic, anthropometric and socioeconomic characteristics and lifestyle behaviors were collected by standardized interviewer-administered questionnaires.

We categorized the participants' educational level as 0 – 5 years (primary school), 6 – 12 years (high school) and > 12 years (university degree).

Psychological distress

Psychological distress was measured by a 12-item General Health Questionnaire (GHQ-12), a self-administered screening tool for assessing psychological distress.¹⁶ GHQ-12 is a consistent and reliable instrument for use in general population studies.¹⁷ Each item is rated on a four-point scale (less than usual, no more than usual, fairly more than usual and much more than usual). The system used to score the GHQ-12 questionnaires was the 0-0-1-1 method. Using this method, a participant could have been scored between 0 and 12 points; a score of 4 or more was used to identify a participant with high stress level.

Term definitions

In our study, weight concern was defined as the individual's perceived weight and body dissatisfaction and was determined by the following dichotomous question: “Are you satisfied with your current body shape and size?” Everyone who gave negative response to the first question was considered weight-concerned and was asked the following: “Are you doing any type of activities to reduce your weight?” The participants' positive responses to the latter question were reported under the title of “weight control behavior”. Individuals with weight concern underwent a complete enquiry about the type and quality of their weight-reducing activities. Healthy behavior was defined as acceptable physical activity and healthy diet (low fat and low calorie foods) which were in compliance with the advice of professionals.^{13,14} In this study, obesity was defined based on a waist circumference ≥ 88 cm in

females and ≥ 102 cm in males.¹⁸

Statistical analysis

The Statistical Package for Social Sciences version 15.0 (SPSS Inc., Chicago, IL, USA) was used for all statistical analyses. Quantitative variables were expressed as mean \pm standard deviation. To compare continuous variables between groups, Student's t-test was used. Categorical variables were compared between groups using Chi-square and Wilcoxon tests. The enter method of logistic regression analysis was used for determining the effect of obesity, weight concern, weight control behavior, healthy diet behavior, and acceptable physical activity behavior as independent variables on the stress level of all subjects and obese individuals separately. Age, gender and education level were considered for adjustment. A *P*-value of less than 0.05 was considered statistically significant in all analyses.

Results

A total of 9544 individuals (mean age: 38.7 ± 15.5 years; 50% male) were enrolled in the study. The participants' characteristics are shown in Table 1. Subjects with higher stress levels were significantly older and more likely to be female or obese. They had healthier diet behavior but had lower education level, less concern about their weight, weight control behavior and acceptable physical activity (Table 1).

The results of analyses in the obese subset are summarized in Table 2. According to the table, subjects with higher levels of stress were older, more likely to be female and had greater waist circumference and healthy diet behavior while their education level, weight concern, and acceptable physical activity was significantly lower (Table 2).

In obese participants, crude and adjusted ORs of high stress to low stress level was higher than 1 for weight concern and acceptable physical activity behavior, but lower than 1 for waist circumference and healthy diet behavior. Weight control behavior was not different between the two groups on multivariate analysis (Table 3). Even after adjustment for education, in addition to age and sex, statistical significant level, the ORs and their 95% CIs did not change significantly.

Discussion

We found that high stress people were more obese. In line with this finding, some other studies have shown the association of higher levels of stress with obesity.¹⁹ Other factors may influence stress effects in weight gain and obesity but in our study, as well as another one,²⁰ high level of psychological distress remained significantly associated with obesity after adjusting for possible confounders; however, it seems that there is a reciprocal association between obesity and stress level, and at least one study has found obesity to be a predisposing factor to experiencing high levels of psychological distress.⁷ In contrast, one study has reported lower levels of stress among young obese Bangladeshi people compared to normal-weight subjects.²¹ This controversy might indicate that cultural differences may play a role in the correlation between psychological distress and obesity.

We found that people with high levels of stress have lower concern about their weight gain, less weight control behavior, and less acceptable physical activity which are important factors in

Table 1. Basic characteristics of participants in study of psychological distress effect on weight concern and control behaviors^a

	Low stress (n = 6289)	High stress (n = 3264)	P
Age (years)	38.4 ± 15.1	39.5 ± 16.2	0.001
Sex			< 0.001
Male	3402 (54.1%)	1375 (42.1%)	
Female	2887 (45.9%)	1880 (57.9%)	
Education level			< 0.001
5 year(s)–0	2677 (42.6%)	1619 (49.7%)	
12 years–6	2699 (43.0%)	1278 (39.2%)	
> 12 years	901 (14.4%)	363 (11.1%)	
Obesity ^b	1795 (28.6%)	1020 (31.2%)	< 0.001
Weight concern	4580/6289 (72.8%)	2137/3264 (65.5%)	< 0.001
Weight control behavior	1956/4580 (42.7%)	880/2137 (41.2%)	0.048
Healthy diet behavior	666/1956 (34.0%)	398/880 (45.2%)	< 0.001
Acceptable physical activity behavior	973/1956 (49.7%)	397/880 (45.1%)	< 0.001

^aData are expressed as mean ± SD for continuous variables and number (percentage) of participants for categorical variables; ^bobesity as: waist circumference ≥ 88 cm for females and ≥ 102 cm for males

Table 2. Characteristics of obese individuals in study of psychological distress effect on weight concern and control behaviors^a

	low stress (n = 1795)	high stress (n = 1020)	P
Age (years)	43.7 ± 14.9	45.8 ± 15.3	< 0.001
Sex			< 0.001
Female	1425 (79.4%)	871 (85.4%)	
Male	370 (20.6%)	149 (14.6%)	
Education level			< 0.001
5 years–0	1077 (60.2%)	689 (67.6%)	
12 years–6	589 (32.9%)	276 (27.1%)	
> 12 years	123 (6.9%)	54 (5.3%)	
Waist circumference (cm)	101.3 ± 8.4	102.2 ± 9.1	< 0.001
Weight concern	1276/1795 (71.1%)	640/1020 (62.7%)	< 0.001
Weight control behavior	616/1276 (48.3%)	320/640 (50.0%)	0.786
Healthy diet behavior	243/616 (40.1%)	171/320 (51.8%)	0.003
Acceptable physical activity behavior	220/616 (36.3%)	102/320 (30.9%)	0.049

^aData are expressed as mean ± SD for continuous variables and number (percentage) of participants for categorical variables

Table 3. The association of weight related factors with high level stress in obese individuals

	Unadjusted	Adjusted ^a
Waist circumference (cm)	1.015 (1.004, 1.580)	1.045 (1.025, 1.956)
Weight concern	0.718 (0.604, 0.854)	0.821 (0.682, 0.988)
Weight control behavior	0.977 (0.827, 1.154)	1.227 (0.996, 2.042)
Healthy diet behavior	1.492 (1.147, 1.940)	1.360 (1.040, 1.780)
Acceptable physical activity behavior	0.757 (0.571, 0.904)	0.833 (0.624, 0.912)

^a Adjusted for age, sex and education

controlling weight. Behavioral weight control intervention with stress management components increases concern about weight and weight control behaviors in overweight/obese women.²² One study found that 80% of adolescent girls with high level disturbance such as psychosocial problems had no real concern about their weight.²³ On the other hand, a study reported that difference in weight concern can account for difference in psychological stress experience.²⁴ In addition, job strain was positively associated with average daily intake of fat in a Japanese population.²⁵ Prolonged stress was associated with lower leisure time physical activity in both men and women.^{26,27} Subjects with significantly lower levels of stress had two to three times more physical activ-

ity.²⁸ However, even among the limited number of available studies, some degree of inconsistency can be seen, as another study did not find any role for health behaviors such as physical activity in the association of stress and weight gain.²⁹

It has been shown that high levels of chronic stress make short-term endocrine changes such as Hypothalamic–pituitary–adrenal (HPA) axis regulation disruption and increased levels of glucocorticoids,³⁰ which may lead to prolonged physiologic consequences even after termination of the stressor. Positive association of obesity and plasma cortisol level is now well-documented.³¹ Cortisol level is positively associated with response of right amygdala in brain. Acute stress can potentiate right amygdala response to food

and stress-related eating which leads to changes favoring high caloric and high-fat diets.³² Excess eating has been reported among stressed subjects and a study has suggested the inhibition of satiety by stress.³³ The association of uncontrolled eating under stress with obesity has been reported by Heatherton, et al. only in females;³⁴ but stressed men ate significantly lower unhealthy foods than non-stressed subjects in another study.³⁵ It is noteworthy that these findings were challenged by some contradictory studies. One study failed to find any significant difference in food intake between high stress and low stress females.⁵ In addition, Shapiro, et al.³⁶ rejected more calorie intake and hunger level among people with higher levels of stress. Although our results show that subjects with higher stress have healthy diet behavior, it has been reported that individuals consuming healthy foods, such as fish, experience half the psychological distress experienced by subjects using high calorie foods.³⁷

According to our findings, stress level was higher in women. Girls had higher mean of stress score in their life compared to boys.²⁸ It seems that females experience higher levels of stress and they show more negative psychological symptoms than males.¹² Previous observations have shown that women are more prone to overeating in high stress situations.³⁸ Therefore, females need more attention when stressed and lack of emotional support is the most important factor predicting stress-related eating in this group.³⁹ These results indicate a gender specific response to stress. We found that higher level of stress is associated with lower level of education.

Overweight and obese people are at higher risk of having lower levels of education and higher rates of unemployment.⁴⁰ Level of stress-related cortisol was positively associated with lower education level in a report by Rosmond, et al.⁴¹ We found that education has an important role in weight concern, because patients with high level of stress were obese and also less educated. Women with lower level of education might have more traditional habits and views that link beauty and health with being slightly overweight.³⁹ However, another study could not find any significant association between education and stress level.⁴²

We found a direct positive correlation between stress level and age. Disturbance of HPA axis is more common among elderly males and females than younger population.⁴³ Therefore, this abnormality causes a mal-adaptation of old people with stress which may make them obese.⁴⁴ Furthermore, baseline cortisol concentration increases with age in women.⁴⁵ In contrast, it is suggested that psychological distress decreases with age, especially in women.⁴⁶

Although the present study benefited from a large and representative sample, the following limitations should be considered when generalizing the results. First, the people of central parts of Iran were our target population whose habits and culture may differ from other parts of Iran. Therefore, this study may not be applicable to all Iranians. Second is the probability of recall bias which may influence the subjects' answers about weight concern, food intake, and physical activity. However, the impact of this bias was estimated negligible. Finally, the design of this study was cross-sectional; hence, associations should be interpreted cautiously.

Our findings showed that people with high levels of stress had less weight concern and physical activity compared with low-stress subjects. It seems that high level of stress distracts people's attention from their body shape and prevents them from doing acceptable physical activity. Accordingly, stress management strategies may contribute to weight control or preventing overweight obesity.

Acknowledgment

We would like to express our thanks to the Iranian Budget and Programming Organization, the Isfahan Cardiovascular Research Institute and Provincial Health Office of the Isfahan University of Medical Sciences that supported the project. The registration number of this project in this Institute was 80117.

Conflict of Interest

There is no conflict of interest.

References

- Lobstein T, Baur L, Uauy R. Obesity in children and young people: a crisis in public health. *Obes Rev.* 2004; **5(Suppl 1)**: 4 – 104. (Epub 2004/04/21).
- Torres SJ, Nowson CA. Relationship between stress, eating behavior, and obesity. *Nutrition.* 2007; **23(11–12)**: 887 – 894. (Epub 2007/09/18).
- Rutters F, Lemmens SG, Born JM, Bouwman F, Nieuwenhuizen AG, Mariman E, et al. Genetic associations with acute stress-related changes in eating in the absence of hunger. *Patient Educ Couns.* 2010; **79(3)**: 367 – 371. (Epub 2010/04/23).
- Stunkard AJ, Faith MS, Allison KC. Depression and obesity. *Biol Psychiatry.* 2003; **54(3)**: 330 – 337. (Epub 2003/08/02).
- Ball K, Lee C, Brown W. Psychological stress and disordered eating: an exploratory study with young Australian women. *Women Health.* 1999; **29(2)**: 1 – 15. (Epub 1999/07/31).
- Chan CW, Choi KC, Chien WT, Cheng KK, Goggins W, So WK, et al. Health-related quality-of-life and psychological distress of young adult survivors of childhood cancer in Hong Kong. *Psycho-oncology.* 2013; **23(2)**: 229 – 236. DOI: 10.1002/pon.3396. (Epub 2013/09/13).
- Walders-Abramson N, Nadeau KJ, Kelsey MM, Schmiege SJ, Ellert S, Cejka A, et al. Psychological functioning in adolescents with obesity co-morbidities. *Child Obes.* 2013; **9(4)**: 319 – 325. (Epub 2013/06/15).
- Holmes ME, Ekkekakis P, Eisenmann JC. The physical activity, stress and metabolic syndrome triangle: a guide to unfamiliar territory for the obesity researcher. *Obes Rev.* 2010; **11(7)**: 492 – 507. (Epub 2009/11/10).
- Stults-Kolehmainen MA, Sinha R. The Effects of Stress on Physical Activity and Exercise. *Sports Med.* 2013; **44(1)**: 81 – 121. DOI: 10.1007/s40279-013-0090-5 (Epub 2013/09/14).
- Parker G, Roy K, Mitchell P, Wilhelm K, Malhi G, Hadzi-Pavlovic D. Atypical depression: a reappraisal. *Am J Psychiatry.* 2002; **159(9)**: 1470 – 1479. (Epub 2002/08/31).
- Kelishadi R, Alikhani S, Delavari A, Alaedini F, Safaie A, Hojatzadeh E. Obesity and associated lifestyle behaviours in Iran: findings from the First National Non-communicable Disease Risk Factor Surveillance Survey. *Public Health Nutr.* 2008; **11(3)**: 246 – 251. (Epub 2007/07/13).
- Janghorbani M, Amini M, Willett WC, Mehdi Gouya M, Delavari A, Alikhani S, et al. First nationwide survey of prevalence of overweight, underweight, and abdominal obesity in Iranian adults. *Obesity (Silver Spring).* 2007; **15(11)**: 2797 – 2808. (Epub 2007/12/12).
- Sarrafzadegan N, Baghaei AM, Kelishadi R, Rabiei K, Sadri GH, Talei M. First annual evaluation of Isfahan Healthy Heart Program (IHHP): full report. 2004; Available from: URL: <http://74.125.39.104/search?q=cache:N9zP2gD3WWEJ:ihhp.mui.ac.ir> (Accessed Date: 24 October 2008).
- Sarrafzadegan N, Baghaei A, Sadri Gh, Kelishadi R, Malekafzali H, Boshtam M, et al. Isfahan healthy heart program: Evaluation of comprehensive, community-based interventions for non-communicable disease prevention. *Prev Control.* 2006; **2(2)**: 73 – 84.
- Sarraf-Zadegan N, Sadri G, Malek Afzali H, Baghaei M, Mohammadi Fard N, Shahrokhi S, et al. Isfahan Healthy Heart Programme: a comprehensive integrated community-based programme for cardiovascular disease prevention and control. Design, methods and initial experience. *Acta Cardiol.* 2003; **58(4)**: 309 – 320.
- Goldberg DP, Hillier VF. A scaled version of the General Health Questionnaire. *Psychol Med.* 1979; **9(1)**: 139 – 145.

17. Montazeri A, Harirchi AM, Shariati M, Garmaroudi G, Ebadi M, Fateh A. The 12-item General Health Questionnaire (GHQ-12): translation and validation study of the Iranian version. *Health Qual Life Outcomes*. 2003; **1**: 66.
18. National Cholesterol Education Program (NCEP) Expert Panel on Detection E, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation*. 2002; **106**: 3143 – 3421.
19. Rutters F, Nieuwenhuizen AG, Lemmens SG, Born JM, Westerterp-Plantenga MS. Acute stress-related changes in eating in the absence of hunger. *Obesity (Silver Spring)*. 2009; **17**(1): 72 – 77. (Epub 2008/11/11).
20. Ostry AS, Radi S, Louie AM, LaMontagne AD. Psychosocial and other working conditions in relation to body mass index in a representative sample of Australian workers. *BMC Public Health*. 2006; **6**: 53. (Epub 2006/03/04).
21. Viner RM, Haines MM, Taylor SJ, Head J, Booy R, Stansfeld S. Body mass, weight control behaviours, weight perception and emotional well being in a multiethnic sample of early adolescents. *Int J Obes (Lond)*. 2006; **30**(10): 1514 – 1521. (Epub 2006/05/24).
22. Cox TL, Krukowski R, Love SJ, Eddings K, Dicarolo M, Chang JY, et al. Stress Management-Augmented Behavioral Weight Loss Intervention for African American Women: A Pilot, Randomized Controlled Trial. *Health Educ Behav*. 2012; **40**(1): 78 – 87. DOI: 10.1177/1090198112439411. (Epub 2012/04/17).
23. Cooper PJ, Goodyer I. Prevalence and significance of weight and shape concerns in girls aged 11-16 years. *Br J Psychiatry*. 1997; **171**: 542 – 544. (Epub 1998/03/31).
24. Allen KL, Byrne SM, Blair EM, Davis EA. Why do some overweight children experience psychological problems? The role of weight and shape concern. *International Journal of Pediatric Obesity : IJPO : An Official Journal of the International Association for the Study of Obesity*. 2006; **1**(4): 239 – 247. (Epub 2007/10/02).
25. Kawakami N, Tsutsumi A, Haratani T, Kobayashi F, Ishizaki M, Hayashi T, et al. Job strain, worksite support, and nutrient intake among employed Japanese men and women. *J Epidemiol*. 2006; **16**(2): 79 – 89. (Epub 2006/03/16).
26. Kouvonen A, Kivimaki M, Elovainio M, Virtanen M, Linna A, Vahtera J. Job strain and leisure-time physical activity in female and male public sector employees. *Prev Med*. 2005; **41**(2): 532 – 539. (Epub 2005/05/27).
27. Lallukka T, Lahelma E, Rahkonen O, Roos E, Laaksonen E, Martikainen P, et al. Associations of job strain and working overtime with adverse health behaviors and obesity: evidence from the Whitehall II Study, Helsinki Health Study, and the Japanese Civil Servants Study. *Soc Sci Med*. 2008; **66**(8): 1681 – 1698. (Epub 2008/02/12).
28. Moljord IE, Moksnes UK, Eriksen L, Espnes GA. Stress and happiness among adolescents with varying frequency of physical activity. *Percept Mot Skills*. 2011; **113**(2): 631 – 646. (Epub 2011/12/22).
29. Serlachius A, Hamer M, Wardle J. Stress and weight change in university students in the United Kingdom. *Physiol Behav*. 2007; **92**(4): 548 – 553. (Epub 2007/06/01).
30. Tamashiro KL, Hegeman MA, Sakai RR. Chronic social stress in a changing dietary environment. *Physiol Behav*. 2006; **89**(4): 536 – 542. (Epub 2006/07/18).
31. Kalra SP, Kalra PS. NPY and cohorts in regulating appetite, obesity and metabolic syndrome: beneficial effects of gene therapy. *Neuropeptides*. 2004; **38**(4): 201 – 211. (Epub 2004/09/01).
32. Rudenga KJ, Sinha R, Small DM. Acute stress potentiates brain response to milkshake as a function of body weight and chronic stress. *Int J Obes (Lond)*. 2012. (Epub 2012/03/21).
33. Wallis DJ, Hetherington MM. Emotions and eating. Self-reported and experimentally induced changes in food intake under stress. *Appetite*. 2009; **52**(2): 355 – 362. (Epub 2008/12/17).
34. Heatherton TF, Herman CP, Polivy J. Effects of physical threat and ego threat on eating behavior. *J Pers Soc Psychol*. 1991; **60**(1): 138 – 143. (Epub 1991/01/01).
35. Zellner DA, Saito S, Gonzalez J. The effect of stress on men's food selection. *Appetite*. 2007; **49**(3): 696 – 699. (Epub 2007/09/11).
36. Shapiro JR, Anderson DA. Counterregulatory eating behavior in multiple item test meals. *Eat Behav*. 2005; **6**(2): 169 – 178. (Epub 2004/12/16).
37. Bhattacharyya M, Marston L, Walters K, D'Costa G, King M, Nazareth I. Psychological distress, gender and dietary factors in South Asians: a cross-sectional survey. *Public Health Nutr*. 2013; **2013**: 1 – 9. (Epub 2013/06/19).
38. Zellner DA, Loaiza S, Gonzalez Z, Pita J, Morales J, Pecora D, et al. Food selection changes under stress. *Physiol Behav*. 2006; **87**(4): 789 – 793. (Epub 2006/03/08).
39. Rasheed P. Perception of body weight and self-reported eating and exercise behaviour among obese and non-obese women in Saudi Arabia. *Public Health*. 1998; **112**(6): 409 – 414. (Epub 1999/01/12).
40. Ali SM, Lindstrom M. Socioeconomic, psychosocial, behavioural, and psychological determinants of BMI among young women: differing patterns for underweight and overweight/obesity. *Eur J Public Health*. 2006; **16**(3): 325 – 331. (Epub 2005/09/16).
41. Rosmond R. Role of stress in the pathogenesis of the metabolic syndrome. *Psychoneuroendocrinology*. 2005; **30**(1): 1 – 10. (Epub 2004/09/11).
42. Ignatyev Y, Assimov M, Dochshanov D, Strohle A, Heinz A, Mundt AP. Social Characteristics of Psychological Distress in a Disadvantaged Urban Area of Kazakhstan. *Community Mental Health Journal*. 2013. (Epub 2013/05/29).
43. Deuschle M, Schweiger U, Weber B, Gotthardt U, Korner A, Schmitter J, et al. Diurnal activity and pulsatility of the hypothalamus-pituitary-adrenal system in male depressed patients and healthy controls. *J Clin Endocrinol Metab*. 1997; **82**(1): 234 – 238. (Epub 1997/01/0).
44. Hajian-Tilaki KO, Heidari B. Prevalence of obesity, central obesity and the associated factors in urban population aged 20-70 years, in the north of Iran: a population-based study and regression approach. *Obes Rev*. 2007; **8**(1): 3 – 10. (Epub 2007/01/11).
45. Martens EA, Lemmens SG, Adam TC, Westerterp-Plantenga MS. Sex differences in HPA axis activity in response to a meal. *Physiol Behav*. 2012; **106**(2): 272 – 277. (Epub 2012/03/01).
46. Brandheim S, Rantakeisu U, Starrin B. BMI and psychological distress in 68,000 Swedish adults: a weak association when controlling for an age-gender combination. *BMC Public Health*. 2013; **13**: 68. (Epub 2013/01/26).

Surf and download all data from SID.ir: www.SID.ir

Translate via STRS.ir: www.STRS.ir

Follow our scientific posts via our Blog: www.sid.ir/blog

Use our educational service (Courses, Workshops, Videos and etc.) via Workshop: www.sid.ir/workshop