

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی



مقاله نویسی علوم انسانی



اصول تنظیم قراردادها



آموزش مهارت های کاربردی در تدوین و چاپ مقاله

Indigestion Leads to Depression: Avicenna's Point of View

Malihe Motavasselian^{1,*}; Mandana Tavakkoli Kakhki²; Mohammad Reza Shams Ardekani¹

¹Department of Traditional Medicine, Faculty of Traditional Medicine, Tehran University of Medical Sciences, Tehran, IR Iran

²Department of Traditional Medicine, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

*Corresponding Author: Malihe Motavasselian, Department of Traditional Medicine, Faculty of Traditional Medicine, Tehran University of Medical Sciences, Tehran, IR Iran. Tel/Fax: +98-5118405210, E-mail: m_motavasselian@yahoo.com

Received: November 3, 2013; Accepted: November 6, 2013

Keywords: Depression; Dyspepsia; Medicine Traditional

Dear Editor,

Depression is a common mental disorder and leading cause of disability worldwide (1). As a definition, depression may be a mood state, a syndrome or a specific disorder that may be secondary to various medical illnesses (2, 3). Avicenna (980-1037 AD), a scholar in traditional medicine, believed that indigestion can lead to depression (4). Indigestion is a common gastrointestinal problem that consists of nausea, vomiting, heartburn, regurgitation and dyspepsia (5). Functional dyspepsia is one of the most common causes of indigestion (6). In the modern medicine the effect of psychological factors on indigestion has been confirmed (2). In the current study we have demonstrated evidences of the effects of indigestion on depression. A recent prospective study in 2012 has stated that people with Functional gastrointestinal disorders (FGIDs) such as functional dyspepsia who had not suffered from anxiety and depression at baseline significantly developed signs and symptoms of anxiety and depression over 12-year follow-up (7). The probable mechanisms for the relationship between depression and dyspepsia have also been studied. Functional brain imaging study has shown neurobiological link between psychological abnormalities and visceral hypersensitivity in FGIDs (8). Another study has approved the role of serotonin and noradrenalin, the two main neurotransmitters involved in the pathophysiology of depression, in indigestion (9). On the other hand, from the Avicenna's point of view the absorbable part of food converts to "chylous" in the stomach that absorbs via the mesenteric vessels to the liver. Then chylous is converted to "chymous" in the liver. Chymous is composed of four humors including "blood", "phlegm", "yellow bile" and "black bile". The

body is also composed of these four humors. Humors enter to the blood circulation and supply food and energy for different organs. Abnormal humors cause functional disturbances in all organs including brain. Therefore the development of depression may be explained by the presence of the abnormal humors in the blood circulation. Production of normal humors requires appropriate composition of the gastric chylous that is delivered to the liver. Some etiological factors affect composition of the gastric chylous including: type of food consumed, the quality of gastric digestion, and also the quality of post-feeding conditions. Hence, indigestion can lead to depression with the following mechanism (4, 10)

The preventive methods have a crucial importance in Iranian Traditional Medicine (ITM). To this end, some items have been considered in association with the three mentioned etiological factors for indigestion as follows:

Dietary habits that cause indigestion

Over eating, low eating, inappropriate time of eating like eating when stomach is full, eating slow-digesting foods before fast-digesting ones and also eating some foods like milk, watermelon, peach and mushroom

Gastric Disorders that cause indigestion

Primary gastric disorders like gastric impairments or secondary gastric disorders like ones due to accumulation of excessive gas in stomach or entering of post nasal discharge to it.

Post-feeding conditions that cause indigestion

Lack of sleep, excessive movement, intercourse, taking a shower, exposure to hot or cold weather, and also little or too much drinking after meal (4). If the Avicenna's perspective would be considered alongside with further complementary investigations especially through observational studies, the prevention of indigestion may lead to significance decrease in the prevalence of depression.

Acknowledgements

This study is a part of Ph.D thesis entitled: "Etiology and clinical manifestation of depression and anxiety based on Iranian Traditional Medicine" (Department of Traditional Medicine, Faculty of Traditional Medicines, Tehran University of Medical Sciences, Tehran, IR Iran) and was supported by a grant from Tehran University of Medical Sciences.

Funding Support

This study was supported by a grant from Tehran University of Medical Sciences.

Financial Disclosure

There is no financial disclosure.

Authors' Contributions

M. Motavasselian developed the original idea and the protocol, abstracted data, wrote the manuscript and is guarantor. M. Tavakkoli-Kakhki contributed to the development of the protocol and prepared the manuscript. M. Shams-Ardekani studied the manuscript for important intellectual content.

References

1. depression. 2013. Available from: <http://www.who.int/mediacentre/factsheets/fs369/en/>.
2. Sadock BJ, Kaplan HI, Sadock VA. *Kaplan & Sadock's synopsis of psychiatry: behavioral sciences/clinical psychiatry*. Philadelphia: Wolter Kluwer/Lippincott Williams & Wilkins; 2007.
3. Gelder MG, Mayou R, Geddes J. *Psychiatry*. London, UK: Oxford University Press Incorporated; 2005.
4. Ibn-e Sina A. *Al-Qanun fi al-Tibb (The Canon of Medicine)*. Lebanon: Alamy le-al-Matbooa institute; 2005.
5. William LH. Nausea, Vomiting and Indigestion. In: Kasper DL, Fauci AS, Hauser SL, Longo DL, Jameson JL, Loscalzo J editors. *Harrison's principle of internal medicine*. New York: McGraw-Hill; 2008. p. 240-5.
6. William LH, Chung O. Approach to the Patient with Gastrointestinal Disease. In: Kasper DL, Fauci AS, Hauser SL, Longo DL, Jameson JL, Loscalzo J editors. *Harrison's principle of internal medicine*. New York: McGraw-Hill; 2008. p. 1831-2.
7. Koloski NA, Jones M, Kalantar J, Weltman M, Zaguirre J, Talley NJ. The brain-gut pathway in functional gastrointestinal disorders is bidirectional: a 12-year prospective population-based study. *Gut*. 2012;**61**(9):1284-90.
8. Van Oudenhove L, Coen SJ, Aziz Q. Functional brain imaging of gastrointestinal sensation in health and disease. *World J Gastroenterol*. 2007;**13**(25):3438-45.
9. O'Mahony S, Dinan TG, Keeling PW, Chua AS. Central serotonergic and noradrenergic receptors in functional dyspepsia. *World J Gastroenterol*. 2006;**12**(17):2681-7.
10. Emtiazy M, Keshavarz M, Khodadoost M, Kamalinejad M, Gooshahgir SA, Shahradsaj H, et al. Relation between Body Humors and Hypercholesterolemia: An Iranian Traditional Medicine Perspective Based on the Teaching of Avicenna. *Iran Red Crescent Med J*. 2012;**14**(3):133-8.

SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

کارگاه های آموزشی مرکز اطلاعات علمی



مقاله نویسی علوم انسانی
تربیه آموزشی

مقاله نویسی علوم انسانی



اصول تنظیم قراردادها
دوره آموزشی

اصول تنظیم قراردادها



آموزش مهارت های کاربردی در تدوین و چاپ مقاله
تربیه آموزشی

آموزش مهارت های کاربردی در تدوین و چاپ مقاله