

//
//

()

*

(salvage)

Archive of SID

()

(%)

(%)

%

(%)

Lactic Acid Dehydrogenase

(DHAP)

DHAP

(.)

%

%

()

(Performance Status)

(Metoclopramide)

(Kytril)

/ Dexamethasone+High-dose :DHAP)

(Cytosine Arabinoside+Cisplatin

% %

()

()

% (Ara-C)

in vitro

(remission)

DHAP

(.)

()

)

(

(Adriamycin)

CHOP

(Cyclophosphamide)

(Vincristine)

/

(restaging)

(hydration)

/ /

() (B- symptom) B

جدول ۱- مشخصات مبتلایان به لنفوم عود کرده تحت درمان با DHAP

B	
) LDH	
()	()
(Low grade)	()
(Intermediate grade)	()
(High grade)	()
()	()

Tumor Lysis)

(Syndrome

()	()	()	()	%
()	()	()	()	(partial remission or PR)
()	()	()	()	complete remission or)
()	()	()	()	(CR
()	()	()	()	" %
()	()	()	()	"(progressive disease or PD)
()	()	()	()	(PR) (PD)
()	()	()	()	(stable disease or SD)
()	()	()	()) chi-square
()	()	()	()	P value (Fisher's exact
()	()	()	()	% (response rate)
()	()	()	()	/(

.(CR)

(intermediate-grade)

(%)

%

DHAP		LDH	
		%	
		(CR)	
	LDH		(CR)
(B-symptoms)	%		
(remission)		(CR)	
		(CR)	
		(CR)	
LDH	(stage)	%	(stage) LDH
	()	(stage)	()
LDH	()	(stage)	
(stage)			
	LDH	()	جدول ۲ - فاکتورهای تعیین کننده پیش آگهی مبتلایان به لنفوم عود کرده تحت درمان با DHAP
(survival)			
		P-value	
		(CR)	
DHAP	(CR)		
			: (B symptoms)
	/	(%)	
		(%)	
			: LDH
	< /	(%)	
		(%)	/
		(%)	
			: (Stage)
	< /	(%)	I - II
		(%)	III - IV
			: LDH
		(CR)	
			LDH
			LDH
			LDH
		(remission)	

جدول ۳ - عوارض مربوط به رژیم درمانی DHAP در مبتلایان
به لنفوم عود کرده

DHAP		%	
LDH	()	()	:
(stage)	()	()	(<500/ mic L)
	(stage)	()	(Hb< 7gr/dL)
		()	(<20,000 / mic L)
		()	:(constitutional)
	LDH	()	
LDH	(CR)	()	
	(stage)	()	:
	(CR)	()	
		()	
	LDH	()	
	(stage)	()	:
		()	
%	(LDH)	()
			:
			()
			()
			()
()		
(CR)	DHAP		
CNS			
	DHAP		
		%	MIME
			:
			(CR)
			DHAP
DHAP			
		MIME	
			%
	(hypokalemia)	(CR)	
			%
			DHAP

(Phillip)

(salvage)

DHAP

()

%

%

%

(salvage)

% (disease-free survival)

DHAP

%

()

()

Kantajian

(stage)

LDH

(%)

Adelstein ()

(PR)

(consolidation)


)

(

(CR)

DHAP

()



REFERENCES

1. Editorial. Treatment for Non-Hodgkin's Lymphoma in relapse. What are alternatives? *N.Eng.J.Med* 1995; 333: 1565-6.
2. Phillip T, Guglielmi C, Hagenbeek A. Autologous bone marrow transplantation as compared with salvage chemotherapy in relapse of chemotherapy-sensitive Non-Hodgkin's lymphoma. *N Eng J Med*, 1995; 333: 1540-5.
3. Velasquez W.S, Cabanillas F, Salvador P. Effective salvage therapy for lymphoma with cisplatin in combination with high-dose Ara-C and dexamethasone (DHAP). *Blood* 1988; 71: 117-22.
4. Cabanillas F, Hagemester FB, Bodey GP. IMVP-16: an effective regimen for patients with lymphoma who have relapsed after initial combination chemotherapy. *Blood* 1982; 60: 693-7.
5. Kantajian H, Barlogie B, Plunkett W, Velasquez WS, McLaughlin P, Riggs S, Cabanillas F. High dose cytosine arabinoside in Non-Hodgkin's lymphoma. *J Clin Oncol* 1983; 1: 689.
6. Adelstein DJ, Lazarus HM, Hines JD, Herzig RH. High-dose cytosine arabinoside in previously treated patients with poor-prognosis Non-Hodgkin's lymphoma. *Cancer* 1985; 56: 1493.
7. Velasquez WS, Jagannath S, Hagemester FB, McLaughlin P, Swan F, Redman JR, et al. Dexamethasone, high dose Ara-C and cisplatin (DHAP) as salvage treatment for relapsing Hodgkin's disease. *Proc Am Soc Hematol* 1986; 68: 242a (abstr).
8. Velasquez WS, McLaughlin P, Swan F, Kantajian J, Cabanillas F, Barlogie B. Dexamethasone, high dose Ara-C and cisplatin (DHAP) combination for progressive chronic lymphocytic leukemia. *Proc Am Soc Hematol* 1986; 68: 234a (abstr).
9. Spitzer G, Armitage J, Jagannath S, Dicke K, Zander A, Kumar P, et al. Treatment of relapsed diffuse large cell lymphoma (DLCL) with high dose Melphalan (HDM), total body irradiation (TBI) and autologous bone marrow transplantation (ABMT). *Eur J Cancer Clin Oncol* 1986; 22: 677.
10. Anderson CC, Goldstone AH, Souhami RL, Linch DC, Harper PG, McLennan KA, et al. Very high dose chemotherapy with autologous bone marrow rescue in adult patients with resistant relapsed lymphoma. *Cancer Chemother Pharmacol* 1986; 16: 170.
11. Phillip T, Armitage JO, Spitzer G, Chauvin F, Jagannath F, Cahn JY, et al. High-dose therapy and autologous bone marrow transplantation after failure of conventional chemotherapy in adults with intermediate-grade or high-grade Non-Hodgkin's lymphoma. *N Eng J Med* 1987; 316: 1493.