

# SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

## کارگاه های آموزشی مرکز اطلاعات علمی



مقاله نویسی علوم انسانی



اصول تنظیم قراردادها



آموزش مهارت های کاربردی در تدوین و چاپ مقاله



مرکز بررسی و مطالعات دریایی

سازمان بنادر و دریانوردی به عنوان تنها مرجع حاکمیتی کشور در امور بندری، دریایی و کشتی‌رانی بازرگانی به منظور ایفای نقش مرجعیت دانشی خود و در راستای تحقق راهبردهای کلان نقشه جامع علمی کشور مبنی بر "حمایت از توسعه شبکه‌های تحقیقاتی و تسهیل انتقال و انتشار دانش و سامان‌دهی علمی" از طریق "استانداردسازی و اصلاح فرایندهای تولید، ثبت، داوری و سنجش و ایجاد بانک‌های اطلاعاتی یکپارچه برای نشریات، اختراعات و اکتشافات پژوهشگران"، اقدام به ارایه این اثر در سایت SID می‌نماید.



سازمان بنادر و دریانوردی



**The 10th International Conference on Coasts, Ports and Marine Structures (ICOPMAS 2012)  
Tehran, Iran, 19-21 Nov. 2012**



## **SIMULATION OF WAVE CHARACTERISTICS IN CASPIAN SEA USING SWAN**

**Mahsa Modiri<sup>1</sup>, Amir Etemed Shahidi<sup>2</sup>, Sanaz Hadadpour<sup>3</sup>**

**Key Words:** wave hindcasting, SWAN, Caspian sea, QuikSCAT wind field

### **Abstract**

Wave parameters are necessary for many marine and coastal operations. Different methods such as empirical, numerical and soft computing have been developed for wave hindcasting. In this study, SWAN model has been used for the prediction of wave parameters in Caspian Sea. The comparison of obtained results with the Anzali buoy measurements, indicated that this global modeling yields an accurate estimation of wave characteristics.

### **Introduction**

Wave characteristics are one of the most important factors in design of coastal and marine structures. For the prediction of wave parameters, different methods have been developed such as empirical methods (e.g. SPM [1], SMB [2], CEM [3] and Donelan [4]), numerical models (e.g. Mike 21 [5], Wavewatch III [6], WAM [7] and SWAN [8]) and soft computing (e.g. Artificial Neural Network, Fuzzy Inference Systems, Decision Trees and Genetic Programming). Until now different numerical models have been investigated for wave hindcasting ([9], [10], [11], [12], [13] and [14]). In this study SWAN is used for wave modeling in Caspian Sea.

### **Study Area and Evaluation of the Data**

In this study, QuikSCAT wind data have a spatial resolution of 25 kilometer and temporal resolution of 12 hour, were used. The recorded wave data by Anzali buoy at 49.52° E and 37.549° N were used for calibration and verification of the model. The study area, Caspian Sea, has a laterally-prolonged scale of about 800 kilometer in the west-east direction between 47° E and 55° E. Its width is about 1000 kilometer in the north-south direction between 36.5° N and 47° N. The geographical domain has a resolution of 0.0166° E in both x and y directions. Figure 1 shows the considered computational area in the Caspian Sea.

---

<sup>1</sup>graduate student, iran university of science and technology, modiri@civileng.iust.ac.ir

<sup>2</sup>associated professor, Griffith University, etemad@iust.ac.ir

<sup>3</sup>graduate student, iran university of science and technology, s\_hadadpour@civileng.iust.ac.ir




Fig. 1) Computational area in the Caspian Sea

### **SWAN Model**

SWAN is a third generation spectral wind wave model solves the spectral action balance equation for the cartesian coordinates:

$$\frac{\partial}{\partial t} N + \frac{\partial}{\partial x} c_x N + \frac{\partial}{\partial y} c_y N + \frac{\partial}{\partial \sigma} c_\sigma N + \frac{\partial}{\partial \theta} c_\theta N = \frac{S}{\sigma} \quad (1)$$

Where  $N$  is action density,  $c_x$  and  $c_y$  is propagation velocities in  $x$  and  $y$  spaces, respectively.  $c_\theta$  and  $c_\sigma$  is propagation velocities in  $\theta$  and  $\sigma$  spaces, respectively. The term  $S$  on the right-hand side of the equation is the source term in terms of energy density representing the effects of generation, dissipation and nonlinear wave-wave interactions.

### **Select the Calibration And Verification Periods**

For model calibration and verification time series from May, 23th to May, 27th, 2003 and June, 15th to June, 21th, 2003 were selected, respectively. According to the time series of wind and wave data for those periods, there are a good agreement between the variation of wind speed and wave parameters for both calibration and verification periods. Statistics of the wind and wave characteristics for these period are illustrated in tables 1 and 2.

Table 1) Statistical of wind and wave characteristics for calibration period


Parameter	Minimum	Average	Maximum
Significant wave height (m)	0.20	0.45	1.18
Peak period (s)	1.89	4.01	5.88
Wind speed (m/s)	0	3.89	12
Wind direction (degree)	13.33	214.81	350.36

**Table 2) Statistical of wind and wave characteristics for verification period**


Parameter	Minimum	Average	Maximum
Significant wave height (m)	0.18	0.41	1.56
Peak period (s)	2	3.85	8.33
Wind speed (m/s)	0.04	3.55	17.24
Wind direction (degree)	0.20	206.73	358.68

**Calibration of the Model**

Time series of the measured and modeled wave characteristics during the calibration period are shown in figures 2 and 3.



**Fig. 2) Time series of the measured and modeled significant wave heights, calibration period**



**Fig. 3) Time series of the measured and modeled peak period, calibration period**

In order to evaluate the results quantitatively, the Bias, correlation coefficient (r), the root mean square error (RMSE) and the scatter index (SI) are calculated:

$$Bias = \bar{p} - \bar{o} \tag{2}$$



$$R = \frac{\sum_{i=1}^N [(o_i - \bar{o}) \times (p_i - \bar{p})]}{\sqrt{\sum_{i=1}^N (o_i - \bar{o})^2 \sum_{i=1}^N (p_i - \bar{p})^2}} \quad (3)$$

$$RMSE = \sqrt{\frac{\sum_{i=1}^N (o_i - p_i)^2}{N}} \quad (4)$$

$$SI = \frac{RMSE}{\bar{o}} \quad (5)$$


where,  $o_i$  and  $p_i$  denote the observed and predicted value, respectively. Moreover,  $\bar{o}$  and  $\bar{p}$  denote the mean value of them.  $N$  shows the number of data.

**Table 3) Error indices for calibration period**

Error index	Peak period	Significant wave height
Bias	-0.020	0.123
R	0.374	0.917
RMSE	1.209	0.272
SI (%)	30.18	60

### Verification of the Model

The results of verification process are indicated in figures 4 and 5 and table 4.



**Fig. 4) Time series of the measured and modeled significant wave heights, verification period**




Fig. 5) Time series of the measured and modeled peak period, verification period

Table 4) Error indices for verification period

Error index	Peak period	Significant wave height
Bias	-0.422	0.002
R	0.534	0.903
RMSE	1.487	0.247
SI (%)	38.66	60.53

### Summary and Conclusion

In this study SWAN is used for wave modeling in Caspian Sea. The results show that the modeled wave characteristics, were correlated well with observed data in the verification period. Moreover, the statistics of the error show that the model performs well in simulating wave parameters.

### References

- [1]-Shore Protection Manual, (1984), 4th ed., 2Vol, U.S. Army Engineer Waterways Experiment Station, U.S. Government Printing Office, Washington, DC.
- [2]-Sverdrup, H.U., Munk, W.H., (1974), Wind, seas and swell: theory of relations for forecasting, Publication 601, U.S. Navy Hydrographic office, Washington, DC.
- [3]-Coastal Engineering Manual, (2006), Chapter II-2, Meteorology and Wave Climate, Engineer Manual 1110-2-1100, U.S. Army Corps of Engineers, Washington, DC.
- [4]-Donelan, M.A., (1980), Similarity theory applied to the forecasting of wave heights, periods and directions, In: Proceedings of Canadian Coastal Conference, National Research Council of Canada, 47–61.
- [5]-MIKE 21, (2009), Wave Modeling Manual, DHI Water and Environment.
- [6]-Tolman, H.L., (1991), A third-generation model for wind waves on slowly varying, unsteady and inhomogeneous depths and currents, *Physical Oceanography*, vol. 21, 782–797.
- [7]-Komen, G.J., Cavaleri, L., Donelan, M., Hasselmann, K., Hasselmann, S., Janssen, P.A.E.M., (1994), *Dynamics and Modeling of Ocean Waves*, Cambridge University Press, UK.
- [8]-Booij, N., Ris, R.C., Holthuijsen, L.H., (1999), A third generation wave model for coastal regions. 1. Model description and verification, *Geophysical Research*, vol. 104, 7649-7666.
- [9]-Moeini, M.H., Etemad-Shahidi, A., Chegini, V., (2010), Wave modeling and extreme value analysis off the northern coast of the Persian Gulf, *Applied Ocean Research*, vol. 32, 209-218.

- [10]-Moeini, M.H., Etemad-Shahidi, A., (2007), Application of two numerical models for wave hindcasting in Lake Erie, *Applied Ocean Research*, vol. 29, 137-145.
- [11]-Cavaleri, L., Bertotti, L., (2006), The improvement of modelled wind and wave fields with increasing resolution, *Ocean Engineering*, vol. 33, 553-565.
- [12]-Bolaños-Sanchez, R., Sanchez-Arcilla, A., Cateura, J., (2007), Evaluation of two atmospheric models for wind-wave modelling in the NW Mediterranean, *Marine Systems*, vol. 65, 336-353.
- [13]-Signell, R.P., Carniel, S., Cavaleri, L., Chiggiato, J., Doyle, J., Pullen, J., et al., (2005), Assessment of wind quality for oceanographic modelling in semi-enclosed basins, *Marine Systems*, vol. 53, 217-233.
- [14]-Kazeminezhad, M.H., Etemad-Shahidi, A., Mousavi, J., (2005), Application of fuzzy inference system in prediction of wave parameters, *Ocean Engineering*, vol. 32, 1709-1725.





# SID



سرویس های ویژه



سرویس ترجمه تخصصی



کارگاه های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم های آموزشی

## کارگاه های آموزشی مرکز اطلاعات علمی



مقاله نویسی علوم انسانی



اصول تنظیم قراردادها



آموزش مهارت های کاربردی در تدوین و چاپ مقاله