





















- [23]- Pedrol N, Ramos P, Reigosa MJ (2000). Phenotypic plasticity and acclimation to water deficits in velvet-grass: a long-term greenhouse experiment. Changes in leaf morphology, photosynthesis and stress-induced metabolites. *J. Plant Physiol.*, 157: 383-393.
- [24]- Moran, J.F., Becana, M., Ormaetxe, I.I., Frechilla, S.L., Klucasc, R.V., and Tejo, D.A. 1994. Drought induces oxidative stress in pea plants. *Plant*, 194: 346-352.
- [25]-Martinez, C. A., C. Guerrero and U. Moreno. 2005. Diurnal fluctuations of carbon exchange rate, proline content and osmotic potential in two water - stressed potato hybrids. *Revista Brasileira de Fisiologia Vegetal*. 7: 27- 33.
- [26]- Arazmjo, A., M. Heidari and A. Ghorbani. 2010. The effect of water stress and three sources of fertilizers on flower yield, physiological parameters and nutrient uptake in chamomile (*Matricaria chamomilla* L.). *Iranian Journal of Medicinal and Aromatic Plants*. 25(4):482-494
- [27]- Hernandez, J.A., Jimenez, A., Mullineaux, P., Sevilla, F. 2000. Tolerance of pea (*Pisum sativum* L.) to long term salt stress is associated with induction of antioxidant defences. – *Plant Cell Environ*. 23: 853-862.
- [28]- Zhang J.X., Kirkham, M.B.. Droughtstress- induced changes in activities of superoxide dismutase, catalase, and peroxidase in wheat species. *Plant Cell Physiol*. 35: 785–791,1994.
- [29]-Bowler, C., Van Montagu, M., Inze, D., 1992. Superoxide dismutase and stress tolerance. *Ann. Rev. Plant Physiol. Mol. Biol*. 43: 83-116. 1161-1168.