

SID



ابزارهای پژوهش



سرویس ترجمه تخصصی



کارگاه‌های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم‌های آموزشی

سامانه ویراستاری (ویرایش متون فارسی، انگلیسی، عربی)

کارگاه‌ها و فیلم‌های آموزشی مرکز اطلاعات علمی



روش تحقیق کمی

روش تحقیق کمی



آموزش مهارت‌های کاربردی در تدوین و چاپ مقالات ISI

آموزش مهارت‌های کاربردی در تدوین و چاپ مقالات ISI



آموزش نرم افزار Word برای پژوهشگران

آموزش نرم افزار Word برای پژوهشگران

Association between Hepatitis G and Unknown Chronic Hepatitis

Saeed Soleiman-Meigooni¹, Ali Asgari¹, Seyyed Javad Hoseini-Shokouh¹, Jalil Rajabi¹,
Mohammad Hassan Kazemi-Galougahi², Mohammad Moshtaghi³

¹Infectious disease specialist, Infectious Disease Research Center, AJA University of Medical Sciences, Tehran, Iran

²Ph.D. Candidate of Epidemiology, Department of Epidemiology & Biostatistics, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

³M.D, General Physician, Tehran, Iran

Type of article: Original

Abstract

Background: Hepatitis G virus (HGV) is a hepatotrope virus with unknown importance. The genome of the virus has been detected in patients with acute or chronic non-A-E hepatitis, cirrhosis, and hepatocellular carcinoma. The aim of this study was to determine the association between hepatitis G and unknown chronic hepatitis.

Methods: This case-control study was performed in Ebne-Sina military hospital in Hamadan, Iran. The cases were 35 military staff with unknown chronic hepatitis. The control group consisted of 59 healthy subjects who had normal levels of serum alanine aminotransferase (ALT). The data were analyzed by SPSS, version 18, using Fisher's exact test, the Student's t-test, and multivariate logistic regression analysis.

Results: Only one patient in the case group (2.9%) tested positive for HGV antibodies, and no one was infected in the control group. There was no association between HGV infection and unknown chronic hepatitis in our study ($P=0.37$). A significant association was found between the male gender and unknown chronic hepatitis ($OR=14.9$, $P=0.01$).

Conclusion: No association between HGV infection and unknown chronic hepatitis was found in our study, so it was not necessary to evaluate these patients for HGV infection.

Keywords: hepatitis G, chronic hepatitis

1. Introduction*1.1. Background*

The asymptomatic elevation of liver enzymes is a common problem, and, if it goes undetected, it can damage the liver over time (1). In a survey of 1959 blood donors in Iran, 5.1% had asymptomatic elevated liver enzymes, and the common diagnoses for half of those affected were nonalcoholic steatohepatitis (NASH) (88%), hepatitis C (7.7%), and alcohol and drug-related liver injury (1.9%) (2). Another study of about 2000 people in Golestan Province in northern Iran found that the prevalence of persistently-elevated liver enzymes was 3.1% with an unknown etiology in about 80%, hepatitis B in 9.3%, hepatitis C in 6.2%, alcoholic liver disease in 4.6%, and fatty liver disease in 2% (3).

1.2. Statement of the problem

HGV is an RNA flavivirus transmitted mainly by the parenteral route. HGV RNA and HGV antibody (anti-E2) seropositivity in blood donors is about 1-4% and 3-14%, respectively (4). The mean prevalence of HGV in blood donors is 4.8% and is different in diverse world regions, e.g., 4.5% in Caucasians, 3.4% in Asians, and 17.2% in Negros (5). It seems that Iran has the least prevalence of HGV, i.e., about 1%, among other countries in the Middle

Corresponding author:

Assistant Professor Dr. Saeed Soleiman-Meigooni, Infectious Disease Research Center, AJA University of Medical Sciences, Tehran, Iran. Tel: +98.9123058830, Fax: +98.2155725615, Email: Dr.saeed.meigooni@gmail.com

Received: March 03, 2014, Accepted: January 20, 2015, Published: March 01, 2015

© 2015 The Authors. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

East, e.g., Turkey (4.1%), Kuwait (24.6%), Jordan (9.8%), and Saudi Arabia (2%) (7-10). Seventy-five percent of infected patients have normal liver enzymes with no symptoms of hepatic disease; however, it can cause acute and chronic hepatitis (11). In a study of 41 acute and 67 chronic hepatitis patients with unknown etiology in China, HGV RNA was detected in six patients (14.6%) and 12 patients (17.9%), respectively (12). A similar study of 55 acute unknown hepatitis patients indicated that 29.1% had HGV RNA seropositivity and very severe clinical syndrome (13). Multiple studies have expressed doubts about the relationship between HGV infection and hepatic damage (14-16). Thus, our aim was to investigate the association between HGV infection and unknown chronic hepatitis.

1.3. Objective of the research

The aim of this research was to evaluate the association of hepatitis G with unknown chronic hepatitis. The specific objectives of the study were:

- 1) To determine the frequency of HGV antibodies in the case group and the control group
- 2) To determine the difference in the HGV seropositivity between the two groups

2. Materials and Methods

2.1. Research design and setting

A case-control study was conducted at the Ebne-Sina Army Hospital in Hamadan, Iran, from September 2011 through October 2012. Patients with unknown chronic hepatitis were evaluated for HGV infection. The cases were 35 military staff with unknown chronic hepatitis, and the controls were 59 patients with normal serum ALT levels who were selected from the orthopedic and gynecologic clinics at the same hospital. Chronic hepatitis was defined as serum ALT \geq 40 IU/L for more than six months (17).

2.2. Data collection

A checklist containing demographic data, ALT levels, and HGV antibody states was designed to collect the data (Table 1). One hundred and seventy-three subjects with serum ALT \geq 40 IU/L were evaluated for:

- 1) Viral hepatitis B and C by the detection of HB antigens and HCV antibodies using the ELISA method with 5-ml samples of serum.
- 2) Non-alcoholic fatty liver disease (NAFLD) using abdominal sonography by an expert radiologist.
- 3) History of alcohol or hepatotoxic drug consumption or recent infectious processes, such as cold, fever, sore throat, myalgia, arthralgia, diarrhea, abdominal pain, and vomiting (2, 3, 17).

Any patients who had any of the three conditions described above were excluded from the study. In this way, 115 subjects were selected for the study, and they were followed for 6-9 months without any intervention, and 35 patients with permanent ALT \geq 40 IU/L were selected as the case group. The control group was comprised of patients with normal serum ALT levels who were referred to the orthopedic and gynecologic clinics at the same hospital. Five-milliliter serum samples were taken from the members of both the case and control groups and sent to a private laboratory to be assessed for HGV infection. The laboratory used the Cortez 4th generation kit (manufactured in the U.S.), and HGV anti-E2 was detected by ELISA with 99.8% sensitivity.

2.3. Ethical considerations

This research was approved by the Ethics Committee at the AJA University of Medical Sciences. Participation in the study was voluntary, and the physicians explained the purpose and procedures of the study to the participants and asked each one to sign a consent form. For the sake of confidentiality and privacy, the serum samples were coded and sent anonymously to the laboratory.

2.4. Statistical analysis

SPSS version 18 (SPSS Inc., Chicago, IL, U.S.) was used to analyze the data. Fisher's exact test, the Student's t-test, and multivariate logistic regression were used for data analysis. An alpha level of 0.05 was considered to be significant.

3. Results

The mean ages of the members of the case group and the control group were 34.9 \pm 8.5 and 40 \pm 12.3 years, respectively. All of patients in the case group were males, and the male-to-female ratio in the control group was 2.28. There was a significant difference between the ages of the patients in the case and control groups, with the former having a significantly lower mean age than the latter based on the Student's t-test (P=0.021) (Table 1).

Among all of the people in the case group and the control group, there was just one person who tested HGV anti-E2 positive in the case group (2.9%); all of the people in the control group tested negative. Thus, Fisher's exact test indicated that there was no significant statistical difference of HGV infection between case group and the control group ($P=0.372$). In multivariate logistic regression analysis, a significant relationship was found between unknown chronic hepatitis and being male ($OR=14.9$; 95% $CI=1.9-117.6$, $P=0.010$), but there was no such relationship for age ($P=0.446$) or HGV infection ($P=0.269$) (Table 2).

Table 1. Reasons for excluding patients with elevated liver transaminases (n = 58)

Reason for exclusion	n	%
Fatty liver syndrome	38	65.5
History of recent infectious processes or symptoms/signs of ongoing infectious processes	17	29.3
HBV infection	2	3.4
Long term methotrexate consumption	1	1.7
Total	58	100

Table 2. Demographic characteristics of the study population (n = 94)

Variables		Cases		Controls		P-Value
		n	%	n	%	
Age group (year)	<30	11	31.4	14	23.7	0.021
	30-39	15	42.8	14	23.7	
	40-49	7	20.0	19	32.2	
	50-59	2	5.7	8	13.6	
	≥60	0	0	4	6.8	
Gender	Male	34	97.1	41	69.5	0.001
	Female	1	2.9	18	30.5	

4. Discussion

In our study, the frequency of HGV antibodies was not different between the patients with unknown chronic hepatitis and the healthy subjects, confirming that the HGV infection did not injure the liver. Our results were in agreement with the results of a similar study conducted in Italy on 50 unknown chronic hepatitis patients and 50 healthy persons (18). Some studies have concluded that it was doubtful that HGV was responsible for chronic hepatitis (19, 20), and several studies have shown the correlation of HGV with hematologic disorders, such as non-Hodgkin's lymphoma (21-23), aplastic anemia (24, 25) and chronic renal failure (26-29). In our study, chronic hepatitis was more frequent in males due to the selection of the cases from active military patients participating in a military health monitoring program.

The main challenges associated with HGV infection are liver injury and chronic hepatitis. It seems that co-infection with other viral forms of hepatitis, such as HBV and HCV, can lead to hepatic damage whereas the HGV infection alone does not. One survey of 64 Japanese children infected with HCV with co-infection with HGV showed no clinical, virological, or pathological differences between those with HCV/HGV co-infection and HCV alone (30). In another study of 130 patients with chronic hepatitis B and 173 patients with chronic hepatitis C, the prevalence of HGV RNA was about 8% and 17%, respectively, and, while HGV co-infection did not worsen the clinical symptoms of disease, it accelerated the progression to chronic hepatitis and increased the risk of hepatocellular carcinoma (31). Regarding the limitation of the study, we did not study patients with unknown chronic hepatitis for auto-immune hepatitis, which might have a role in liver inflammation and chronic hepatitis in these patients.

5. Conclusion

This study showed no association between a history of HGV infection and chronic hepatitis in the patients with unknown chronic hepatitis. Therefore, it seems that screening for HGV antibodies is not recommended. However, we believe that HGV RNA measurement in the patients with unknown chronic hepatitis may help identify any ongoing inflammation. Further studies are recommended to evaluate HGV RNA seropositivity in patients with unknown chronic hepatitis.

Acknowledgments

We thank the Day Private Laboratory in Hamadan, Iran, for collecting and analyzing the serum samples for HGV antibodies. The authors do not have any conflict of interest, and corresponding author self-funded this research.

Conflict of Interest:

There is no conflict of interest to be declared.

Authors' contributions:

All of authors contributed to this project and article equally. All authors read and approved the final manuscript.

References

- 1) Pratt DS, Kaplan MM. Evaluation of abnormal liver-enzyme results in asymptomatic patients. *N Engl J Med* 2000; 342:1266-1271. April 27, 2000. DOI: 10.1056/NEJM200004273421707. PMID:10781624
- 2) Pourshams, A., Malekzadeh, R., Monavvari, A., Akbari, M. R., Mohamadkhani, A., Yarahmadi, S, et al. Prevalence and etiology of persistently elevated alanine aminotransferase levels in healthy Iranian blood donors. *Journal of Gastroenterology and Hepatology*, 20: 229–233. DOI: 10.1111/j.1440-1746.2004.03511.x. PMID:15683425
- 3) Jamali R, Khonsari M, Merat S, Khoshnia M, Jafari E, Bahram Kalhori A, et al. Persistent alanine aminotransferase elevation among the general Iranian population: prevalence and causes. *World J Gastroenterol*. 2008; 14(18):2867-71. DOI:10.3748/wjg.14.2867. PMID:18473412
- 4) Kleinman S. Hepatitis G virus biology, epidemiology, and clinical manifestations: Implications for blood safety. *Transfus Med Rev*, 2001; 15(3):201-12. DOI:10.1053/tm.2001.24589. PMID:11471122
- 5) Wiwanitkit V. Hepatitis G virus RNA positivity among the voluntary blood donors: a summary. *Ann Hepatol*. 2005; 4(1):43-6. PMID:15798660
- 6) Kafi-Abad SA, Samiei Sh, Talebian A, Maghsudloo M, Gharehbaghian A. Hepatitis G virus infection in Iranian blood donors and high-risk groups. *Hepat Mon*. 2009; 9(4):282-86.
- 7) Ramezani A, Gachkar L, Eslamifar A, Khoshbaten M, Jalilvand S, Adibi L, et al. Detection of hepatitis G virus envelope protein E2 antibody in blood donors. *Int J Infect Dis*. 2008; 12(1):57-61. DOI: /10.1016/j.ijid.2007.04.010. PMID:17629534
- 8) Kalkan A, Ozdarendeli A, Bulut Y, Saral Y, Ozden M, Kelestimur N, et al. Prevalence and genotypic distribution of hepatitis GB-C/HG and TT viruses in blood donors, mentally retarded children and four groups of patients in eastern Anatolia, Turkey. *Jpn J Infect Dis*. 2005; 58(4):222-7. PMID:16116255
- 9) Odeh RA, Yasin S, Nasrallah G, Babi Y. Rates of infection and phylogenetic analysis of GB virus-C among Kuwaiti and Jordanian blood donors. *Intervirol*. 2010; 53(6):402-7. DOI: 10.1159/000317290. PMID:20606462
- 10) Al-Ahdal MN, Rezeig MA, Kessie G, Chaudhry F, Al-Shammary FJ. GB virus C/hepatitis G virus infection in Saudi Arabian blood donors and patients with cryptogenic hepatitis. *Arch Virol*. 2000; 145(1):73-84. DOI: 10.1007/s007050050006. PMID:10664407
- 11) Alter HJ. The cloning and clinical implications of HGV and HGBV-C. *The N Engl J Med*, 1996; 334(23):1536-7. DOI:10.1056/NEJM199606063342310. PMID:8618611
- 12) Wang HL, Jin DY. Prevalence and genotype of hepatitis G virus in Chinese professional blood donors and hepatitis patients. *J Infect Dis*. 1997; 175(5):1229-33. DOI: 10.1086/593676
- 13) Tassopoulos NC, Papatheodoridis GV, Delladetsima I, Hatzakis A. Clinicopathological features and natural history of acute sporadic non-(A-E) hepatitis. *J Gastroentrol Hepatol*, 2008 Aug;23(8 Pt 1):1208-15. doi: 10.1111/j.1440-1746.2008.05454.x. Epub 2008 Jun 28. PMID:18554239
- 14) Uygun A, Kadayifci A, Kubar A, Tuzun A, Erdil A, Gulsen M, et al. Insignificant role of hepatitis G virus infection in patients with liver enzyme elevations of unknown etiology *J Clin Gastroenterol*. 2000 Jul;31(1):73-6. DOI: 10.1097/00004836-200007000-00017. PMID:10914782
- 15) Hyams KC. Chronic liver disease among U.S. military patients: the role of hepatitis C and G virus infection. *Mil Med*. 2000; 165(3):178-9. PMID:10741077
- 16) Jeon MJ, Shin JH, Suh SP, Lim YC, Ryang DW. TT virus and hepatitis G virus infections in Korean blood donors and patients with chronic liver disease. *World J Gastroenterol*. 2003; 9(4):741-4. PMID:12679923
- 17) Mohamadnejad M, Pourshams A, Malekzadeh R, Mohamadkhani A, Rajabiani A, Asgari AA, et al. Healthy ranges of serum alanine aminotransferase levels in Iranian blood donors. *World J Gastroenterol*. 2003; 9(10):2322-4. PMID:14562401

- 18) Grassi M, Raffa S, Traditi F, Ferrazzi M, Cioschi S, Fontana M, et al. Detection and clinical evaluation of GBV-C/HGV in plasma from patients with chronic hepatitis of unknown etiology. *Clin Ter.* 2000 Jul-Aug; 151(4):241-5. PMID:11107672
- 19) Hollingsworth RC, Minton EJ, Fraser-Moodie C, Metivier E, Rizzi PM, Irving WL, et al. Hepatitis G infection: role in cryptogenic chronic liver disease and primary liver cell cancer in the UK. Trent Hepatitis C virus Study Group. *Journal J Viral Hepat.* 1998 May;5(3):165-9. DOI: 10.1046/j.1365-2893.1998.00102.x. PMID:9658369
- 20) Guilera M, Saiz JC, Lopez-Labrador FX, Olmedo E, Ampurdanes S, Fornis X, et al. Hepatitis G virus infection in chronic liver disease. *Gut.* 1998; 42(1):107-11. DOI: 10.1136/gut.42.1.107. PMID: 9505895
- 21) A, Chow R, et al. GBV-C/hepatitis G virus infection and non-Hodgkin lymphoma: a case control study. *Int J Cancer.* 2010 Jun 15;126(12):2885-92. doi: 10.1002/ijc.25035. PMID: 19904755
- 22) Giannoulis E, Economopoulos T, Mandraveli K, Giannoulis K, Nikolaides C, Zervou E, et al. The prevalence of hepatitis C and hepatitis G virus infection in patients with B cell non-Hodgkin lymphomas in Greece: a Hellenic Cooperative Oncology Group Study. *Acta Haematol.* . 2004; 112(4):189-93. DOI: 10.1159/000081270. PMID:15564729
- 23) Cividari G, Tanzi E, Ferrari B, Vallisa D, Zanetti A, Cavanna L. High prevalence of anti-HGV/E2 antibodies in HCV-positive patients with non Hodgkin's lymphoma. *Haematologica.* 1998; 83(10):957-8. PMID:9830812
- 24) Riaz Shah SA, Idrees M, Hussain A. Hepatitis G virus associated aplastic anemia: a recent case from Pakistan. *Virology.* 2011 Jan 21; 8:30. DOI: 10.1186/1743-422X-8-30. PMID:21251303 PMCID:PMC3032715
- 25) Zaidi Y, Chapman CS, Myint S. Aplastic anaemia after HGV infection. *Lancet.* 1996; 348(9025):471-2. DOI: 10.1016/S0140-6736(05)64561-8. PMID: 8709794
- 26) Masuko K, Mitsui T, Iwano K, Yamazaki C, Okuda K, Meguro T, et al. Infection with hepatitis GB virus C in patients on maintenance hemodialysis. *N Engl J Med.* 1996 Jun 6; 334(23):1485-90.. DOI: 10.1056/NEJM199606063342301. PMID:8618602
- 27) Fallahian F, Alavian SM, Rasoulinejad M. Epidemiology and transmission of hepatitis G virus infection in dialysis patients. *Saudi J Kidney Dis Transpl.* 2010 Sep; 21(5):831-4., PMID: 20814115
- 28) Hammad AM, Zaghoul MH. Hepatitis G virus infection in Egyptian children with chronic renal failure (single centre study). *Ann Clin Microbiol Antimicrob.* 2009 Dec 16; 8:36. DOI: 10.1186/1476-0711-8-36. PMID: 20015406.
- 29) Hinrichsen H, Leimenstoll G, Stegen G, Schrader H, Folsch UR, Schmidt WE. Prevalence of and risk factors for hepatitis G (HGV) infection in haemodialysis patients: a multicentre study. *Nephrology, dialysis, transplantation: Nephrol Dial Transplant.* 2002 Feb; 17(2):271-5. DOI: 10.1093/ndt/17.2.271. PMID:11812878
- 30) Komatsu H, Fujisawa T, Inui A, Sogo T, Morinishi Y, Miyagawa Y, et al. GBV-C/HGV infection in children with chronic hepatitis C. *J Med Virol.* 1999 Oct; 59(2):154-9. DOI: 10.1002. PMID:10459149
- 31) Yang JF, Dai CY, Chuang WL, Lin WY, Lin ZY, Chen SC, et al. Prevalence and clinical significance of HGV/GBV-C infection in patients with chronic hepatitis B or C. *Jpn J Infect Dis.* 2006 Feb;59(1):25-30.. PMID:16495630

SID



ابزارهای پژوهش



سرویس ترجمه تخصصی



کارگاه‌های آموزشی



بلاگ مرکز اطلاعات علمی



سامانه ویراستاری STES



فیلم‌های آموزشی

سامانه ویراستاری (ویرایش متون فارسی، انگلیسی، عربی)

کارگاه‌ها و فیلم‌های آموزشی مرکز اطلاعات علمی



روش تحقیق کمی

روش تحقیق کمی



آموزش مهارت‌های کاربردی در تدوین و چاپ مقالات ISI

آموزش مهارت‌های کاربردی در تدوین و چاپ مقالات ISI



آموزش نرم افزار Word برای پژوهشگران

آموزش نرم افزار Word برای پژوهشگران